

Application of Ovsynch-CIDR vs 2 Consecutive Injections of PGF 2 α , 14 Days Apart in Dairy Holstein Cows and Comparison of Reproductive Parameters, Plasma and Milk E2 and P4 Concentrations

A.H. Fallah Rad and G. Ajam

Department of Clinical Sciences, Faculty of Veterinary Medicine,
Ferdowsi University of Mashhad, Mashhad, Iran

Abstract: In the present study, effects of ovsynch-CIDR protocol was compared with the conventional method of 2 consecutive injections of PGF2 α 14 days apart, on some reproductive indices and concentrations of E2 and P4 on day 1, day 5 and 21 after insemination. All the cows assigned into the study were in their second parity onward and had no history of peri-parturient diseases including: retained placenta, dystocia, lameness, clinical mastitis and metritis. In the nPG group (n = 27), on day 40 \pm 5 postpartum, 2 doses of PG were injected 14 days apart and if after the 2nd injection, cow was inseminated upon showing signs of estrus. In the OC group (n = 25), Ovsynch-CIDR protocol along with timed AI was applied. Milk and blood samples were obtained from all the cows on day1, day 5 and 21 post AI. E2 and P4 of the plasma and P4 of the milk samples were measured by ELISA and reproductive indices were calculated. In case of return to estrus, the program was repeated for up to 3 times. Conception Rate (CR) was lower in the nPG than OC but the difference was not significant (p>0.05). In the 2nd and 3rd AI, CR was higher in OC than nPG (71.42 and 85.71% Vs 23.8 and 61.9% in the OC and nPG, respectively, p<0.05). Number of AI/conception and overall CR were 3.13, 40.3, 2.23 and 55.71% for nPG and OC, respectively (p<0.05). Days Open (DO) and predicted calving intervals were 138 and 418 and 106 and 387 for nPG and OC, respectively (p<0.05). Milk P4 was higher on day 5 and 21 in the OC and plasma E2 was lower in OC on day 5 than nPG (p<0.05). Results show that, ovulation synchronization by Ovsynch-CIDR, as compared with 2 consecutive PG injections reduced DO, increased CR in the 2nd and 3rd inseminations, increased overall CR and reduced number of inseminations per conception. It can not be claimed that results of this experiment might be similar in all the cows of the same herd or other herds because, cows assigned into this study were selected.

Key words: Ovsynch-CIDR, PGF2 α , P4, E2, dairy holstein

INTRODUCTION

Milk production of the dairy breeds had increase tremendously at the expense of decrease in reproductive performance (López-Gatius *et al.*, 2001), increase on the incidence of reproductive and metabolic diseases (Francos, 1998; Macmillan *et al.*, 1996). From 1973-1995, First Service Conception Rate (FSCR) in the dairy cows has been declining at a rate of 0.4 and 1% per year in the US and UK, respectively (Gröhn and Rajala-Schultz, 2000). The main reasons are, negative impacts of poor heat detection, early embryonic death, poor management and nutrition (Butler, 2001; Peters, 1996). In industrial dairy farming, heat detection rarely reaches a 50% rate. Moreover, after induction of estrus by using different hormone therapy regimens, CR is around 30-45% that is much lower than 60-70% of the natural conditions (López-Gatius, 2000).

With regard to the expansion of the dairy farms, difficulty of heat detection and also greater use of AI, estrus and ovulation synchronization methods are increasingly applied (Burke *et al.*, 1999; Pursley *et al.*, 1995; Xu and Burton, 2000). Application of some of these techniques along with fixed time AI (FTAI), had obviated estrus detection (Rhodes *et al.*, 2002). One of the main features of estrus synchronization is short standing heat. In the more sophisticated protocols for FTAI, ovulation is synchronized (Pursley *et al.*, 1995; 1997), CR is lower but, most of the cows can be inseminated within a reasonable period after calving while there is no need to see the estrus signs of individual cows (Refsdal, 2000). Ideal synchronization program should have CR similar to the natural rate (Vasconcelos *et al.*, 1999). Goal of the present study was to compare effects of 2 different estrus synchronization protocols on the reproductive indices, plasma E2, P4 and milk P4 concentrations in dairy cows.

Corresponding Author: A.H. Fallah Rad, Department of Clinical Sciences, Faculty of Veterinary Medicine, Ferdowsi University of Mashhad, Mashhad, Iran

MATERIALS AND METHODS

Lactating Holstein dairy cows in a dairy farm in Mashhad, Iran, were randomly assigned into either nPG (n = 27) or OC (n = 25) groups. Average daily milk production of the cows in the barn was 38 kg with 4% milk fat with a VWP of 40±5 days. All the cows were in their 2nd parity on, had no history of dystocia, downer syndrome, RFM, clinical mastitis, metritis, vaginal laceration or vaginitis in their last parturition. If metritis was seen after start of the experiment, the cow was replaced with a healthy one. Cows in the nPG group received 2 consecutive IM injections of 0.5mg of Cloprostenol (Estrumate®, Schering-Plough, NL) 14 days apart. Upon observation of estrus after 2nd injection of PG, cows were inseminated. If there were no signs of estrus within 14 days, a 3rd, 4th or 5th injection was given. Ovsynch CIDR protocol was applied to cows in the OC group as explained by others (Pursley *et al.*, 1997). In case of return to estrus, the program was repeated for up to 3 times. In non-return cases, non pregnant cows, on day 45, were removed from the experiment.

On day 1, 5 and 21 post insemination, 8 mL of blood and 20 mL of milk were collected, blood plasma was separated and stored at -20°C until P4 and E2 were determined. CR from the 1st, 2nd and 3rd AI, DO, number of AI/conception and interval from the 1st AI to conception were calculated. ELISA kits (DRG Instruments, GmbH), were read by ELISA reader and Washer (Bio-Tek Instruments, USA). Milk P4 was measured by the method described by Waldmann *et al.* (2001).

Data were analyzed by SPSS software. For the analysis of non-normal data, non parametric tests were used. Concentrations of E2 and P4 in the 2 groups and within the groups were compared by independent and paired t-test, respectively. Levels of E2 and P4 in 3 different samplings were compared by repeated measures of GLM. For the correlation of plasma and milk P4 and also plasma E2 and P4 in each stage, Pearson correlation test was used. For the analysis of other factors, Mann-Whitney U test was used. Values of p≤0.05 were considered significant.

RESULTS

Table 1 and 2 shows reproductive indices of the 2 groups and concentrations of E2 and P4 in the plasma and P4 of the milk, respectively. Significant differences are shown by different letters (a and b; p<0.05).

On day 21, correlation coefficient for CR and plasma P4 was 0.622 (p<0.05) and 0.723 for milk P4 (p<0.05). Plasma E2 on day 5 and 21 in nPG group was lower in the

Table 1: Comparison of the reproductive indices of the 2 groups

Factor	nPG	OC
Number of cows	27	25
Pregnant cows	22 (81.48%)	21 (84%)
Number of previous pregnancies (Mean±SEM)	3.17±0.31	3.17±0.28
Corrected milk production (kg)	8093.34	8606.88
Calving to start of experiment (days, Mean±SEM)	41±1.08	41.82±0.97
Number of AI/conception (Mean±SEM)	3.13±0.26 ^a	2.23±0.23 ^b
1st service CR (%)	14.28	23.80
2nd service CR (%)	23.8 ^a	71.42 ^b
3rd service CR (%)	61.90 ^a	85.71 ^b
Total CR (Mean±SEM)	40.30±5.44 ^a	55.71±5.93 ^b

Table 2: Plasma E2 and P4 and milk P4 of the 2 groups (Mean±SEM)

Concentration of E2 (pg mL ⁻¹) and/or P4 (ng mL ⁻¹)	nPG group	OC group
Plasma E2 on day 1	33.38±4.05	26.28±1.17
Plasma E2 on day 5	26.19±3.12 ^a	36.28±2.41 ^b
Plasma E2 on day 21	25.85±6.02	26.61±2.81
Plasma P4 on day 1	4.47±0.87	3.07±0.77
Plasma P4 on day 5	2.82±0.24	2.95±0.54
Plasma P4 on day 21	6.17±1.23	5.95±1.50
Milk P4 on day 5	15.37±2.36 ^a	8.44±0.97 ^b
Milk P4 on day 21	24.17±3.57 ^a	15.89±2.15 ^b

pregnant as compared to the non pregnant cows (p<0.05). On day 21, in the pregnant OC cows, P4 of the plasma was higher than non pregnant cows (p<0.05).

DISCUSSION

FSCR in both groups were less than 30% and was similar to the results of others. It has been shown that FSCR in the ovsynch groups were lower than the cows inseminated after showing estrus signs (Alnimer *et al.*, 2002). In the herds with low estrus detection rate, use of estrus and/or ovulation synchronization along with FTAI may generally increase CR but, FSCR may be lower than 2nd or 3rd service (Xu *et al.*, 2000b). In the beginning of the present study, only 16.66% of the cows had less than 1n mL⁻¹ P4 in their plasma, Ovsynch protocols give the best results on day 5-9 of the cycle but, if it is applied late in the luteal phase, a large follicle is produced with lower CR (Cordoba and Fricke, 2001). Most of the cows with low FSCR might be in their mid luteal phase. Similar to results of others (Xu *et al.*, 2000a), CR from the 2nd and 3rd insemination and total CR in the CO group in the present study were higher than the nPG group (Table 1). Effectiveness of Ovsynch protocol may be due to better control of follicular development and ovulation (Mialot *et al.*, 1999).

When exogenous P4 is used in the synchronizing protocols, in the next cycle a durable follicle is produced and a larger CL with more P4 is produced naturally, therefore, CR may increase (Xu *et al.*, 2000a). CR from the

2nd AI was higher in the present research which confirms the above concept. Xu *et al.* (2000a) stated that total CR was not different when Ovsynch protocol was compared with repeated injections of PG (38.9 against 37.8%) but, in our experiment, total CR was higher in the OC than nPG while number of AI per conception was lower. By using Ovsynch protocol, Seguin (1997) could reduce DO from 121-98 days with no reduction in the CR. In the present experiment, mean DO and predicted calving interval was lower in the OC than nPG (106 and 387 days, for OC and, 138 and 418 days for nPG, respectively).

There are plenty of plausible reasons for the lower CR in the PG group which include: heat detection mistakes, AI in un-appropriate time, interference of the synchronization protocol with the follicular development, incomplete regression of the CL (Drillich *et al.*, 2000), short half life of the PG in some cows, long follicular phases after PG injections and cows with inactive cycles (Peters and Ball, 1995). PG treatment is not followed by luteolysis in at least 10% of the cows and within 24-48 h P4 production returns back. If CL is not active, it doesn't respond to PG. Negative energy balance and heat stress (Alnimer *et al.*, 2002), might be other reasons for low CR from the 1st insemination in both groups. Day 5 post insemination milk P4 was higher in OC than nPG probably due to the delay in ovulation, follicles and the subsequent CLs were larger and more P4 was produced (4 and 32). Highest CR was seen in cows with P4 concentrations between 3-9 ng mL⁻¹ of plasma on day 5 post AI (Starbuck *et al.*, 2001; Thatcher *et al.*, 2001).

Mean plasma E2 on day 5 after AI in nPG group was distinctly higher than OC which shows higher number of pregnant cows in the nPG from the 1st AI. Some cows have high E2 even on day 5-6 after standing heat which might due to the delayed ovulation. This phenomenon had been noticed in cows treated with PGF2 α (Humblot, 2001). When ovulation is close, 2nd injection of GnRH in Ovsynch protocol helps surge of LH, therefore, hastens ovulation (Berber *et al.*, 2002).

Mean milk fat P4 on day 21 in the OC was higher than nPG, showing higher CR in this group. Cows having less than 5 ng mL⁻¹ plasma P4 levels and not showing signs of estrus had lower chance of pregnancy and may need up to 4 inseminations (Francos, 1998). In the present study, there was a significant correlation between P4 on day 20-23 post AI with FSCR which is similar to the results of others (Alvarez *et al.*, 1989; Karagiannidis, 1990; Pieterse *et al.*, 1990).

CONCLUSION

From the results of the present study it could be concluded that ovulation synchronization by using Ovsynch-CIDR protocol as compared with 2 consecutive injections of PGF2 α , had decreased DO and CI, increased

CR from the 2nd and 3rd insemination, increased total CR and number of inseminations/ pregnancy. Cows assigned into Ovsynch-CIDR protocol had higher P4 on day 5 and 21 post insemination. In the present experiment cows were selected with specific conditions, therefore, these results may not be applicable to all cows of the herd or other herds.

ACKNOWLEDGEMENT

This research was funded by the Ferdowsi University of Mashhad, Mashhad, Iran.

REFERENCES

- Alnimer, M., De G. Rosa, F. Grasso, F. Napolitano and A. Bordi, 2002. Effect of climate on the response to 3 oestrous synchronization techniques in lactating dairy cows. *Anim. Reprod. Sci.*, 71 (3): 157-168.
- Alvarez, R.H., J.P. Massat and C.F. Meirelles, 1989. Early pregnancy diagnosis in cows using a milk progesterone immunoassay (ELISA) field test. *Revista Brasileira Reprod. Anim.*, 13: 25-32.
- Berber, R.C.A., E.H. Madureira and P.S. Barucelli, 2002. Comparison of 2 ovsynch protocols (GnRH versus LH) for fixed timed insemination in buffalo (*Bubalus bubalis*). *Theriogenology*, 57 (5): 1421-1430.
- Burke, C.R., M.P. Boland and K.L. Macmillan, 1999. Ovarian responses to Progesterone and Oestradiol benzoate administered intravaginally during dioestrus in cattle. *Anim. Reprod. Sci.*, 55 (1): 23-33.
- Butler, W.R., 2001. Nutritional effects on resumption of cyclicity and on conception rate. *Proc. Dairy Ferti. Northern Ireland*, pp: 13.
- Cordoba, M.C. and P.M. Fricke, 2001. Evaluation of 2 hormonal protocols for synchronization of ovulation and timed artificial insemination in dairy cows managed in grazing-based dairies. *J. Dairy Sci.*, 84 (12): 2700-2708.
- Drillich, M., B.A. Tenhagen and W. Heuwiezer, 2000. Effect of one spontaneous estrus cycle (after synchronization with PGF2 α) on reproductive performance in dairy cows. *Theriogenology*, 54 (9): 1389-1394.
- Francos, G., 1998. Association between milk Progesterone concentration after 1st insemination and conception in dairy cattle in Israel. *Vet. Rec.*, 142 (3): 63-64 21.
- Gröhn, Y.T. and P.J. Rajala-Schultz, 2000. Epidemiology of reproductive performance in dairy cows. *Anim. Reprod. Sci.*, 60-61: 605-614.
- Humblot, P., 2001. Use of pregnancy specific proteins and Progesterone assays to monitor pregnancy and determine the timing, frequencies and source of embryonic mortality in ruminants. *Theriogenology*, 56 (9): 1417-1433.

- Karagiannidis, A.K., 1990. Factors affecting the accuracy of early pregnancy diagnosis in cattle by RIA of Progesterone in milk. *Bull. Hellen. Vet. Med. Soc.*, 41 (4): 84-99.
- López-Gatius F., 2000. Reproductive performance of lactating dairy cows treated with cloprostenol, hCG and Estradiol benzoate for synchronization of estrus followed by timed AI. *Theriogenology*, 54 (4): 551-558.
- López-Gatius F., P. Santolaria, J. Yáñez, J. Rullant and M. López-Béjar, 2001. Persistent ovarian follicles in dairy cows: A therapeutic approach. *Theriogenol.*, 56 (4): 649-659.
- Macmillan, K.L., I.J. Lean and C.T. Westwood, 1996. The effect of lactation on the fertility of dairy cows. *Aus. Vet. J.*, 73 (1): 141-147.
- Mialot, J.P., G. Laumonier, C. Ponsart, H. Fauxpoint, E. Barassin, A.A. Ponter and F. Deletang, 1999. Postpartum subestrus in dairy cows: Comparison of treatment with prostaglandin F_{2α} or GnRH + prostaglandin F_{2α} + GnRH. *Theriogenology*, 52 (5): 901-911.
- Peters, A.R. and P.J.H. Ball, 1995. *Reproduction in Cattle*. 2nd Edn. Blackwell Science Ltd. London, UK, pp: 89-104.
- Peters, A.R., 1996. Herd management for reproductive efficiency. *Anim. Reprod. Sci.*, 42 (1-4): 455-464.
- Pieterse, M.C., O. Szenic, A.H. Willemse, C.S.A. Bajcsy, S.J. Dieleman and M.A.M Taverne, 1990. Early pregnancy diagnosis in cattle by means of linear-array real-time ultrasound scanning of the uterus and qualitative and quantitative milk Progesterone test. *Theriogenology*, 33 (3): 697-707.
- Pursley, J.R., M.O. Mee and M.C. Wiltbank, 1995. Synchronization of ovulation in dairy cows using PGF_{2α} and GnRH. *Theriogenology*, 44 (7): 915-923.
- Pursley, J.R., M.C. Wiltbank, J.S. Stevenson, J.S. Ottobre, H.A. Garverick and L.L. Anderson, 1997. Pregnancy rates per artificial insemination for cows and heifers inseminated at a synchronized ovulation or synchronized estrus. *J. Dairy Sci.*, 80 (2): 295-300.
- Refsdal, A.O., 2000. To treat or not to treat: A proper use of hormones and antibiotics. *Anim. Reprod. Sci.*, 60-61: 109-119.
- Rhodes, F.M., C.R. Burke, B.A. Clark, M.L. Day and K.L. Macmillan, 2002. Effect of treatment with Progesterone and Oestradiol benzoate on ovarian follicular turnover in postpartum anoestrus cows and cows which have resumed estrus cycles. *Anim. Reprod. Sci.*, 69 (3-4): 139-150.
- Seguin, B., 1997. Ovsynch: a method for breeding dairy cows without doing heat detection. *Bovine Practitioner*, 31 (1): 11-16.
- Starbuck, G.R., A.O. Darwash, G.E. Mann and G.E. Lamming, 2001. The detection and treatment of post insemination Progesterone insufficiency in dairy cows. *Proceedings of Dairy Fertility in Northern Ireland*, pp: 78.
- Thatcher, W.W., F. Moriera, J.E.P. Santos, R.C. Mattos, F.L. Lopez, S.M. Pancarci and C.A. Risco, 2001. Effects of hormonal treatments on reproductive performance and embryo production. *Theriogenol.*, 55 (1): 75-89.
- Vasconcelos, J.L.M., R.W. Silcox, G.J.M. Rosa, J.R. Pursley and Wiltbank, M.C. 1999. Synchronization rate, size of the ovulatory follicle and pregnancy rate after synchronization of ovulation breeding on different days of the estrous cycle in lactating dairy cows. *Theriogenol.*, 52 (6): 1067-1078.
- Waldmann, A., O. Reksen, K. Landsverk, E. Kommisrud, E. Dahl, A.O. Refsdal and E. Ropstad, 2001. Progesterone concentrations in milk fat at 1st insemination-effects on non-return and repeat-breeding. *Anim. Reprod. Sci.*, 65 (1-2): 33-41.
- Xu, Z.Z. and L.J. Burton, 2000. Estrus synchronization of lactating dairy cows with GnRH, Progesterone and prostaglandin F_{2α}. *J. Dairy Sci.*, 83 (3): 471-476.
- Xu, Z.Z. and L.J. Burton, S. McDougall and P.D. Jolly, 2000a. Treatment of non-cyclic lactating dairy cows with Progesterone and Estradiol or with Progesterone, GnRH, prostaglandin F and Estradiol. *J. Dairy Sci.*, 83 (3): 464-470.
- Xu, Z.Z., G.A. Verkerk, J.F. Mee, S.R. Morgan, B.A. Clark, C.R. Burke and L.J. Burton, 2000b. Progesterone and follicular changes in post partum non-cyclic dairy cows after treatment with progesterone and estradiol or with progesterone, GnRH, PGF_{2α} and Estradiol. *Theriogenology*, 54 (2): 273-282.