

## Urinary Urea Elimination Following Feeding of Low and High Protein Diets to Camels (*Camelus dromadarius*)

A.M. Homeida and S.A. AL-Shami

Department of Veterinary Public Health and Animal Husbandry,  
College of Veterinary Medicine and Animal Resources, King Faisal University,  
Camel Research Center, P.O. Box 1069, AL-Ahsa 31982, Saudi Arabia

**Abstract:** Feeding of urea-treated wheat straw to camels resulted in significantly increased concentration of plasma urea compared to their counterparts that fed straw only. P-aminohippuric acid, inulin and urea clearances and filtered urea were also significantly higher in animals fed urea treated straw than in animals fed straw. It is suggested that in camels fed straw, the renal urea spare could serve for ruminal recycling of urea to compensate for low nitrogen diet.

**Key words:** Urea, urine, protein, diet, camel, wheat straw

### INTRODUCTION

It is well known that in ruminants, endogenous urea is partly recycled in the forestomach. This process is nutritionally advantageous for ruminants because ruminal bacteria are able to use urea nitrogen to synthesis protein that will be absorbed in the small intestine (Von Engelhardt and Schnider, 1977). Urea treatment of poor quality roughages has emerged as promising technology for improving their feeding value (Majumdar *et al.*, 2003). Urea is hydrolysed to ammonia due to the action of urease enzyme. The generated ammonia, react with the cell walls of roughages (Klopferstein, 1978), increasing the crude protein content (Roseler *et al.*, 1993; AL-Shami and AL-Sultan, 2006).

Wheat straw is the most abundant agriculture in Saudi Arabia. It is widely used as cheap source of bulk feed to ruminants. The feed value of wheat straw is, however limited by its low available energy, low protein, minerals and vitamins (Saddullah *et al.*, 1981). It is now established practice in some parts of the world to upgrade the protein content of low quality roughage by urea treatment (AL-Shami and AL-Sultan, 2006).

Camels graze different kinds of pastures with variable levels of nitrogen content throughout the year. This study was undertaken to investigate the importance of renal mechanisms to control urea losses in camels fed low or high urea-treated straw.

### MATERIALS AND METHODS

**Urea treatment of wheat straw:** One ton of wheat straw (tbin) was treated with 4% w w<sup>-1</sup> urea (Saudi Basic Industries Corporation) solution (12.5%) and subjected to ureolysis by anaerobic storage for 8 weeks. This treatments of urea was expected to increase the crude protein content of straw from 3% in straw to 8% in urea treated straw (AL-Shami and AL-Sultan, 2006).

**Experimental animals and treatments:** Twelve mature (2-4 years old) and weigh 300-400 kg body weight camels were used in the study. They were divided equally into 3 groups.

**Group 1:** Animals were given wheat straw as a sole diet for 21 days. Water and mineral salt licks were provided *ad libitum*.

**Group 2:** Animals were treated similar to group1 but in addition were given each 1 kg urea-treated straw and 2 kg of concentrates.

**Group 3:** Animals were treated similar to group 1 but in addition were given each 2 kg urea-treated straw and 2 kg of concentrates.

On the last day of the feeding period the renal function was studied. Both jugular veins of each animal

were catheterized. One vein was used for perfusion and the other one for blood sampling. In order to measure renal plasma flow and the glomerular filtration rate, a primed constant infusion of 0.5 mL min<sup>-1</sup> of P-aminohippuric acid (PAH, 8 mg/ml/min) and inulin (70 mg/ml/min) (Sigma, UK) in 0.9% saline, was made into a jugular vein. Bladder urine was collected for each period and a heparinized blood sample (5 mL) was obtained at the midpoint of each period. Blood samples were centrifuged at 3000 rpm for 5 min, to obtain plasma which was stored until analysis.

Plasma and urine were analysed for urea (urease Kit, Wiener, Austria), inulin and PAH (Tebot *et al.*, 2002). Renal plasma flow (as PAH clearance), glomerular filtration rate (as inulin clearance) and urea clearance were calculated as the product of urine flow and urinary concentration divided by plasma concentration. Fractional excretion of urea (as urea/inulin clearances) and filtered load of urea (as inulin clearance by plasma urea concentration) were determined.

Results were subjected to Analysis of Variance (ANOVA) using least significance difference for comparing differences between means (SAS, 1985).

## RESULTS

Effects of feeding urea-treated straw on plasma urea concentrations and kidney functions are given in Table 1. Feeding of urea treated straw resulted in significantly ( $p < 0.05$ ) increased concentration of plasma urea in group 2 and 3 animals compared to group 1. Significantly ( $p < 0.05$ ) increased losses of nitrogen in urine was seen in animals in group 2 and 3 compared to those in group 1. PAH, inulin and urea clearances and filtered urea were also significantly ( $p < 0.05$ ) higher in group 2 and 3 animals compared to those in group 1. Fractional urea excretion was significantly ( $p < 0.05$ ) higher in animals of group 1 than in animals of group 2 and 3. No adverse effect of feeding urea-treated straw was shown in camels.

Table 1: Urea kinetics in camels fed urea-treated wheat straw

| Parameter                                        | Group 1    | Group 2    | Group 3    |
|--------------------------------------------------|------------|------------|------------|
| Plasma urea concentration (mg dL <sup>-1</sup> ) | 10.1±0.6a  | 30.2±2.1b  | 35.3±2.3c  |
| Nitrogen losses in urine (g day <sup>-1</sup> )  | 16.2±1.3a  | 29.0±2.6b  | 34.0±2.5c  |
| PAH clearance (mL min <sup>-1</sup> )            | 602±22.0a  | 405±20.0b  | 426±21.0b  |
| Inulin clearance (mL min <sup>-1</sup> )         | 16.3±13.0a | 52.1±2.3b  | 54.3±2.4b  |
| Urea clearance (mL min <sup>-1</sup> )           | 13.1±0.6a  | 36.6±1.4b  | 38.3±1.6b  |
| Filtered urea (mg min <sup>-1</sup> )            | 3.2±0.06a  | 16.2±1.3b  | 17.6±1.5b  |
| Fractional urea excretion                        | 0.81±0.03a | 0.65±0.04b | 0.61±0.03b |

a,b,c means bearing different letters in the same row differ significantly  $p < 0.05$

## DISCUSSION

The results of the present study indicate that feeding of urea-treated straw increased blood urea concentration. The increased value of blood urea level in animal fed urea-treated straw was due to consumption of more amount of easily degradable non-protein nitrogen available through ammoniated straw which was in normal physiological range, thus no adverse effect of feeding urea treated straw diets were visible (Bhatt *et al.*, 2004; Singh *et al.*, 2007). Animal fed wheat straw of low protein content were adapted by reducing glomerular filtration, urea clearance and filtered load of urea. Similar finding were reported in sheep on restricted protein intake (Tebot *et al.*, 2002). Data on nitrogen balance indicated that loss of nitrogen through urine was significantly more in urea supplemented groups, indicating that ammonia-nitrogen could not be utilized completely due to less available energy. Similar nitrogen excretion was reported in crossbred calves and buffaloes fed diet containing higher than optimal amount of protein (Dass and Arora, 1989) and with various levels of urea with molasses (Ali and Mirza, 1986). This suggests that the ammonia-nitrogen utilization depends on the activity of rumen microflora and rumen microbes are maximum in quantity when there is matching supply of energy to the available nitrogen (Toppo *et al.*, 1997). Changes in tubular handling of urea (fractional excretion) induced by feeding wheat straw was also evident. An enhanced capacity for urea reabsorption from tubules and the renal pelvis was noticed when protein intake was restricted (Cirio and Biovin, 1990). This adaptive process prevents an excessive reduction of blood urea pool (Mcknight, 1994). The camel often re-uses nitrogenous waste by returning urea to the stomach via bloodstream and liver, rather than voiding it.

## ACKNOWLEDGEMENT

The authors thank the Deanship of Scientific Research of King Faisal University and Saudi Basic Industries Corporation (SABIC) for financial support.

## REFERENCES

- Ali, A. and I.H. Mirza, 1986. Feeding of urea molasses blocks buffalo chocolate to ruminants in the tropic. *Asian Livest.*, 6: 160-166.
- AL-Shami, S.A. and S.A. AL-Sultan, 2006. Effect of treating wheat straw with urea on its crude protein content. *J. Anim. Vet. Adv.*, 3: 106-108.

- Bhatt, C., R.S. Jiswal, T.C. Thakur and Y.P. Joshi, 2004. Effect of urea treated blade wheat straw based diets on the performance of crossbred heifers. *Indian J. Anim. Nutr.*, 21: 236-239.
- Cirio, A. and R. Boivin, 1990. Urea recycling for the renal pelvis in sheep: A study with <sup>14</sup>C urea. *Am. J. Physiol.*, 258: F1196-F1202.
- Dass, R.S. and S.P. Arora, 1989. Effect of feeding aflatoxin on N-kinetics in crossbred and buffalo calves fed different levels of protein. *J. Nuclear Agric. Biol.*, 18: 233-238.
- Klopferstein, T., 1978. Chemical treatment of crop residues. *J. Anim. Sci.*, 46: 841.
- Majumdar, A.B., M.M. Dar, S.B. Maity and A. Kumar, 2003. Nutrient utilization and performance of crossbred cows fed urea treated straw based diets. *J. Anim. Nutr. Feed Technol.*, 3: 173-178.
- Mcknight, T.L., 1994. *The camel in Australia*, Melbourne University Press.
- Roseler, D.K., J.D. Ferguson, C.J. Sniffen and J. Herrema, 1993. Dietary Protein Degradability effect on plasma and milk urea nitrogen in Holstein cows. *J. Dairy Sci.*, 76: 525-534.
- Saddullah, M., M. Haque and F. Dolberg, 1981. Effectiveness of ammonia-treated straw in feeding of dairy cows and growing cattle. *Agric. Environ.*, 6: 267-271.
- Singh, A., R.S. Jawishal, S.S. Chauhan, T.C. Thakur, V. Singh and Y. Joshi, 2007. Impact of feeding urea ammoniated paddy straw on utilization of nutrients blood, metabolites and cost effectiveness on crossbred lactating cows. *Indian J. Anim. Sci.*, 77: 595-598.
- SAS (Statistical Analysis System Institute), 1985. SAS. User guide. SAS. Inc. Cary, NC.
- Tebot, I., A. Britos, J.M. Codeau and A. Cirio, 2002. Microbial Protein production determined by urinary allantoin and renal urea sparing in normal and low protein fed Corriedale sheep. *Vet. Res.*, 33: 101-106.
- Toppo, S., A.K. Verma, R.S. Dass and U.R. Mehra, 1997. Nutrient utilization and rumen fermentation pattern in crossbred cattle fed different planes of nutrition supplemented with urea molasses mineral block. *Anim. Feed Sci. Technol.*, 29: 333-341.
- Von Engelhardt, W. and W. Schnider, 1977. Energy and nitrogen metabolism in the llama. *Anim. Res. Develops*, pp: 68-71.