

Influence of Dietary Phytoadditive as Polyherbal Combination on Performance of Does and Respective Litters in Cross Bred Dairy Goats

F. Mirzaei and S. Prasad
Livestock Production and Management Division, National Dairy Research Institute,
Karnal, 132-001 Haryana, India

Abstract: The aim of the present research was to study the effects of a polyherbal supplement on cross bred does, starting from the last month of pregnancy to weaning on milk yield, kid birth weight and growth rate. A total of 30 does were divided into three treatments of ten each in individual pens: Low level Supplementation (LS), High level Supplementation (HS) and Non-Supplemented treatment (NS) as control. Low supplemented goats were given 125 mg kg⁻¹ BW day⁻¹ of polyherbal combination; high supplemented goats were given 250 mg kg⁻¹ BW day⁻¹. The study was carried out in 2008. About 59 kids were born from all the experimental animals based on treatment wise. There was no difference on milk yield between supplemented groups and control ($p>0.05$) although, polyherbal supplementation had positive effect on litter birth weight and growth rate compared to control. Weaning weights were higher ($p<0.001$) in LS and HS compare to NS does. In both supplemented treatments compared to control, mortalities and morbidities were also lower in kids born. It is concluded that pre-partum to weaning supplementation increases kids growth rates and weaning weights as well as reduces kid mortalities but it doesn't have significant effect on milk production.

Key words: Doe performance, goat kid growth, polyherbal supplement, mortalities, treatment, India

INTRODUCTION

The goat is among the earliest species to be domesticated which records show happened around 7000 years B.C. in Southeast Asia along the present Iraq-Iran borders (Mason, 1981). Goats are kept for various reasons of which milk and meat are the most important. The use of goat milk for human consumption is as old as the domestication of animals of economic importance (Dubeuf, 2005). The importance of goats as providers around the world of essential food in meat and dairy products has been discussed and documented in many recent proceedings of national and international conferences (Bouvier *et al.*, 2000; Haenlein and Fahmy, 1999; Haenlein, 1992, 2001; Morand-Fehr and Boyazoglu, 1999; Rubino *et al.*, 1999). Under this system, inaccessibility and cost of timely veterinary health care are major constraints to the viability and productivity of goat production in many regions of the world. As recognized by the World Health Organization, local ethno-veterinary medicines could play an important role in ensuring general well-being and welfare of livestock in the developing world. Since extracting the effective agent (s) from these herbs is uneconomic, it is better to use them in the form of mixture

as few herbs have the galactogogue property besides the capacity of improving the palatability and carminative so aiding the digestion process. Therefore, it is desirable to use the herbs as these are a good source of plant secondary metabolites and available in the country abundantly at cheaper rate. Farmers are also aware of their use and they can adopt this approach easily. Keeping these facts in view, the potential of a herbal mixture based on commonly available and cheap herbs was evaluated for its efficacy for increasing of dairy goat performance. The present study was therefore undertaken to evaluate efficacy of polyherbal supplement in kid performance and doe productivity. It was hypothesized that polyherbal combination would result in higher kid growths and doe weight gain.

MATERIALS AND METHODS

Study area and experimental diets: The experiment was carried out at livestock farm of the National Dairy Research Institute, Deemed University (NDRI), Karnal, situated in eastern zone of Haryana and in the Trans Gangetic Plain Region of India at an altitude of 250 m above mean sea level on 29°42'N latitude and 75°94'E longitude. The minimum ambient temperature falls near to

freezing point in winter and maximum goes approximately up to 45°C in May/June. The annual rainfall is close to 700 mm most of which is received from July-September. The climate of the farm is typically sub-tropical and the land area is very productive with sufficient irrigation facilities for growing green fodder for animals year round. The relative humidity of this farm varies from 41-85% and vapour pressure ranges from 7.0-25 mm of Hg. Thus this farm receives the extreme climates due to the wide range of variation in various meteorological factors. The polyherbal supplement contained; *Asparagous racemosus* (Shatavari), *Leptadenia reticulata* (Jivanti), *Nigella sativa* (Kolonji), *Cuminum cyminum* (Jeera) and *Pueraria tuberosa* (Vidarikand). Individual herbs were procured from local market after assessing their quality in consultation with ayurvedic practitioners and drug manufacturers. Used parts of each herb were as follows for Asparagous and Pueraria (root), Nigella and Cuminum (seed) and Leptadenia (leaf). They were purchased as dried (sun-dried). Each sun-dried herb was pulverized separately. The polyherbal supplement was prepared after mixing powdered specific parts of five herbs in same proportion based on weight of dry matter. The dried samples of the feeds (concentrates and herbs) were ground through 1 mm sieve and further dried at 105°C for 1 h to determine the dry matter. The Crude Protein (CP) in and dried samples of the feeds and herbs was determined according to Kjeldahl procedure (AOAC, 1990) while the Neutral Detergent Fibre (NDF) and Acid Detergent Fibre (ADF) of feeds were determined according to the procedure of Van Soest *et al.* (1991). The AOAC (1990) procedures were followed to determine the ash content of the feeds and herbs and also that of the Crude Fibre (CF) and Ether Extract (EE) for feeds. The samples were ashed by charring in a Muffle furnace at 500°C for about 3 h or until a whitish ash remained. Total phenolics and condensed tannins were analyzed colorimetrically (Milton Roy 401 Spectronic, spectrophotometer) using modified Folin-Dennis procedures (AOAC, 1990) and vanillin-HCL procedures (Price *et al.*, 1978), respectively.

Experimental design and animal management: Beetal is one of the heaviest dairy type goat breeds of Northern India. The animals are characterized by a large size, long drooping ears and roman nose. The Beetal breed has been used for cross-breeding with Saanen and Alpine breeds in the all India coordinated research project on goats both for milk and meat components (Rana *et al.*, 1981). Lactation responses to the supplements were tested with 30 cross bred of Alpine×Beetal (AB) goats (Mean body

weights 44.6±4.20 kg). A general management program for de-worming, disease prevention and hoof trimming was followed throughout. Goats were selected on the basis of milk production records from a herd at the station and assigned randomly (10 doe each) to a supplement of polyherbal combination at 125 mg kg⁻¹ Body Weight (BW) as low level polyherbal supplement (LS) and 250 mg kg⁻¹ BW as high level polyherbal supplement (HS) for 6 weeks before kidding till weaning time of their kids. One treatment with equal numbers of doe served as control without supplement (NS). To avoid dominance behavior and to ensure equal access to the supplement each goat from every treatment has been randomly assigned to separate pens, eliminating of possible biases due to environmental variation within the animal house. Pregnant goats were kept on isocaloric and isonitrogenous diet according to NRC (1981) feeding standard. The animals were kept in the shade in individual feeding pens. The experimental diets offered to the goats consisted of concentrate mixture according to requirements in advanced pregnancy and green fodder (Berseem) as *ad libitum*. Clean and fresh water was always available for consumption. The feed was given to the animals twice a day; in the morning (9:00 am) and afternoon (2:00 pm). Goats were adapted for 10 days to the experimental diets before measurements. After kidding, goats were hand milked twice daily (6.00 and 16.00 h) that milking commenced on day 8 post partum. Daily milk yield of individual goats was recorded throughout the experiment. BW changes were determined by weighing each goat for three consecutive days at the beginning and subsequently at fortnightly intervals early in the morning before feeding. About 30 advanced pregnant cross bred goats of an average age of 2.3 years were randomly allocated to three dietary treatments using a Complete Randomized Design (CRD). Goats in different treatments were tagged with different colours for easy identification. The does were mated in June/July 2008 and kidded in November/December 2008. In September, a month pre-partum they were vaccinated against pulpy kidney, dosed against internal parasites. Animals were allowed access to green fodder in open area and water during the day for 7 h and penned overnight under the same environment throughout the trial. Doe post-partum weights, kids weights, sex and type of birth (single or twin) were recorded on day one postpartum. Thereafter, weekly weights for both does and kids were recorded in between 8 and 9 am up to weaning. In addition, the kids mortalities and morbidities were recorded. The does were left with their respective kids for 7 days in the kidding

paddock. Milking started on the 8 day post-partum to weaning at an average of 100 days of kids' age. After kidding, the dams were weighed to know the parturition weights which were taken as the initial weights. Milk yield measurement was commenced after the kids were allowed to suckle the dams for the first 7 days postpartum to consume colostrum and to establish strong dam-kid relationship to forestall rejection of kids by their dams after overnight separation to measure milk yield. The kids were separated from their dams for 12 h over night (6 pm to 6 am) and only re-introduced to their dams after milking.

Statistical analysis: Data were analyzed using the least squares means Harvey. Analyses of Variance (ANOVA) were performed to establish effects of supplementation on sex and type of birth and effects of supplemented feed on kid growth rate and milk output. The kid parameters analyzed were birth weights, growth rates and weaning weights and doe parameters analyzed were milk yield, post-partum weight and doe weight at weaning. Duncan's multiple range was used to determine significant differences between the means. The results were expressed as the mean±standard error of the mean. For reproduction performance only the effect of treatment was analyzed using descriptive statistics.

RESULTS

Supplementation period: Phytochemical and proximate composition of offered polyherbal combination and feeds have been shown in Table 1 and 2. The efficacy of supplementation on milk production, kid birth weight, doe post-partum weight and doe weight at weaning as well as the effect of milking on kid weight from birth to weaning and doe performance have been shown in Table 3. There was no effect of treatment ($p>0.05$) on doe body weight post partum or at weaning and milk yield. The fortnightly pattern of milk yield was shown in Fig. 1. Total milk produced was 2.41 ± 0.12 kg in LS, $1.932.37\pm0.12$ kg in HS and 2.37 ± 0.12 kg in NS does ($p>0.05$). The litters birth weight and daily BW gain were similar between treatments but weaning BW were higher ($p<0.001$) at lower dose as well as high level of supplement compared to control. At birth, litters Live Weight (LW) were similar between the low level supplement and high level supplement compared to non-supplement but was tended to be higher in LS rather than HS. Similarly, average daily growth rate was higher for two supplemented treatments compared to control and ranged from 0.147 ± 0.02 - 0.198 ± 0.02 kg day⁻¹.

Table 1: Physical composition of concentrate mixture

Ingredients	Parts (%)
Maize	33
Groundnut cake	20
Mustard cake	13
Wheat bran	20
Deoiled rice bran	11
Mineral mixture	2
Common salt	1
Total	100

Table 2: Plant secondary metabolites and nutritional properties of polyherbal supplement and feeds

Parameters (%DM basis)	Polyherbal supplement	Concentrate	Berseem
OM	90.80	91.30	89.90
CP	6.46	20.50	17.80
EE	0.35	4.23	3.34
Total ash	2.50	8.67	10.10
NDF	38.10	43.50	31.30
ADF	13.50	13.50	26.50
Total phenolics	4.57	-	-
Total tannin	3.69	-	-

Table 3: Summary of litter and doe growth performance due to polyherbal supplementation

Variables	N	NS	LS	HS	SEM	p-value
Milk yield (kg day ⁻¹)	30	2.37 ⁰	2.41 ⁰	1.93 ⁰	0.12	0.010
Litter birth W (kg)	59	5.120	6.670	6.400	0.65	0.220
Doe postpartum W (kg)	30	39.600	40.100	39.000	1.84	NS
Litter weaning W (kg)	58	13.76 ⁰	17.42 ⁰	17.02 ⁰	0.68	0.001
Doe W at weaning (kg)	30	42.850	42.300	43.500	2.06	NS
Kids daily gain (kg day ⁻¹)	58	0.147	0.178	0.198	0.02	0.070

NS: Non polyherbal Supplemented group; LS: Low polyherbal supplemented group; HS: High polyherbal supplemented group; N: Number of animal; values in the rows with different superscripts are significantly different




Fig. 1: Fortnightly changes in milk yield (LSM±SEM) in different treatments groups (NS: Non polyherbal supplemented group; LS: Low polyherbal supplemented group; HS: High polyherbal supplemented group)

The results of the reproductive performance of lactating goats observed during the experiment are shown in Table 4, results on total number of kids born, weaned and those that died in different treatments indicate there is slightly improvement of reproduction performance in supplemented groups rather than control group, it might be caused by immunological aspects of used medicinal plants.

Table 4: Total number of kids born, weaned and those that died in different treatments

Variables	NS	LS	HS
Total kids born	17	24	18
Single	2	1	2
Twin	6	7	5
Triple	1	2	2
Overall kid mortality (%)	6	0	0
Mortality of single (%)	50	0	0
Mortality of twin (%)	0	0	0
Mortality of triple (%)	0	0	0
Total number weaned	16	24	18
Single weaned	1	1	2
Twin weaned	6	7	5
Triple weaned	1	2	2
Male (%)	47	46	55
Female (%)	53	54	45

NS: Not polyherbal supplemented group; LS: Low polyherbal supplemented group; HS: High polyherbal supplemented group

Table 5: Mean±SE of reproduction parameters of lactating goats in different treatment groups

Parameters	NS (control)	LS	HS
Gestation length (day)	151.33±7.54	146.33±1.89	145.40±2.11
Litter size and survivability/animal	1.80±0.29	2.20±0.20	2.00±0.21
Kid birth weight (kg)	2.99±0.10	3.20±0.15	3.21±0.16
Litter birth weight (kg)	5.58±0.49	6.64±0.56	5.88±0.49
Twining rate (%)	90	110	100
Pregnancy rate (%)	70	80	90
Kidding disorders (case)	2	0	1

NS: Non polyherbal supplemented group; LS: Low polyherbal supplemented group; HS: High polyherbal supplemented group

Supplementation withdrawal period: The performance of does after stopping of polyherbal supplement was studied up to next parturition by December, 2009. The results of the reproductive performance of lactating goats observed during the experiment are shown in Table 5. The average gestation length (day) was 151.33±7.54, 146.33±1.89 and 145.40±2.11 for goats in NS, LS and HS, respectively. The goats in HS and LS were better for the length of gestation compared to goats in NS. The pregnancy rates (%) were 70.00, 80.00 and 90.00 for goats in NS, LS and HS, respectively. Goats in HS had higher pregnancy rate followed by the goats in LS. Two goats in NS aborted whereas in LS and HS all goats gave successful births. There were variations in the average litter size (2 kids/head) of lactating goats so it was 1.80±0.29, 2.20±0.20 and 2.00±0.21 for goats in NS, LS and HS, respectively. The results were in agreement with Beyan (2009) findings. The average litter BW (kg) per goat was 5.58±1.00, 6.64±0.56 and 5.88±0.49 for goats in NS, LS and HS, respectively while the corresponding average birth weight (kg) of kids was 2.99±0.10, 3.20±0.15 and 3.21±0.16, respectively. The goats in LS recorded higher litter body weight and HS showed higher kid birth weight compared to NS. Kid birth weights were similar ($p>0.05$). Litter body weight growth rates and weaning weights were different between treatments ($p<0.05$), the heaviest being from LS




Fig. 2: Effects of supplementation on litter live weight (kg) at different times (LSM±SEM); (NS: Non polyherbal supplemented group; LS: Low polyherbal supplemented group; HS: High polyherbal supplemented group)

does followed by HS does then the NS does (Fig. 2). Kid disorders were highest in NS (12%) and lowest in LS group (0%) which it has been shown in Table 5.

DISCUSSION

The study showed that phytochemical additives nutrition had an effect on milk production with low polyherbal supplemented does producing more milk than the non-supplemented and high supplemented does. This trend shows that supplementation is mostly of benefit in the 1st months of lactation period. No difference in supplementation and non-supplementation was however shown in doe weight post-partum and at weaning suggesting that supplementation had no effect on doe weight changes. Polyherbal combination nutritional treatment effects were also evident on kid performance with kids from low supplemented does having the highest average daily weight gains and weaning weights as well as lowest disorders. Goat productivity as measured by increased conception rates, kid survival and kid growth rate has been enhanced by polyherbal supplementation of does which is agreement with findings of Sikosana *et al.* (1990). The constraint of high kid disorders and mortalities (30-50%) up to weaning has been reported by Matika and Maphosa (1992), reduction in kid disorders in this study therefore, shows that supplementation of does can improve kid survival. Kids from polyherbal supplemented does however had higher gains than those from non-supplemented. The similarity in mean birth weights of kids in all supplemented treatments in following year which had been stopped supplementation of polyherbal combination suggests that the survival strategy of supplemented goats was higher than that of non-supplemented ones. It has also been shown in sheep that efficiency of foetal uptake of maternal glucose is only about 69% as a result of a substantial backflow of glucose

from foetus to mother (Greyling, 2000). Type of birth had an effect on kid growth. There were also more twins on LS and HS does than NS does. This might explain the higher daily weight gains and weaning weights in kids from supplemented than from non-supplemented does. Sex also had an effect on kid growth with males being heavier than females at birth. Consistent superiority of males has been widely reported with similar trends being reported by Tawonezvi and Ward (1987). This has been attributed to hormonal differences between sexes and their resultant effects on growth (Meyer, 2001). Overall doe productivity was measured using kids weaning weights and total milk production. Does that were supplemented gave higher milk yields and their average kids weaning weights at studied weeks were higher than that of non-supplemented group suggesting that supplementation improves overall doe productivity. The results of the present study depicted that application of polyherbal supplementation both in higher as well as lower level had better reproductive performance compared to control treatment. The improvements in reproductive performance achieved by supplementing polyherbal combination at the rate of 0.125 and 0.250 g kg⁻¹ BW could be due to stimulation of the reproductive process by the steroidal saponins contained in supplementation.

Berhane and Singh (2002) reported that highest reproductive performances (onset of estrus, pregnancy rate) were recorded in dairy cows supplemented with fenugreek as compared to the control group without supplementation. Rajkhowa *et al.* (2001) studied the effect of *Saraca asoca* stem bark and *Trigonella foenum-graecum* seeds on reproductive performance, serum progesterone and micro minerals profile in anoestrus cows and reported that the percentage of animals induced in oestrus and overall pregnancy rate using fenugreek seeds were higher (83.33 and 80.00%) as compared to *Saraca asoca* (66.66 and 50.00%) at doses of 50 and 100 g level.

Asparagus racemocos (Shatavari) is an immunomodulation and immunopotential in late gestation have been shown to shortening the uterine involution period (Qureshi *et al.*, 1997; Sattar *et al.*, 2003a, b; Sattar *et al.*, 2007; Amer and Badr, 2008). Estrogenic (Mitra *et al.*, 1999; Pandey *et al.*, 2005) property of shatavari, one of the polyherbal ingredient which stimulate the ovarian function, improves uterine tonicity thus helps in early uterine involution which consequently results into early initiation of estrus cycle. Additionally, resumption ovarian cyclicity after parturition depends on the nutritional status, body energy reserved and blood glucose level of the animal. Mitra *et al.* (1999) also reported that shatavari based herbal formulation did not possess oxytocin like activity which might be useful in uterine hypermotility associated early abortion.

CONCLUSION

In the study, feed additive sources like polyherbal combination that are cheaper than commercial feed additives can be stored and used during time of need when doe nutritional requirements are high but feed resources are scarce. Feeding of does pre and post-partum with 125 mg kg⁻¹ BW of polyherbal combination gave satisfactory results on overall doe productivity and kid performance. Supplementary feeding would however improve milk production as well as overall doe productivity, measured by prolificacy, kid survival and weaning rate and kid weaning weight. Smallholder farmers would therefore benefit from more milk for their home consumption and at the same time get more kids per kidding season thereby increasing flock sizes and consequently higher income. Supplementation was however important in early lactation due to the effect of negative energy balance at that time. There is however, need for further studies on kid growth rates post weaning to 18 months as well as doe complete lactation period and conception rates to determine if supplementation have any effect after weaning.

The beneficial effects of supplementary feeding with polyherbal combination on goat production in India were demonstrated in this study. It helps substantially reducing the incidence of abortion and increases the overall yield of kids per animal. Pregnancy rate was higher in LS group than HS and NS groups.

IMPLICATIONS

Goat is considered as poor people cow so researchers should consider economic management of this domestic animal. Therefore for enhancement of their productivity by using of eco-friendly and non side effect feed additives like medicinal plants as locally available resources and timely veterinary health care, researchers may help their holders who are almost from developing countries with low income to sustain their livelihood.

ACKNOWLEDGEMENTS

The researchers are grateful to the livestock farm for the goats and facilities which were used in this study. The study was funded by the division of Livestock Production Management of National Dairy Research Institute (NDRI).

REFERENCES

- AOAC, 1990. Official Methods of Analysis. 15th Edn., Association of Official Analytical Chemists, Arlington, VA., USA.

- Amer, H.A. and A.M. Badr, 2008. Influence of antepartum administration of immunopotentiators on reproductive efficacy of buffalo and viability of their newborn. *Vet. Ital.*, 44: 373-382.
- Berhane, M.B. and V.P. Singh, 2002. Effect of feeding indigenous galactopoietics feed supplements on milk production in crossbred cows. *Indian J. Anim. Sci.*, 72: 609-611.
- Beyan, M., 2009. Effect of fenugreek (*Trigonella foenum graecum*) seeds as feed additive on production performance of dairy goats. Ph.D. Thesis, National Dairy Research Institute, Deemed University, Karnal, India.
- Bouvier, F., E. Manfredi, F. Pilla and M. Heyman, 2000. Allergic sensitization to milk proteins in guinea pigs fed cow milk and goat milks of different genotypes. Proceedings of the 17th International Conference on Goats, (ICG'00), Institute de l'Elevage, Tours, France, pp: 874-874.
- Dubeuf, J.P., 2005. Structural, market and organisational conditions for developing goat dairy production systems. *Small Rumin. Res.*, 60: 67-74.
- Greyling, J.P.C., 2000. Reproduction traits in the Boer goat doe. *Small Rumin. Res.*, 36: 171-177.
- Haenlein, G.F.W. and M.H. Fahmy, 1999. Role of small ruminants in the supply of animal products. *Small Rumin. Res.*, 34: 175-308.
- Haenlein, G.F.W., 1992. Role of goat meat and milk in human nutrition. Proceedings of 5th International Conference on Goats, Mar. 1-8, ICAR Publ., New Delhi, India, pp: 575-580.
- Haenlein, G.F.W., 2001. Past, present and future perspectives of small ruminant dairy research. *J. Dairy Sci.*, 84: 2097-2115.
- Mason, I.L., 1981. Classification and Distribution of Goat Breeds. In: Genetic Resources of Pig, Sheep and Goats. Maijala, K. (Ed.), World Animal Science B8. Elsevier, Amsterdam, pp: 405-411.
- Matika, O. and V. Maphosa, 1992. Assessment of kid mortality in goats reared under a semi-extensive system. Matopos Research Station, Bulawayo, Annual Report, Government of Zimbabwe, pp, 24-27.
- Meyer, H.H.D., 2001. Biochemistry and physiology of anabolic hormones used for improvement of meat production. *APMIS*, 109: 1-8.
- Mitra, S.K., S. Gopumadhavan, M.V. Venkataranganna, D.N.K. Sharma and S.D. Anturlikar, 1999. Uterine tonic activity of U-3107 (EveCare), a herbal preparation in rats. *Indian J. Pharmacol.*, 31: 200-203.
- Morand-Fehr, P. and J. Boyazoglu, 1999. Present state and future outlook of the small ruminant sector. *Small Rumin. Res.*, 24: 175-188.
- NRC., 1981. Nutrient Requirements of Goats: Angora, Dairy and Meat Goats in Temperate and Tropical Countries. National Academy Press, Washington, DC.
- Pandey, S.K., A. Sahay, R.S. Pandey and Y.B. Tripathi, 2005. Effect of *Asparagus racemosus* rhizome (Shatavari) on mammary gland and genital organs of pregnant rat. *Phytother. Res.*, 19: 721-724.
- Price, M.L., S. van Scoyoc and L.G. Butler, 1978. A critical evaluation of the vanillin reaction as an assay for tannin in sorghum grain. *J. Agric. Food Chem.*, 28: 1214-1218.
- Qureshi, Z.I., L.A. Lodhi and A. Sattar, 1997. An apparent effect of immunopotential during late gestation on the postpartum reproductive performance of Nili-Ravi buffaloes (*Bubalus bubalis*). *Vet. Res. Commun.*, 21: 375-380.
- Rajkhowa, S., G.C. Hazarika, G.N. Dutta and B.N. Neog, 2001. Comparative efficacy of Ehb, a herbo mineral complex and Chelated iron in the prevention and control of piglet anaemia. *Phytomedica*, 2: 31-35.
- Rana, Z.S., D.S. Balaine and S. Ram, 1981. Defining Beetal goats. A useful breed of North India. *Livestock Adviser*, 6: 26-28.
- Rubino, R., P. Morand-Fehrb, C. Renieric, C. Perazad and F.M. Sarti, 1999. Typical products of the small ruminant sector and the factors affecting their quality. *Small Rumin. Res.*, 34: 289-302.
- Sattar, A., L.A. Lodhi, I. Ahmad, Z.I. Qureshi and N. Ahmad, 2003b. Study of some production traits in cows and their calves after immunostimulation of pregnant dams during late gestation. *Pak. Vet. J.*, 23: 169-172.
- Sattar, A., L.A. Lodhi, Z.I. Qureshi, I. Ahmad and N.A. Naz, 2003a. Efficacy of immunopotentiators on postpartum reproductive performance in Sahiwal cows. *Pak. Vet. J.*, 23: 130-133.
- Sattar, A., R.H. Mirza and S.M.I. Hussain, 2007. Effect of prepartum treatment of vitamin e-selenium on postpartum reproductive and productive performance of exotic cows and their calves under subtropical conditions. *Pak. Vet. J.*, 27: 105-108.
- Sikosana, J.L.N., J. Mangena and T. Vela, 1990. Growth and reproductive performance of young indigenous does given supplementary feed during the dry season. Matopos Research Station, Bulawayo, Annual Report, Government of Zimbabwe, pp, 36-41.
- Tawonezvi, H.P.R. and H.K. Ward, 1987. Productivity of indigenous sheep and goats. 1. Environmental influences on performance of sheep. *Zimbabwe J. Agric. Res.*, 25: 41-49.
- Van Soest, P.J., J.B. Robertson and B.A. Lewis, 1991. Methods for dietary fiber, neutral detergent fiber and non starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.*, 74: 3583-3597.