

Microarray Analysis on the Differences of Gene Expression in Longissimus Dorsi Muscle Tissue Between 1 and 24 Months Chinese Red Steppes

¹Lihong Qin, ²Guoliang Zhang, ²Yang Cao, ¹Jiabao Zhang,
²Yumin Zhao and ¹Zhihui Zhao

¹Keylab of Animal Embryo Engineering,
College of Animal Science and Veterinary Medicine,
Jilin University, 130062 Changchun, P.R. China

²Branch of Animal Science, Jilin Academy of Agricultural Sciences,
136100 Gongzhuling, P.R. China

Abstract: Gene expression in different growth stages showed differences. In order to establish the correlation between gene expression mechanisms and beef traits formation which controlled by polygenes, microarray analysis was performed in Longissimus dorsi muscle tissue between 1 and 24 months Chinese red steppes. About 1,282 (5.6%) probes showed significant differences at the 2 growth stages and 126 genes showing strong correlation with beef traits formation were gained by the GO analysis. With the KEGG analysis, 63 pathways were found to be related to beef traits formation which involved 73 genes. About 28 genes were found in a single pathway while 35 genes were found in 2-16 pathways, respectively. The panel of transcripts and gene pathway analysis in different growth stages may be helpful for the study on beef traits formation and the gene expression profile construction in Longissimus dorsi muscle tissues would also make a model for screening candidate genes which have genetic effect on meat quality in bovine.

Key words: Microarray, gene expression profile, different growth stage, beef traits formation, Chinese red steppe, muscle

INTRODUCTION

Beef traits controlled by polygenes are affected by many factors such as genetic factors, nutrition, age, slaughtering and processing technology but the genetic factors are the most influential. In-depth studies of the molecular mechanisms of beef traits formation are expected to develop. Kee *et al.* (2008) analyzed the correlation between beef transcriptome and 4 different beef traits: Shear Force (SF), Water Holding Capacity (WHC), Cooking Loss (CL) and Loin Eye Area (LEA) in Korean native cattle. The results showed that SF was mainly related to energy metabolism, LEA to rRNA processing, WHC to protein metabolism. Thus energy, protein metabolism and meat quality establishes a causal relationship after the aging process. In Ponsuksili *et al.* (2008), the researcher showed that in 1,279 transcripts with trait correlated expression to WHC, 897 expressions QTL (eQTL) with 104 eQTL coinciding with QTL regions for WHC, 96 transcripts had trans acting and 8 transcripts had cis acting regulation. These results showed that the differences of traits may eventually be linked to genes expressions. Thus, we may predict the

different traits by analyzing gene expression mechanisms. Te Pas *et al.* (2007) showed that 88 pathways of porcine myogenesis are involved by Gene Ontology (GO) and Kyoto Encyclopaedia of Genes and Genomes (KEGG) analysis. In their research, most genes were found in a single pathway, even some were found in up to seven pathways. Combining the pathways and the microarray information, about 21 pathways showed sufficient information content for further analysis. These pathways were related to the regulation of several steps in myogenesis and energy metabolism. The experiment indicates that combining microarray results and biological pathway information will give insight into biological trait formation.

The present study try to relate gene expression changes to beef traits formation at two stages (1 and 24 months of age) by the GO and the KEGG analysis. Knowing these relationships may provide a better understanding of gene expression in beef traits controlled by polygenes. It was necessary to develop method to select candidate genes of beef traits by searching Gene Ontology (GO) and biochemical pathways databases (KEGG).

MATERIALS AND METHODS

Animals and muscle samples: A total of 18 bulls of the same breed (Chinese red steppe) were included in the study; 9 were 1 month old while remaining 9 were 24 months old and all provided by Jilin Academy of Agricultural Sciences. They were maintained in standard conditions and fed with standard diets. The bulls were humanly killed at the slaughter house of the academy and fresh longissimus dorsi muscle tissues were obtained during the process which were immediately frozen in liquid nitrogen and stored at -80°C for microarray analysis.

Total RNA isolation and validation: Total RNA from muscle samples was extracted using Trizol reagent (Invitrogen, USA) and purified using QIAGEN RNeasy Total RNA Isolation kit (QIAGEN, USA). The RNA quantity was assessed by measure the optical density at 260/280 nm using Nanodrop ND-1000 (Nano-Drop Technology, Wilmington, DE) and the integrity of total RNA samples were assessed qualitatively on an agilent bio analyzer by inspection of 18 and 28S rRNA bands after gel electrophoresis.

Microarray target preparations and hybridization: The microarray analysis (probe labeling, hybridization and scanning) was performed using Affymetrix GeneChip Bovine Genome Array (Affymetrix, Santa Clara, CA, USA) following the manufacturer's instruction in the CapitalBio Corporation, Beijing, China. Total RNA (10 µg) or mRNA (0.2 µg) was first reverse transcribed using a T7-Oligo (dT) promoter primer in the first-strand cDNA synthesis reaction. Following RNase H-mediated second-strand cDNA synthesis, the double-stranded cDNA was purified and served as a template in the subsequent *In vitro* Transcription (IVT) reaction. The IVT reaction was carried out in the presence of T7 RNA Polymerase and a biotinylated nucleotide analog/ribonucleotide mix for complementary RNA (cRNA) amplification and biotin labeling. The biotinylated cRNA targets were fragmented in 1 x fragmentation buffer solution provided with the

genechip sample cleanup module (Affymetrix) at 94°C for 35 min. A total of 10 µg of fragmented biotin-labeled cRNA per replicate in hybridization mixture then was hybridized to Bovine Genome Array from Affymetrix GeneChips and incubated overnight at 45°C in Affymetrix GeneChip Hybridization Oven 640, all according to the manufacturer's instructions. The mixture was removed 16 h after hybridization in several cycles; the chips were washed with nonstringent buffer and stained with streptavidin-phycoerythrin antibody solution (Affymetrix) on an automated Affymetrix GeneChip Fluidic Station 450 station. The data were collected by using affymetrix genechip scanner 3000. Microarray images quantified using Affymetrix GeneChip Operating Software (GCOS1.4).

Quantitative real-time RT-PCR: To validate the results of the microarrays, 6 genes (IGF2 (down-regulated), PSPH (down-regulated), IGFBP2 (no significant change), IGFBP5 (no significant change), GPAM (up-regulated), CAPN3 (up-regulated)) showing differential expression were selected and analyzed by real-time RT-PCR. Genes were chosen in different pathways. Primers were designed using Primer Premier5 software (Table 1). Relative mRNA expression levels were determined by using the cDNA for quantitative real-time PCR amplification with SYBR Green I kit (Biotechs, China). Specificity of all individual amplification reactions was confirmed by melting curve analysis. Real-time expression values were calculated by using the relative standard curve method. Standard curves were generated for each mRNA by using 10-fold serial dilutions. The reactions were denatured at 95°C for 5 min followed by 45 cycles at 95°C for 10 sec and 60°C for 30 sec in the ABI PRISM 7500 Sequence detection system (Applied biosystems, USA). Data are reported as values normalized to the housekeeping gene GAPDH. One way ANOVA analysis was performed using SPSS (13.0) software. The significance of differences between the means for each gene was evaluated using LSD or Tamhane's T2 test (for significance levels of $p < 0.05$). All the values are presented as means \pm SEM.

Table 1: Primer pairs used for real-time PCR

Gene symbols	GenBank accession number	Sequence (5'-3')	Product size (bp)
IGF2	NM_174087	F-CCTTCGCCTCGTGCTGCTATG R-GTCGGTTTATGCGGCTGGATG	135
PSPH	NM_001046355	F-TGAGACGCAGCCAACAG R-CCGAATCCAATGAAAGC	150
IGFBP2	NM_174555	F-ACTCCCTACACATCCCCAACTG R-GGATCAGCTTCCCGGTGTTAG	123
IGFBP5	S52657	F-GGTTTGCCCTGAACGAAAAGA R-CTTGGGCGAGTAGGCTCC	111
GPAM	CK977603	F-GATGCCAAGTCTCAAGTTCCT R-CCCTCAATGTGTCTGGGTTTT	244
CAPN3	NM_174260	F-ATGGAGACTGGGTGGACG R-CATTGCGATGGTTGGAT	87

Microarray data analysis: The 6 chips rectified were made by normalization processing using dChip (Affymetrix) software (Li and Wong, 2001a, b). False discovery rates for the genes were calculated by using t-test and p-values. Fold changes were calculated based on unadjusted data's means.

Genes were annotated with NetAffx (Liu *et al.*, 2003) and differentially expressed genes were identified using Significance Analysis of Microarrays (SAM). Using the criterion of cutoff limitation as a fold change ≥ 2 or ≤ 0.5 and q-value $\leq 5\%$, differential expression genes were screened and clustered with the Cluster 3.0 software. The selected genes were further analyzed in the context of Gene Ontology (GO) biological process and KEGG biological pathway using the molecule annotation system 2.0 (MAS 2.0, <http://bioinfo.capitalbio.com/mas3/>) software (CapitalBio, Beijing, China).

RESULTS AND DISCUSSION

Microarray analysis: The Affymetrix GeneChip Bovine Genome Array which contains 24,128 probe sets representing >23,000 transcripts and 19,000 UniGene clusters was used to profile gene expression in the longissimus dorsi muscle of 1 and 24 months old Chinese red steppes.

The microarray hybridization results showed that 22,810 probe sets were used to detect which accounts 94.5% of the total number of probes and 16,321 (71.6% of the total number) probe sets were detected. At the cutoff criteria of the signed fold change ≥ 2 or ≤ 0.5 and $q \leq 5\%$, a total of 1,282 transcripts (5.6% of total probe sets on the array) were differentially expressed at the two stages, 959 (4.2%) of which were up-regulated while 323 (1.4%) were down-regulated (GEO number: GSE21782). About 1,008 differential gene annotations obtained using the molecule annotation system 2.0 (CapitalBio, Beijing, China) among of which 4 gene transcripts are unknown proteins in the functions (0.4%) while 274 of which show no homology to any known sequence.

BLAST analysis revealed that 1,001 probe genes (745 up-regulated and 256 down-regulated) share significant similarity in amino acid sequence with other functional genes accounting for 78.1% of the total number of probe sets. However, 281 (21.9%) probe genes (214 up-regulated and 67 down regulated) did not show similarity with other functional genes (June 2009).

Quantitative real-time RT-PCR: To confirm the microarray data, 6 genes (2 down-regulated genes (IGF2 and PSPH), 2 up-regulated genes (GPAM and CAPN3) and 2 genes with no significant change (IGFBP2 and IGFBP5)) with differentially expressed were selected for quantitative real-time PCR. Absolute quantification method was chosen for each of the 6 genes in the experiment (Tichopad *et al.*, 2003). Linearized plasmid DNA was used to plot a standard curve to quantify absolute gene expression changes and the R^2 for each standard curves was >0.99. The values were presented as means \pm SEM (Table 2). SPSS analysis showed that the expression of IGF2 (Sig., 0.008) and PSPH (Sig., 0.032) are down-regulated, GPAM (Sig., 6.919e-008) and CAPN3 (Sig., 0.004) are up-regulated, IGFBP2 (Sig., 0.808) and IGFBP5 (Sig., 0.748) with no significant change (Fig. 1). These results were exactly consistent with the microarray experiment.

Gene Ontology (GO) analysis: The selected genes with significant expression difference at the 2 stages were analyzed in the context of GO biological process. The GO analysis dynamically assigns different genes to different GO biological process categories and provides a z score for each category.

The z score represents a statistical measurement of relative gene expression dynamic in a given GO term. This approach revealed very interesting phenomena. When we choose GO class of level 0, gene expression fall in 4 Gene ontology categories including molecular functions (464), biological processes (747) ontology categories and cellular components (201) but when it comes to the GO class of level 1, 282 categories were obtained (Fig. 2).

Analysis of the genes may affect beef traits: About 585 genes in 1,282 genes of significant expression difference were provided with gene annotations in relation to cellular component. About 126 genes related to cellular component and metabolism out of 457 genes with definite biological process showed correlation with meat quality traits.

Based on the functional analysis of 126 genes showing correlation with beef traits, we identified a few common biological metabolism categories including glucose (16 genes, 10 up-regulated and 6 down-regulated), lipid (19 genes, 17 up-regulated and 2 down-regulated), protein and amino acid (40 genes, 32 up-

Table 2: Number of copies of mRNA in different genes (copies μ LG¹)

Periods	IGF2	PSPH	IGFBP2	IGFBP5	GPAM	CAPN3
1 month old bull	86676 \pm 11028	747797 \pm 56197	1289908 \pm 141173	1413683 \pm 179104	736994 \pm 27502	58866 \pm 5967
24 months old bull	46014 \pm 5765	518622 \pm 76179	1351482 \pm 217665	1492156 \pm 160146	2246000 \pm 99830	209489 \pm 37816

(p<0.05)

regulated and 8 down-regulated), DNA and RNA (4 genes, 1 up-regulated and 3 down-regulated), energy (9 genes, 6 up-regulated and 3 down-regulated) and others (38 genes, 30 up-regulated and 8 down-regulated) (Table 3). KEGG analysis showed 63 pathways related to metabolism. About 73 genes expressed with significant difference. We also find 28 genes in a single pathway, 35 genes in 2-16 pathways (Table 4).

Meat quality traits were quantitative traits controlled by polygenes. Due to the special characteristic, many of genes could not be clearly established links with traits, although they played an important role. Some of them regulating the expression of other genes could not simply be correlated with in the way of measuring traits and gene expression. Therefore, measurement seemed to be pointless in this case. Thus, new methods were expected to develop.




Fig. 1: The gene expression of IGF2, PSPH, IGFBP2, IGFBP5, GPAM, CAPN3 of 1 and 24 months Chinese red steppe longissimus dorsi muscle in quantitative real-time RT-PCR




Fig. 2: The results of Gene Ontology (GO) analysis

Table 3: Genes of significant differential expression related to beef traits in microarray analysis

Gene ID	Representative public ID	Gene title	Fold changes	q-value (%)
Lipid metabolism (up-regulated)				
Bt.5319.2.S1_a_at	AF080228.1	Peroxisomal protein 6	2.60	0.000
Bt.136.1.S1_at	NM_174210.1	Uncoupling protein 3 (mitochondrial, proton carrier)	3.63	0.000
Bt.3372.2.S1_a_at	BF040491	1-acylglycerol-3-phosphate O-acyltransferase 6 (lysophosphatidic acid acyltransferase, zeta)	2.05	0.150
Bt.19423.2.S1_at	CF923200	ATP-binding cassette, sub-family A (ABC1), member 1	2.57	0.150
Bt.19423.1.S1_at	CB462017	ATP-binding cassette, sub-family A (ABC1), member 1	2.39	0.750
Bt.23550.1.S1_at	NM_205793.1	Diacylglycerol O-acyltransferase homolog 2 (mouse)	2.37	0.180
Bt.5467.1.S1_at	NM_174161.2	Prosaposin	3.86	0.000
Bt.5467.1.S2_a_at	CK848991	Prosaposin	2.11	0.000
Bt.5467.2.S1_a_at	BM365269	Prosaposin	2.11	0.000
Bt.5467.1.S2_at	CK848991	Prosaposin	2.11	0.068
Bt.29570.1.S1_at	CK977603	Glycerol-3-phosphate acyltransferase, mitochondrial	2.72	0.150
Bt.21724.2.S1_a_at	CK982401	Farnesyl-diphosphate farnesyltransferase 1	4.52	0.000
Bt.21724.1.S1_at	CK961813	Farnesyl-diphosphate farnesyltransferase 1	2.31	0.120
Bt.997.2.S1_a_at	BM986149	Cytochrome b5 reductase 1	2.57	0.000
Bt.997.1.S1_at	CK774098	Cytochrome b5 reductase 1	2.04	0.000
Bt.22590.1.S1_at	CB435331	1-acylglycerol-3-phosphate O-acyltransferase 2 (lysophosphatidic acid acyltransferase, beta)	2.28	0.000
Bt.12805.1.S1_at	AJ299408.1	LAMA-like protein 1	3.59	0.000
Lipid metabolism (down-regulated)				
Bt.22672.1.A1_at	CK845888	Hydroxyprostaglandin dehydrogenase 15-(NAD)	0.26	0.000
Bt.13315.1.S1_at	CK771506	Hypothetical protein LOC768237	0.34	0.210
Protein and amino acid metabolism (up-regulated)				
Bt.6774.2.S1_at	CK951538	Microtubule-associated protein 1 light chain 3 beta	2.10	0.150
Bt.6774.1.S1_at	CB169021	Microtubule-associated protein 1 light chain 3 beta	2.03	0.300
Bt.23034.2.S1_at	CK771675	SMT3 suppressor of mif two 3 homolog 3 (<i>S. cerevisiae</i>)	2.22	0.000
Bt.20330.1.S1_at	CK773920	Protease, serine, 23	6.00	0.000
Bt.8947.1.S1_at	NM_174101.2	Legumain	2.19	0.070
Bt.13730.2.S1_at	AV595619	ATG4 autophagy related 4 homolog A (<i>S. cerevisiae</i>)	3.03	0.150
Bt.13730.1.A1_at	BF775534	ATG4 autophagy related 4 homolog A (<i>S. cerevisiae</i>)	2.14	0.150
Bt.15675.1.S1_at	NM_181667.1	ADAM metalloproteinase with thrombospondin type 1 motif, 4	2.15	0.750
Bt.24474.1.S1_at	BF429904	AFG3 ATPase family gene 3-like 2 (yeast)	3.30	0.000
Bt.1657.1.A1_at	CK963042	AFG3 ATPase family gene 3-like 2 (yeast)	2.02	1.520
Bt.11239.3.S1_at	BM287651	Spastic paraplegia 7 (pure and complicated autosomal recessive)	2.12	0.150
Bt.3961.2.S1_a_at	AF148714.1	Calpain 3 (p94)	3.51	0.000
Bt.22536.1.S1_at	BI849209	Potassium channel modulatory factor 1	3.14	0.000
Bt.22536.2.S1_at	AV613194	Potassium channel modulatory factor 1	2.63	0.000
Bt.21166.1.S1_at	CK949278	F-box protein 25	2.56	0.000
Bt.21681.1.A1_at	CK777827	F-box and leucine-rich repeat protein 4	3.10	0.000
Bt.21681.2.S1_at	CB532552	F-box and leucine-rich repeat protein 4	4.50	0.000
Bt.18003.1.S1_at	CB437779	Cullin 3	5.52	0.000
Bt.6455.1.S2_at	CB534413	Cullin 4A	2.14	0.000
Bt.6455.1.S1_at	CK772096	Cullin 4A	3.09	0.000
Bt.28873.1.S1_at	CK943939	Cullin 2	2.35	0.000
Bt.3528.2.S1_at	BG359079	Cullin 3	2.41	0.070
Bt.20830.1.A1_at	CB531128	Cullin 5	2.27	0.120
Bt.14158.1.S1_at	BG358477	Cullin 5	2.49	0.120
Bt.5161.1.S1_at	CK951722	Similar to ubiquitin specific protease 24	2.15	0.150
Bt.23662.2.S1_at	CK956837	Ring finger protein 34	2.03	1.080
Bt.6796.1.A1_at	BF440255	Ring finger protein 144B	3.21	0.000
Bt.29888.1.S1_at	BF041956	Cullin 1	3.56	0.000
Bt.15790.2.S1_at	CF615274	Ankyrin repeat and SOCS box-containing 8	2.85	0.000
Bt.27401.1.A1_at	CK849322	Ring finger protein 128	4.36	0.000
Bt.19533.1.A1_at	CB464082	Glutamate-cysteine ligase, modifier subunit	2.54	0.000
Bt.436.1.S1_at	NM_175811.2	Aldehyde dehydrogenase 6 family, member A1	2.19	0.000
Protein and amino acid metabolism (down-regulated)				
Bt.5313.1.S1_at	NM_174745.2	Matrix metalloproteinase 2 (gelatinase A, 72kDa gelatinase, 72kDa type IV collagenase)	0.49	2.390
Bt.5096.1.S1_at	CK849662	Chaperonin containing TCP1, subunit 3 (gamma)	0.45	2.020
Bt.6427.1.S1_at	CB451598	Chaperonin containing TCP1, subunit 4 (delta)	0.49	4.110
Bt.5397.1.S1_at	NM_174465.2	SEC11 homolog A (<i>S. cerevisiae</i>)	0.48	0.210
Bt.2120.1.S1_at	AW485920	Plasma glutamate carboxypeptidase	0.49	0.120
Bt.16188.1.S1_at	CK945725	ADAM metalloproteinase with thrombospondin type 1 motif, 5	0.38	1.080
Bt.23218.1.S1_at	CK849357	Cathepsin K	0.27	2.390
Bt.8549.1.S2_at	CK940477	Speckle-type POZ protein	0.07	2.020

Table 3: Continued

Gene ID	Representative public ID	Gene title	Fold changes	q-value (%)
Glucose metabolism (up-regulated)				
Bt.4857.2.S1_at	BG687989	Prune homolog (Drosophila)	2.15	0.000
Bt.20931.1.S1_at	CF765643	Protein phosphatase 1, catalytic subunit, beta isoform	4.58	0.000
Bt.20281.3.S1_a_at	BE808500	Phosphoglucosyltransferase 1	3.03	0.000
Bt.562.1.S1_at	NM_174572.2	6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 1	2.08	0.150
Bt.8679.3.S1_at	CF613505	Dihydrolipoamide S-acetyltransferase	4.74	0.000
Bt.18533.1.S1_at	CK846562	Activating transcription factor 3	4.74	0.150
Bt.19161.1.S1_at	AV601057	Activating transcription factor 4 (tax-responsive enhancer element B67)	2.49	0.000
Bt.2347.2.S1_at	AV615616	Phosphofructokinase, muscle	7.08	0.000
Bt.19415.1.A1_at	CB461885	Pyruvate dehydrogenase (lipoamide) alpha 1	6.06	0.000
Bt.1129.1.S1_at	CK967288	Citrate synthase	4.06	0.000
Glucose metabolism (down-regulated)				
Bt.7251.1.S1_at	BE477050	UDP-Gal:betaGlcNAc beta 1,3-galactosyltransferase, polypeptide 2	0.27	4.110
Bt.5131.1.S1_at	CK771744	Chitinase domain containing 1	0.36	0.000
Bt.1798.1.S1_at	AF098512.1	Mannosidase, beta A, lysosomal	0.48	2.390
Bt.24309.1.A1_at	CB449936	Lymphatic vessel endothelial hyaluronan receptor 1	0.31	0.450
Bt.23267.1.S1_at	NM_205815.1	Lymphatic vessel endothelial hyaluronan receptor 1	0.44	1.520
Bt.5878.2.S1_at	BM258870	Serine dehydratase	0.49	2.390
Energy metabolism (up-regulated)				
Bt.16782.2.S1_at	BM482702	Adenylosuccinate synthase like 1	4.00	0.000
Bt.5848.2.S1_at	CF614616	ATPase, Ca++ transporting, cardiac muscle, fast twitch 1	2.76	0.150
Bt.8872.1.S1_at	NM_174696.2	ATPase, Ca++ transporting, plasma membrane 1	2.46	0.000
Bt.16296.1.A1_at	CB453612	ATPase, Ca++ transporting, cardiac muscle, slow twitch 2	18.30	0.000
Bt.26179.1.S1_at	CK776897	ATPase, (Na+)/K+ transporting, beta 4 polypeptide	2.10	1.520
Bt.21546.1.S1_at	AW429997	Similar to ATPase, Class II, type 9A	2.15	0.000
Energy metabolism (down-regulated)				
Bt.20852.1.A1_at	CB468425	GNAS complex locus	0.44	0.210
Bt.5340.1.S1_s_at	CK770903	Non-metastatic cells 1, protein (NM23A) expressed in	0.36	0.000
Bt.1582.1.S1_at	CK972542	Non-metastatic cells 2, protein (NM23B) expressed in	0.45	0.000
DNA and RNA metabolism (up-regulated)				
Bt.28559.1.S1_a_at	CK944988	Deoxyribonuclease I-like 1	3.07	
DNA and RNA metabolism (down-regulated)				
Bt.15601.1.A1_at	CB446288	X-ray repair complementing defective repair in Chinese hamster cells 3	0.45	0.750
Bt.5092.1.S1_at	AV614758	Similar to PRDX3 protein /// similar to PRDX3 protein /// peroxiredoxin 3	0.45	1.080
Bt.14038.1.A1_at	CB227291	5', 3'-nucleotidase, cytosolic	0.38	4.110
Other metabolism (up-regulated)				
Bt.4317.1.S1_at	NM_174076.2	Glutathione peroxidase 1	2.62	0.000
Bt.23174.1.S1_at	NM_174799.1	CD74 molecule, major histocompatibility complex, class II invariant chain	2.09	0.560
Bt.26692.1.S1_a_at	CK774343	BCL2/adenovirus E1B 19kDa interacting protein 3	2.99	0.070
Bt.6958.1.A1_at	CK955033	BCL2/adenovirus E1B 19kDa interacting protein 3	2.51	1.520
Bt.3349.1.S1_at	AV615323	Citrate lyase beta like	2.09	0.000
Bt.2737.1.S1_at	CK771684	Carbonic anhydrase III, muscle specific	2.96	0.000
Bt.5152.1.S1_at	NM_174054.2	Ferrochelatase (protoporphyrin)	2.19	0.150
Bt.15857.1.S1_at	CK848612	Aminolevulinic acid, synthase 1	2.88	0.000
Bt.26688.2.S1_a_at	BF429567	FERM domain containing 3	3.25	0.000
Bt.28907.1.S1_at	CK945732	NADH dehydrogenase (ubiquinone) Fe-S protein 1, 75kDa (NADH-coenzyme Q reductase)	3.22	0.150
Bt.28243.1.S1_a_at	CK951402	Vanin 1	2.38	0.150
Bt.28243.2.S1_at	BP106249	Vanin 1	2.07	0.560
Bt.1059.3.S1_a_at	CK945537	ATPase, Ca++ transporting, cardiac muscle, slow twitch 2	2.05	0.000
Bt.18840.1.S1_at	CK846230	Adenylate cyclase 2 (brain)	2.04	0.000
Bt.28878.1.S1_at	CK944325	Aspartoacylase (<i>Canavan disease</i>)	2.56	0.000
Bt.23551.2.S1_at	BM967523	N-acetyltransferase 13 (GCN5-related)	2.24	0.300
Bt.23551.3.S1_at	CB535034	N-acetyltransferase 13 (GCN5-related)	2.19	0.450
Bt.22170.1.S1_at	BE505063	1-acylglycerol-3-phosphate O-acyltransferase 5 (lysophosphatidic acid acyltransferase, epsilon)	2.33	0.000
Bt.22170.1.S1_a_at	BE505063	1-acylglycerol-3-phosphate O-acyltransferase 5 (lysophosphatidic acid acyltransferase, epsilon)	2.37	0.000
Bt.12211.1.S1_at	CB165818	Peroxisomal D3,D2-enoyl-CoA isomerase	2.19	0.070
Bt.22170.2.A1_at	CB454188	1-acylglycerol-3-phosphate O-acyltransferase 5 (lysophosphatidic acid acyltransferase, epsilon)	2.10	0.150
Bt.9507.1.S1_at	BM365221	Haloacid dehalogenase-like hydrolase domain containing 3	2.05	0.250
Bt.29926.2.S1_at	CK961230	Transmembrane protein 68	2.07	0.000

Table 3: Continued

Gene ID	Representative public ID	Gene title	Fold changes	q-value (%)
Bt.4732.1.S1_at	NM_174239.2	Aldehyde dehydrogenase 1 family, member A1	2.01	0.750
Bt.16325.1.A1_at	CK847813	Glutaryl-Coenzyme A dehydrogenase	2.02	0.000
Bt.9777.1.S1_at	CB172625	3-hydroxybutyrate dehydrogenase, type 1	2.02	1.080
Bt.4659.1.S1_at	CK953255	Aldehyde dehydrogenase 2 family (mitochondrial)	2.78	0.000
Bt.5530.1.S1_at	NM_174180.2	Dehydrogenase/reductase (SDR family) member 3	2.49	0.000
Bt.14570.1.A1_at	CB165420	4-aminobutyrate aminotransferase	5.14	0.000
Bt.14570.2.S1_at	AW307709	4-aminobutyrate aminotransferase	3.75	0.000
Other metabolism (down-regulated)				
Bt.546.1.S1_at	NM_174680.2	Potassium large conductance calcium-activated channel, subfamily M, alpha member 1	0.26	2.020
Bt.9047.1.S1_at	CK847263	D-dopachrome tautomerase	0.47	0.170
Bt.3890.1.S1_at	NM_174674.2	Tumor necrosis factor receptor superfamily, member 1A	0.48	1.520
Bt.23171.2.S1_at	CK774607	Pterin-4 alpha-carbinolamine dehydratase/dimerization cofactor of hepatocyte nuclear factor 1 alpha	0.23	3.100
Bt.7490.1.A1_at	CK778231	Methylenetetrahydrofolate dehydrogenase (NADP+ dependent) 1-like	0.19	0.000
Bt.4897.1.S1_at	NM_174094.2	inhibin, alpha	0.09	0.080
Bt.11069.1.S1_at	CK771476	Adenylate cyclase 6	0.48	0.750
Bt.18945.1.A1_at	CB454235	Pleckstrin homology domain containing, family G (with RhoGef domain) member 1	0.39	0.080

Table 4: The KEGG analysis on genes of significant differential expression related to beef traits in microarray analysis

N*	Pathways with N gene(s) of significant differential expression(representative public ID)
1	Methane metabolism (AF080228.1), Alkaloid biosynthesis II (AF080228.1), Inositol metabolism (NM_175811.2), 3-Chloroacrylic acid degradation (CK953255), Ascorbate and aldarate metabolism (CK953255), Limonene and pinene degradation (CK953255), Synthesis and degradation of ketone bodies (CB172625), Glyoxylate and dicarboxylate metabolism (CK967288), Glycosphingolipid biosynthesis – lactoseries (BE477050), Benzoate degradation via CoA ligation (CK847813), N-Glycan degradation (AF098512.1), Glycosphingolipid biosynthesis – ganglioseries (BP102248), Bile acid biosynthesis (CK953255), Glycosylphosphatidylinositol(GPI)-anchor biosynthesis (CK729543), Nicotinate and nicotinamide metabolism (CB227291), Galactose metabolism (AV615616), Glycan structures – degradation (AF098512.1), Pentose phosphate pathway (AV615616), Carbon fixation (CK770445), Starch and sucrose metabolism (CK846492), Ether lipid metabolism (BF040491), Proteasome (CK957649), Citrate cycle (TCA cycle) (CK967288), O-Glycan biosynthesis (NM_177519.2), Arginine and proline metabolism (NM_173892.2), Arachidonic acid metabolism (NM_174076.2), N-Glycan biosynthesis (CK972901), Glycan structures - biosynthesis 1 (NM_177519.2)
2	Valine, leucine and isoleucine biosynthesis (CK775511, CB461885), Lysine degradation (CK847813, CK953255), Terpenoid biosynthesis (CK961813, CK982401), Vitamin B6 metabolism (CB166901, CF763176), Pantothenate and CoA biosynthesis (CK951402, BP106249), Glycerophospholipid metabolism (BF040491, CK977603), Glutathione metabolism (NM_174076.2, CB464082), Fatty acid metabolism (CK847813, CK953255), Aminoacyl-tRNA biosynthesis (CK775511, CB534569), Tyrosine metabolism (NM_177944.1, CK775534), Drug metabolism-cytochrome P450 (NM_177944.1, CK775534)
3	Phenylalanine metabolism (AF080228.1, NM_177944.1, CK775534), Porphyrin and chlorophyll metabolism (CK770847, CK848612, NM_174054.2), beta-Alanine metabolism (CB165420, AW307709, CK953255), Glutamate metabolism (CB165420, AW307709, CB464082), Pyruvate metabolism (CK953255, CB461885, CK770445), Biosynthesis of steroids (CK969564, CK961813, CK982401), Retinol metabolism (NM_205793.1, NM_174239.2, NM_174180.2), Glycolysis / Gluconeogenesis (AV615616, CK953255, CB461885), Glycan structures - biosynthesis 2 (CK729543, BE477050, BP102248)
4	Histidine metabolism (NM_177944.1, CK775534, CK944325, CK953255), Propanoate metabolism (CB165420, AW307709, CK953255, NM_175811.2), Glycerolipid metabolism (NM_205793.1, BF040491, CK953255, CK977603), Urea cycle and metabolism of amino groups (NM_177944.1, CK775534, CK953255, NM_173892.2), Fructose and mannose metabolism (AV615616, NM_174812.2, NM_174434.2, NM_174572.2), Aminosugars metabolism (CK774098, BM986149, CK982168, BM967525)
5	Alanine and aspartate metabolism (CB165420, AW307709, CK944325, CB461885, NM_173892.2), Butanoate metabolism (CB172625, CB165420, AW307709, CK953255, CB461885), Valine, leucine and isoleucine degradation (CB165420, AW307709, CK982441, CK953255, NM_175811.2), Pyrimidine metabolism (CK770903, NM_174428.2, CK730171, BM435937, CB227291)
6	Tryptophan metabolism (CK847813, NM_177944.1, CK775534, CB534569, CK953255, CK849867), Oxidative phosphorylation (NM_176676.1, NM_175807.1, CK945732, CK775109, CB538935, M38520.1)
8	Glycine, serine and threonine metabolism (CK772398, AW478353, CB166901, CF763176, CK848612, CK771294, NM_177944.1, CK775534)
10	Purine metabolism (CB439779, CK770903, NM_174428.2, CB167755, CK983189, CK846230, BM435937, CK773933, CB227291, CK771476)
134	Ubiquitin mediated proteolysis (BG358233, BM251234, CK974661, BF041956, CK943939, CB437779, BG359079, BG358477, CB531128, CB534413, CK772096, CB419023, CB530307, BF039092)

*N = The number of genes with significant differential expression

In this study, microarray analysis was carried out to profile gene expression in Chinese red steppes Longissimus dorsi muscle at two different stages: 1 and 24 months old (the experiments have been achieved recognition of GEO and successfully obtained access number. Fortunately, 1,282 (5.6%) probes showed significant differences between the two growth stages in bovine and 126 genes among of which showing strong

correlation with beef traits' formation were gained by GO and KEGG analysis. These genes related to various metabolisms may be the reason which accounts for the differences of beef traits between the two stages. If we put the cutoff limitation scale down to a fold change ≥ 1.5 or ≤ 0.67 and q-value $\leq 5\%$, 2,923 (12.8%) probe sets were selected but in the case of the majority, the criterion of cutoff limitation may be not strict enough.

We also analyzed several genes which thought to be important and influential beef traits such as those encode Ca^{2+} /calmodulin-dependent protein Kinases (CaMKs), myogenic regulatory factors (MRFs), Fatty Acid-binding Proteins (FABPs) and Peroxisome Proliferator-activated Receptors (PPARs). Calpains are a family of calcium activated cysteine proteases that are widely distributed in cell cytoplasm and sub-organelles.

They go along with their endogenous inhibitor, calpastatin. In calpain families, calpain 1-10 are in relation to tenderization of meat (Kar *et al.*, 2010). In this study, calpain 1 (fold change = 0.97, q-value = 46.28%) has no significant changes. This indicates that calpain 1 may be unresponsive to the differences of beef traits in the experiment. Meanwhile, the experiment shows that Calpastatin (fold change = 2.13, q-value = 0.07%) was significantly up-regulated which indicates that the differences may be due to Calpastatin. The experiment also shows that calpain 3 was significantly up-regulated but in literature there are controversies on whether calpain3 has any effect on tenderization of meat. Correlation analysis between the relative rate of tenderization and mRNA expression of calpains revealed a strong relationship with calpain 3 in both species, bovine and ovine [r (bovine) = 0.522, r (ovine) = 0.706] (Ilian *et al.*, 2001). In contrast, Parr *et al.* (1999) did not find a link between calpain 3 and postmortem tenderization in the porcine Longissimus Thoracis et lumborum (LT).

The discrepancy between them may be due to the methodology or animal species. In addition in this research, the calpain 10 also show no significant changes. Evidence is mounting that the Ca^{2+} /calmodulin-dependent Kinases (CaMK) II and IV play important role in regulating oxidative enzyme expression, mitochondrial biogenesis and expression of fibre-type specific myofibrillar proteins.

Examination of Ca^{2+} -dependent and Ca^{2+} -independent CaMK II activity in chicken anterior latissimus dorsi muscle sample after 7 days of stretch overload and 2 weeks of voluntary wheel running have shown that stretch overload and wheel running result in a 122% increase in muscle protein content and are associated with 47% increase in Ca^{2+} -independent CaMKII activity (Fluck *et al.*, 2000).

Results indicate that CaMK IIg not CaMK IIa or -b is up-regulated in aged and denervated soleus muscle and CaMK IV is absent in skeletal but not cardiac muscle (Chin, 2004). Similarly this study showed that CaMK Iid (fold change = 2.97, q-value = 0) and CaMK IIg (fold change = 3.06, q-value = 0) have significant changes in contrast with CaMK IIa, CaMK IIb or other members of

CaMKs. Thus, the differences in beef traits may be related to the regulations of CaMK Iid or CaMK IIg.

The MyoD gene family members encoding basic helixloop-helix (bHLH) transcription factors (MyoD, myogenin, Myf5 and MRF4) contain one or two transactivation domains (at the N- and C-terminus). They are able to make nonmuscle cells to have muscle phenotype in culture (Wei and Paterson, 2001). It is reported that MRFs have relation to tenderness, Loin Eye Area (LEA) and other beef traits (Kim *et al.*, 2009; Liu *et al.*, 2008; Zhu and Li, 2005), however the experiment showed that MyoD, MyoG, Myf6 and Myf5 have no significant changes. For MyoD (fold change = 1.33, q-value = 5.31%), although the q-value (5.31%) was in accordance with the criterion of cutoff limitation, the fold change (1.33) was not within the scope of screening.

So, it showed that the result of gene expression comparison has no significant difference even if the relaxation of screening criteria to fold change is relaxed to ≥ 1.5 or ≤ 0.67 and q-value $\leq 5\%$. So, the MyoD gene expression showed that differences in beef traits may be unrelated to regulation of MRFs.

Intramuscular fat content (IMF) was correlated with meat quality traits and showed significant effects on three aspects of meat: tenderness, flavor and juicy (De Vol *et al.*, 1988). Fatty Acid-binding Proteins (FABPs) are small intracellular proteins involved in fatty acid transport from the plasma membrane to the sites of β oxidation and/or triacylglycerol or phospholipid synthesis (Veerkamp and Maatman, 1995). The experiment showed that FABPs have no significant changes even the relaxation of screening criteria to fold change is relaxed as above. So, the IMF expression results showed that the differences in beef traits may be unrelated to the regulation of FABPs.

The Peroxisome Proliferator-activated Receptors (PPARs) are members of the nuclear receptor superfamily and play an important regulatory role in lipid metabolism (Latruffe and Vamecq, 1997). The PPAR gamma (PPARG) subtype is a key transcription factor that controls adipocyte differentiation and fat deposition in mammals but in this experiment PPARG (fold change = 0.96, q-value = 46.28%) as well as other members of this family have no significant changes. The PPAR expression results showed that the differences in beef traits may be unrelated to the regulation of PPARs.

CONCLUSION

In this study, we have shown several CaMKs encoded genes play an important role in beef traits using Affymetrix GeneChip Bovine Genome Array while genes

encoding MRFs, FABPs and PPARs do not have influential impact on beef traits' formation. We also found, it was an effective way to exclude invalid genetic pathways when the subject (beef traits in the model) was controlled by various genes. It is no doubt that we need to improve the efficiency of searching for candidate genes. At the same time, we also see the interaction of gene pathways and analyze the mechanisms of gene expression may provide some new insight in regulatory network characters of quantitative traits controlled by polygenes and the result will provided in due course.

ACKNOWLEDGEMENTS

We thank Melvin Korkor (Jilin University, P.R. China) and Naizhong Cui (Dalian University of Technology, P.R. China) for critical reading the manuscript, Yonghe Ma (Jilin University, P.R. China) for technical help and Supertech (Enuo Technology, P.R. China) for revision. This research was supported by the 863 hightech project (2008AA101010) and the National Scientific and Technological Supporting Project (2007BAD55B01).

REFERENCES

- Chin, E.R., 2004. The role of calcium and calcium/calmodulin-dependent kinases in skeletal muscle plasticity and mitochondrial biogenesis. *Proc. Nutr. Soc.*, 63: 279-286.
- De Vol, D.L., F.K. McKeith, P.J. Bechtel, J. Novakofski, R.D. Shanks and T.R. Carr, 1988. Variation in composition and palatability traits and relationships between muscle characteristics and palatability in a random sample of pork carcasses. *J. Anim. Sci.*, 66: 385-395.
- Fluck, M., M.N. Waxham, M.T. Hamilton and F.W. Booth, 2000. Skeletal muscle Ca(2+)-independent kinase activity increases during either hypertrophy or running. *J. Applied Physiol.*, 88: 352-358.
- Ilian, M.A., J.D. Morton, M.P. Kent, C.E. le Couteur, J. Hickford, R. Cowley and R. Bickerstaffe, 2001. Intermuscular variation in tenderness: association with the ubiquitous and muscle-specific calpains. *J. Anim. Sci.*, 79: 122-132.
- Kar, P., K. Samanta, S. Shaikh, A. Chowdhury, T. Chakraborti and S. Chakraborti, 2010. Mitochondrial calpain system: An overview. *Arch. Biochem. Biophys.*, 495: 1-7.
- Kee, H.J., E.W. Park and C.K. Lee, 2008. Characterization of beef transcripts correlated with tenderness and moisture. *Mol. Cells*, 25: 428-437.
- Kim, J.M., B.D. Choi, B.C. Kim, S.S. Park and K.C. Hong, 2009. Associations of the variation in the porcine myogenin gene with muscle fibre characteristics, lean meat production and meat quality traits. *J. Anim. Breed. Genet.*, 126: 134-141.
- Latruffe, N. and J. Vamecq, 1997. Peroxisome proliferators and peroxisome proliferator activated receptors (PPARs) as regulators for lipid metabolism. *Biochimie*, 79: 81-94.
- Li, C. and W.H. Wong, 2001a. Model-based analysis of oligonucleotide arrays: Model validation, design issues and standard error application. *Genome. Biol.*, Vol. 2, No. 8.
- Li, C. and W.H. Wong, 2001b. Model-based analysis of oligonucleotide arrays: Expression index computation and outlier detection. *Proc. Natl. Acad. Sci. USA.*, 98: 31-36.
- Liu, G., A.E. Loraine, R. Shigeta, M. Cline and J. Cheng *et al.*, 2003. NetAffx: Affymetrix probesets and annotations. *Nucleic Acids Res.*, 31: 82-86.
- Liu, M., J. Peng, D.Q. Xu, R. Zheng and F.E. Li *et al.*, 2008. Association of MYF5 and MYOD1 gene polymorphisms and meat quality traits in Large White x Meishan F2 pig populations. *Biochem. Genet.*, 46: 720-732.
- Parr, T., P.L. Sensky, G.P. Scothern, R.G. Bardsley, P.J. Buttery, J.D. Wood and C. Warkup, 1999. Relationship between skeletal muscle-specific calpain and tenderness of conditioned porcine longissimus muscle. *J. Anim. Sci.*, 77: 661-668.
- Ponsuksili, S., E. Jonas, E. Murani, C. Phatsara and T. Srikanchai *et al.*, 2008. Trait correlated expression combined with expression QTL analysis reveals biological pathways and candidate genes affecting water holding capacity of muscle. *BMC Genomics.*, 9: 367-367.
- Te Pas, M.F., I. Hulsegge, A. Coster, M.H. Pool, H.H. Heuven and L.L. Janss, 2007. Biochemical pathways analysis of microarray results: Regulation of myogenesis in pigs. *BMC Dev. Biol.*, 7: 66-66.
- Tichopad, A., M.W. Pfaffl and A. Didier, 2003. Tissue-specific expression pattern of bovine prion gene: Quantification using real-time RT-PCR. *Mol. Cell Probes*, 17: 5-10.
- Veerkamp, J.H. and R.G. Maatman, 1995. Cytoplasmic fatty acid-binding proteins: Their structure and genes. *Prog. Lipid. Res.*, 34: 17-52.
- Wei, Q. and B.M. Paterson, 2001. Regulation of MyoD function in the dividing myoblast. *FEBS Lett.*, 490: 171-178.
- Zhu, L. and X.W. Li, 2005. The genetic diversity and genetic effects of Myf-6 gene in different pig breeds. *Yi Chuan*, 27: 887-892.