

Linalool Improve Biochemical Damage and Fatty Acids Composition of Testes on Fasting Male Rats

¹Alpaslan Dayangac, ²Muammer Bahsi, ³Ahmet Ozkaya and ⁴Okkes Yilmaz

¹Department of Biology, Faculty of Art and Science, Ahi Evran University, 40100 Kirsehir, Turkey

²Department of Science Teacher, Faculty of Education, Firat University, 23169 Elazig, Turkey

³Department of Chemistry, Faculty of Art and Science, Adiyaman University, Adiyaman, Turkey

⁴Department of Biology, Faculty of Science, Firat University, 23119 Elazig, Turkey

Abstract: The aim of this study was to evaluate fatty acids compositions, oxidative stress levels and some antioksidant levels in the testes of 24 h fasted rats and linalool application. Adult male Wistar albino rats were assigned as Control (C), 24 h fasted (24 h F) group and added linalool (120 mg kg⁻¹) during 24 h fasted (24 h F+L) groups. After applications, the animals were killed and the testes were excised for determination of tissue malondialdehyde, Glutathione Peroxidase (GSH-Px), the activity of superoxide dismutase and determination of tissue fatty acids compositions. The activities of Superoxide Dismutase (SOD), Glutathione peroxidase (GSH-Px) and the level of Malondialdehyde (MDA) in testicular tissue specimens were determined spectrophotometrically. The fatty acid compositions in the testes were analyzed as percent by gas chromatography (GC). Researchers calculated between C16:0 (palmitic acid) and C22:6 (docosahexaenoic acid) fatty acids in the testes of all experimental groups. Treatment with fasted rats resulted in a significant reduction in C16:1 (palmitoleate) fatty acids when compared to the control rats (p<0.05). C16:1 fatty acids were higher in linalool application than fasted groups (p<0.05). C22:4 (docosatetraenoic Acid), C22:5 (docosapentaenoic acid), C22:6 n3 (docosahexaenoic acid) fatty acids were lower in the fasted group than the control group. In addition, C22:4, C22:5, C22:6 fatty acids compositions in the linalool treatment group were higher than the fasted group (p<0.05). Saturated fatty acids levels in all experiment groups were not change as statistically (p>0.05). SOD and GSH-Px levels in the fasted rats were statistically lower than control groups (p<0.05). Both SOD and GSH-Px levels in the linalool application groups were not statistically a change compared to fasted groups (p>0.05). In addition, MDA levels in the fasted groups were higher than control groups. Besides, MDA levels in the linalool application groups were lower compared to fasted groups (p<0.05). These findings indicate that fasted state has oxidative effects on testicular tissue and linalool has protective effects on male reproductive system.

Key words: Linalool, fasting, testes, rat, fatty acids, SOD, MDA, GSH-Px, oxidative stress

INTRODUCTION

Linalool is a fragrance ingredient used in decorative cosmetics, fine fragrances, shampoos, toilet soaps and other toiletries as well as in non-cosmetic products such as household cleaners and detergents. Its worldwide use is in the region >1000 metric tonnes per annum (Lapczynski *et al.*, 2008; Bickers *et al.*, 2003). Linalool, a monoterpene alcohol is also present in essential oils of many medicinal and aromatic plants that are endowed with many biological activities, including antioxidant, antimicrobial, antiviral, anti-inflammatory and antitumor properties (Tepe *et al.*, 2004; Pattnaik *et al.*, 1997; Mitic-Culafic *et al.*, 2009; Koroch *et al.*, 2007; Celik and Ozkaya, 2002). Besides, Linalool possesses several

depressant effects on the central nervous system and some finding is important in showing the potential usefulness of linalool as a pharmacotherapeutic agent (Leal-Cardoso *et al.*, 2010).

Fasting states occur oxidative stress in some animal tissue. In several studies, lipid peroxidation, nitric oxide synthase activity and hydrogen peroxide accumulation have been reported as a result of fasted. Some researchers have indicated that total Glutathione (GSH), mitochondrial GSH and the activities of major antioxidant enzymes were lower in the fasted mice (Abdelmegeed *et al.*, 2009; Brown *et al.*, 1995). To protect the integrity of biological membranes from detrimental oxidative processes caused by free radicals both enzymatic and non-enzymatic defense mechanisms are present in the cell (Droge, 2002;

Kara *et al.*, 2007). All animal cells have intense phospholipid layer in both cell membrane and organelle membrane. These phospholipids contain PUFA (Polyunsaturated Fatty Acids) and they are directly effected by reactive oxygen species (Bourre *et al.*, 1992; Halliwell, 1994). α -linolenic acid, Eicosapentaenoic Acid (EPA) and Docosahexaenoic acid (DHA) are known as omega-3 fatty acids and these fatty acids levels are effected by oxidative stress state (Lonergan *et al.*, 2002). Testes which are the source of fertility and heredity in male mammals are particularly sensitive feeding (Santos *et al.*, 2004; Conn, 1986). Secretion of testosterone is impaired due to excessive oxidative stress and the degeneration of Leydig cells (Santos *et al.*, 2004; Bandyopadhyay *et al.*, 1999). Lipid peroxidation that is caused by reduced oxygen directly effect the spermatozoon membrane where contain polyunsaturated fatty acids (Sikka, 1996). In this study, Researchers investigated the fatty acids changes, some antioxidant enzyme and lipid peroxidation both fasting state and the linalool application to fasting on rat testes.

MATERIALS AND METHODS

Animals and experimental application groups: In this study, 24 adult male Wistar albino (Firat University Experimental Research Center, Turkey) rats at 12 weeks of age were used (in the each group, n = 8/group). The animals were group housed in cages in a climate-controlled room with a 12 h light/12 h dark cycle and in room temperature (20±3°C) has been provided. Animals which have been used in the study are obtained from Elazig-Turkey Animal Health and Research Institute. During experiment, criteria which are determined by NIH (National Institutes of Health) are obeyed conscientiously as related with animal rights. Animals were randomly divided into three groups. The first group consisted of normally-fed controls, the second group was fasted for 24 h, the third group was fasted for 24 h+linalool (120 mg kg⁻¹). Drinking water was given to fasting animal groups *ad libitum*. Rats in the second group were not given any food for 24 h while rats in the third group were given linalool during 24 h fasting. Linalool was given at the first 12 h fasting. At the end of the experiment, the animals were anesthetized by intramuscular (i.m.) injection of ketamine (70 mg kg⁻¹) (Ketalar, Parke-Davis, Eczacibasi, Istanbul, Turkey) and xylazine (5 mg kg⁻¹) (Rompun, Bayer, Istanbul, Turkey) and blood was drawn from the heart by cardiac puncture and the investigated testes tissues were removed from animals and frozen until analyzed.

Chemicals: Linalool were purchased from Sigma (St. Louis, MO). All the other chemicals used in the study were of analytical grade and were purchased from Merck (Darmstadt, Germany).

Extraction of lipids and preparation of fatty acid methyl esters:

The lipids of testes tissue samples were extracted by the method of Hara and Radin (Hara and Radin, 1978). Testes tissue samples (0.5 g) were homogenized in 5 mL hexane/isopropanol mixture at 3:2 (v/v) for 30 sec. Samples were centrifuged at 4500 rpm for 10 min; supernatants were transferred to covered tubes and fatty acids of the lipid extract were converted to methyl esters by using 2% sulfuric acid (v/v) in methanol (Christie, 1992). Fatty acid methyl ester forms were extracted with n-hexane. Analysis was performed in a Shimadzu GC-17A V3 (Kyoto, Japan) instrument gas chromatograph equipped with a Flame Ionization Detector (FID) and a 25 m, 0.25 mm i.d. Permabond fused-silica capillary column (Macherey-Nagel, Germany). The oven temperature was programmed between 120-220°C, 5°C min⁻¹. Injector and FID temperatures were 240 and 280°C, respectively. The nitrogen carrier gas flow was 1 mL min⁻¹. The methyl esters of fatty acids were identified by comparison with authentic external standard mixtures analyzed under the same conditions. Class GC 10 software version 2.01 (Shimadzu, Kyoto, Japan) was used to process the data. The results were expressed as percent.

Determination of superoxide dismutase activity: Total SOD activity was determined by the method of Sun *et al.* (1988). The technique is based on inhibition of Nitro Blue Tetrazolium (NBT) reduction by the xanthine-xanthine oxidase system as the superoxide generator. The SOD activity was measured in the ethanol phase of the supernatant after 1 mL of ethanol-chloroform mixture (5:3, v/v) was added to the same volume of sample and centrifuged. One unit of SOD was defined as the amount of enzyme causing 50% inhibition in the NBT reduction rate. The SOD activity was expressed as U g⁻¹ protein.

Glutathione peroxidase (GSH-Px) activity: GSH-Px activity was evaluated by the Flohe and Gunzler method (Flohe and Gunzler, 1984; Tasset *et al.*, 2008). The testes tissues were homogenized in ice-cold buffer (0.1 M KH₂PO₄/K₂HPO₄, pH 7.0 plus 29.2 mg EDTA in 100 mL of distilled water and 10.0 mg digitonin in 100 mL of distilled water, final volume, 2000 mL) to produce a homogenate. The homogenates were then centrifuged at 10,000×g for 10 min at 4°C. The GSH-Px assay is based on the oxidation of NADPH to NAD⁺, catalyzed by a limiting concentration of glutathione reductase with maximum absorbance at

340 nm. The activity of GSH-Px is expressed as units per milligram of protein ($U\ mg^{-1}\ protein$) and absorbance was evaluated in a spectrophotometer (UV-1603 Shimadzu, Kyoto, Japan).

Determination of tissue Malondialdehyde (MDA) levels: Esterbauer method that is lipid peroxidation measurement method was used to determination of tissue malondialdehyde levels (Esterbauer and Cheeseman, 1990). Tiobarbituric acid which is reacting with MDA at 90-95°C was used in order to occur pink kromogen. Then tissue samles was cooled for 30 min and absorbances was read as spectrophotometrical at 532 nm. All results was calculated as $nmol\ g^{-1}\ tissue\ protein$.

Statistical analysis: The SPSS software for Windows, v.10 (SPSS, Chicago, IL) was used for the statistical treatment of the data. The results are expressed as mean values±standard deviation. Differences between means were established by analysis of variance with $p<0.05$ considered significant. Significant differences among treatment and control groups were interpreted using the Tukey’s honest significant difference post-hoc test.

RESULTS

The present study examined the MDA, SOD, GSH-Px and fatty acid composition on testes of Wistar rats which were administered with normal fed, fasted state and linalool applications. The results were shown in Table 1 and Fig. 1-3.

Table 1: Fatty acid composition of rat testes (%g); a) $p<0.05$ significant, compared to control; b) $p<0.05$ significant, compared to 24 h F

Fatty acids	Control	24 h F	24 h F+L
C16:0	23.82±1.52	24.45±1.37	22.41±1.74
C16:1 n7	4.98±1.16	2.46±0.54 ^a	3.68±0.42 ^{ab}
C17:0	0.41±1.02	0.37±0.12	0.36±0.05
C18:0	5.40±0.48	5.98±0.52	6.02±1.23
C18:1 n9	16.78±1.56	15.43±1.43	15.23±1.42
C18:1 n11	4.68±1.23	4.21±0.12	4.34±0.19
C18:2 n6	12.18±1.24	11.98±1.15	12.14±1.32
C18:3 n6	1.35±0.22	1.45±0.17	1.22±0.21
C20:1 n9	0.59±0.21	0.64±0.14	0.49±0.02
C20:2 n6	0.42±0.84	0.47±0.04	0.43±0.16
C20:3 n6	0.84±0.12	0.90±0.06	0.89±0.18
C20:4 n6	12.10±1.54	13.02±1.23	13.42±0.34
C22:2 n6	0.43±0.17	0.50±0.08	0.46±0.46
C22:4 n6	1.18±0.14	0.65±0.18 ^a	1.05±0.52 ^b
C22:5 n6	12.65±1.23	8.47±1.58 ^a	10.94±1.22 ^{ab}
C22:5 n3	0.47±0.04	0.51±0.05	0.49±0.07
C22:6 n3	1.08±0.15	0.62±0.06 ^a	1.24±0.16 ^b
∑Saturated	29.63±0.20	30.80±0.54	28.79±1.08
∑Unsaturated	72.73±1.23	61.31±1.06 ^a	66.02±1.17 ^{ab}
∑MUFA	26.44±0.12	22.74±1.10	23.51±1.22
∑PUFA	46.29±1.02	38.57±1.23 ^a	42.51±1.09 ^b
Satur./unsatur.	0.40±0.03	0.50±0.02	0.43±0.07

Fatty acid composition of rat testes on experimental groups: The results of the different fatty acids values are shown in Table 1. The saturated fatty acids which are C16:0, C17:0 and C18:0 fatty acids composition were not statistical significant among all experimental groups ($p>0.05$). Adversely, the monounsaturated fatty acids like C16:1 were lower compared to the control group. Likewise, polyunsaturated fatty acids like C22:4, C22:5 C22:6 in the fasted groups were lower compared to control group. Both monounsaturated fatty acids and polyunsaturated fatty acids compositions in the linalool application groups were higher than fasted groups ($p<0.05$). C22:6 n3 and C22:4 n6 fatty acids levels in the linalool application groups were not statistically significant compared to control ($p>0.05$).

The activites of Superoxide Dismutase (SOD), Glutathione Peroxidase (GSH-Px) and the level of Malondialdehyde (MDA) in testicular tissues: The results indicate that the GSH-Px level decreased in the both fasted group and linalool application group compared to control. Similarly, SOD levels were lower in the both fasted and linalool application group compared to control ($p<0.05$). Both SOD and GSH-Px levels in linalool




Fig. 1: The levels of GSH-Px in the testes; a) $p<0.05$ significant compared to control




Fig. 2: The levels of SOD in the testes; a) $p<0.05$ significant compared to control




Fig. 3: The levels of MDA in the testes; a) $p < 0.05$ significant compared to control; b) $p < 0.05$ significant compared to 24 h F

application were not statistical significant compared to fasted group ($p > 0.05$, Fig. 1 and 2). Besides, the finding indicate that MDA level increased in the fasted group compared to control group. Diametrically, the MDA level in the linalool group was lower as statistically than the fasted group ($p < 0.05$, Fig. 3).

DISCUSSION

Fasting state or caloric restriction effect testicular cell metabolism, sex behavior and rogen receptors and testosterone concentration (Bandyopadhyay *et al.*, 1999). Liver glycogen stores are virtually depleted within 24 h (Rothman *et al.*, 1991). As liver glycogen stores are virtually depleted within 24 h, gluconeogenesis supplies the requirements of the brain and other glucose-requiring tissues. Falling insulin concentrations lead to both net proteolysis in muscle with release of alanine and glutamine and lipolysis in adipose tissue with release of glycerol and Non-Esterified Fatty Acids (NEFAs) (Nygren, 2006; Awad *et al.*, 2009). The oxidative stres in testicular milieu is associated with DNA damage, lipid peroxidation and produces higher frequency of abnormal sperms with significant effect on male fertility (Kumar *et al.*, 2002). Under normal physiological conditions, free radicals are generated in testis subcellular compartments, particularly mitochondria which are subsequently scavenged by antioxidant defense systems of the corresponding cellular compartments (Bergendi *et al.*, 1999). Further, the mitochondrial membrane is more susceptible to Lipid Peroxidation (LPO) as this compartment is rich in polyunsaturated fatty acids and has been shown to contain low amounts of antioxidants (Selvakumar *et al.*, 2005). Adversely, polyunsaturated fatty acids levels were calculated as low level by Gavazza and Catala (2003) in rat testes during

Fe²⁺-iduced oxidative stres (Gavazza and Catala, 2003). In paralel in the study, total Polyunsaturated Fatty Acid (PUFA) levels were low rate on fasting groups compared to controls. Particularly, C22:4 n6 (Docosatetraenoic Acid), C22:5 n6 (Eicosapentaenoic Acid) and C22:6 n3 (Decosahexaenoic Acid) fatty acids were low in fasting groups compared to control. We also calculated that total PUFA levels were high in linalool groups compared to fasting group. PUFAs of linalool application were not statistically significant difference compared to control ($p > 0.05$, Table 1). Abul *et al.* (2002) expressed that membranes lipids destruction made with aldehydes occurs endoperoksit (Abul *et al.*, 2002; Rosenblum *et al.* (1989) also indacted that lipids peroxidations were occured by reactive oxygen species (Rosenblum *et al.*, 1989). In the study, malondialdehyde that is marker of oxidative stress were high in fasting groups and GSH-Px, SOD that are antioxidant enzyme in fasting groups were low compared to control (Fig. 2 and 3). Linalool applications improved MDA levels. That is MDA level in the linalool groups was low level compared to fasting groups ($p < 0.05$, Fig. 3). Fasting state or starvation effect some hormone levels in the rat tissues. Hormones such as insulin, glucagon, leptin, ghrelin and the gonadal hormones determine energy partitioning that is they direct metabolic fuels into or out of storage (Schneider and Watts, 2009; Wells, 2009). Especially, ghrelin peptide that is the endogenous ligand for the Growth Hormone (GH) Secretagogue Receptor (GHS-R) increase during fasting state (Kojima and Kangawa, 2005; Kojima *et al.*, 1999). The ghrelin gene is expressed in stomach, small intestine, brain, pituitary, salivary gland, adrenal, ovary and testis with maximum expression occurring in the stomach (Ghelardoni *et al.*, 2006; Aydin *et al.*, 2005). It has recently demonstrated the ability of ghrelin has anti-proliferative effects on different testicular cell types and is a negative modulator of male reproductive system (Kheradmand *et al.*, 2009). Ghrelin effects to fatty acid synthesis and related transcription factor mRNA levels (Buyse *et al.*, 2009; Muccioli *et al.*, 2004). Lopez *et al.* (2008) expressed that fasting state reduced malonyl-CoA and fatty acid synthase levels (Lopez *et al.*, 2008). Theander-Carrillo *et al.* (2006) reported that ghrelin action in the brain controls adipocyte metabolism (Theander-Carrillo *et al.*, 2006). The Stearoyl-CoA Desaturases (SCD) are important on unsaturated fatty acids metabolism in animal tissues. SCD can act on chain lengths from C12:0-C19:0 (Mauvoisin and Mounier, 2011; Zolfaghari and Ross, 2003). However, SCD1 preferred desaturation substrates are palmitoyl-CoA (C16:0) and stearoyl-CoA (C:18:0) which are converted to palmitoleoyl-CoA (C16:1) and oleoyl-CoA (C18:1)

respectively (Flowers and Ntambi, 2008). Ambati *et al.* (2010) reported that in rat, injection of ghrelin decreases the level of Stearoyl-CoA Desaturase 1 (SCD1) mRNA in adipose tissue (Ambati *et al.*, 2010). In parallel, we determined that C16:1 was lower in the 24 h F groups than control ($p < 0.05$). Many researcher stated that no histopathologic abnormalities were observed in animals in the high-dose linalool in the liver, adrenals, brain, heart, kidneys, thyroids, mesenteric lymph node, spinal cord, testes, ovaries, spleen, urinary bladder, sternal bone marrow, pituitaries or sciatic nevre (Letizia *et al.*, 2003; Tepe *et al.*, 2004; Pattnaik *et al.*, 1997; Mitic-Culafic *et al.*, 2009). The results of the study are consistent with these reports, demonstrating that linalool effects MDA levels that is indicator for lipid peroksidation. At the same time we determined that linalool protected both monounsaturated and polyunsaturated fatty acids levels (Table 1). Researchers also determined that both C16:1 C22:6 fatty acids were higher as statistically in 24 h F+Linalool groups than 24 h F groups ($p < 0.05$).

CONCLUSION

In the study, we suggest that linalool was a potential candidate in combating the testicular abnormalities induced by fasted. According to results of the study, the linalool can protect polyunsaturated fatty acids composition in the rat testes against to fasted state. Although the exact mechanisms remain to be clarified, linalool could be an protective effects to testicular cell metabolism.

ACKNOWLEDGEMENTS

Thanks to Firat University Animal Research Center where provide the experimental animals and thanks to Dr. Mahmut Yilmaz who accomplished the sataistical analysis and graphical arrangement.

REFERENCES

Abdelmegeed, M.A., K.H. Moon, J.P. Hardwick, F.J. Gonzalez and B.J. Song, 2009. Role of peroxisome proliferator-activated receptor- α in fasting-mediated oxidative stress. *Free Radic. Biol. Med.*, 47: 767-778.

Abul, H.T., T.C. Mathew, F. Abul, H. Al-Sayer and H.M. Dashti, 2002. Antioxidant enzyme level in the testes of cirrhotic rats. *Nutrition*, 18: 56-59.

Ambati, S., Q. Li, S. Rayalam, D.L. Hartzell, M.A. Della-Fera, M.W. Hamrick and C.A. Baile, 2010. Central leptin versus ghrelin: Effects on bone marrow adiposity and gene expression. *Endocrine*, 37: 115-123.

Awad, S., D. Constantin-Teodosiu, I.A. Macdonald and D.N. Lobo, 2009. Short-term starvation and mitochondrial dysfunction- a possible mechanism leading to postoperative insulin resistance. *Clin. Nutr.*, 28: 497-509.

Aydin, S., I. Halifeoglu, I.H. Ozercan, F. Erman and N. Kilic *et al.*, 2005. A comparison of leptin and ghrelin levels in plasma and saliva of young healthy subjects. *Peptides*, 26: 647-652.

Bandyopadhyay, U., D. Das and R.K. Banerjee, 1999. Reactive oxygen species: Oxidative damage and pathogenesis. *Curr. Sci.*, 77: 658-666.

Bergendi, L., L. Benes, Z. Durackova and M. Ferencik, 1999. Chemistry, physiology and pathology of free radicals. *Life Sci.*, 65: 1865-1874.

Bickers, D., P. Calow, H. Greim, J.M. Hanifin and A.E. Rogers *et al.*, 2003. A toxicologic and dermatologic assessment of linalool and related esters when used as fragrance ingredients. *Food Chem. Toxicol.*, 41: 919-942.

Bourre, J.M., M. Bonneil, J. Chaudiere, M. Clement and O. Dumont *et al.*, 1992. Structural and functional importance of dietary polyunsaturated fatty acids in the nervous system. *Adv. Exp. Med. Biol.*, 318: 211-229.

Brown, B.L., J.W. Allis, J.E. Simmons and D.E. House, 1995. Fasting for less than 24 h induces cytochrome P450 2E1 and 2B1/2 activities in rats. *Toxicol. Lett.*, 81: 39-44.

Buyse, J., S. Janssen, S. Geelissen, Q. Swennen, H. Kaiya, V.M. Darras and S. Dridi, 2009. Ghrelin modulates fatty acid synthase and related transcription factor mRNA levels in a tissue-specific manner in neonatal broiler chicks. *Peptides*, 30: 1342-1347.

Celik, S. and A. Ozkaya, 2002. Effects of intraperitoneally administered lipoic acid, vitamin E and linalool on the level of total lipid and fatty acids in guinea pig brain with oxidative stress induced by H₂O₂. *J. Biochem. Mol. Biol.*, 35: 547-552.

Christie, W.W., 1992. *Gas Chromatography and Lipids: A Practical Guide*. The Oil Press, Glaskow, pp: 307.

Conn, P.M., 1986. The molecular basis of gonadotropin-releasing hormone action. *Endocrinol. Rev.*, 7: 3-10.

Droge, W., 2002. Free radicals in the physiological control of cell function. *Physiol. Rev.*, 82: 47-95.

Esterbauer, H. and K.H. Cheeseman, 1990. Determination of aldehydic lipid peroxidation products: Malonaldehyde and 4-hydroxynonenal. *Methods Enzymol.*, 186: 407-421.

Flohe, L. and W.A. Gunzler, 1984. Assays of glutathione peroxidase. *Methods Enzymol.*, 105: 114-121.

Flowers, M.T. and J.M. Ntambi, 2008. Role of stearoyl-coenzyme A desaturase in regulating lipid metabolism. *Curr. Opin. Lipidol.*, 19: 248-256.

- Gavazza, M. and A. Catala, 2003. Melatonin preserves arachidonic and docosapentaenoic acids during ascorbate-Fe²⁺ peroxidation of rat testis microsomes and mitochondria. *Int. J. Biochem. Cell Biol.*, 35: 359-366.
- Ghelardoni, S., V. Carnicelli, S. Frascarelli, S. Ronca-Testoni and R. Zucchi, 2006. Ghrelin tissue distribution: Comparison between gene and protein expression. *J. Endocrinol. Invest.*, 29: 115-121.
- Halliwell, B., 1994. Free radicals, antioxidants and human disease: Curiosity, cause, or consequence. *Lancet*, 344: 721-724.
- Hara, A. and N.S. Radin, 1978. Lipid extraction of tissues with a low-toxicity solvent. *Anal. Biochem.*, 90: 420-426.
- Kara, H., A. Cevik, V. Konar, A. Dayangac and M. Yilmaz, 2007. Protective effects of antioxidants against cadmium-induced oxidative damage in rat testes. *Biol. Trace Element Res.*, 120: 205-211.
- Kheradmand, A., L. Roshangar and M. Taati, 2009. The role of ghrelin on the morphometry and intracellular changes in the rat testis. *Tissue Cell*, 41: 105-111.
- Kojima, M. and K. Kangawa, 2005. Ghrelin: Structure and function. *Physiol. Rev.*, 85: 495-522.
- Kojima, M., H. Hosoda, Y. Date, M. Nakazato, H. Matsuo and K. Kangawa, 1999. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature*, 402: 656-660.
- Koroch, A.R., H.R. Juliani and J.A. Zygadlo, 2007. Bioactivity of Essential Oils and their Components. In: *Flavours and Fragrances*, Berger, R.G. (Ed.). Springer-Verlag, Berlin, Heidelberg, pp: 87-115.
- Kumar, T.R., K. Doreswamy, B. Shrilatha and Muralidhara, 2002. Oxidative stress associated DNA damage in testis of mice: Induction of abnormal sperms and effects on fertility. *Mutation Research Genet. Toxicol. Environ. Mutagen.*, 513: 103-111.
- Lapczynski, A., C.S. Letizia and A.M. Api, 2008. Addendum to fragrance material review on linalool. *Food Chem. Toxicol.*, 46: S190-S192.
- Leal-Cardoso, J.H., K.S. da Silva-Alves, F.W. Ferreira-da-Silva, T. dos Santos-Nascimento and H.C. Joca *et al.*, 2010. Linalool blocks excitability in peripheral nerves and voltage-dependent Na⁺ current in dissociated dorsal root ganglia neurons. *Eur. J. Pharmacol.*, 645: 86-93.
- Letizia, C.S., J. Cocchiara, J. Lalko and A.M. Api, 2003. Fragrance material review on linalool. *Food Chem. Toxicol.*, 41: 943-964.
- Lonergan, P.E., D.S. Martin, D.F. Horrobin and M.A. Lynch, 2002. Neuroprotective effect of eicosapentaenoic acid in hippocampus of rats exposed to gamma-irradiation. *J. Biol. Chem.*, 277: 20804-20811.
- Lopez, M., R. Lage, A.K. Saha, D. Perez-Tilve and M.J. Vazquez *et al.*, 2008. Hypothalamic fatty acid metabolism mediates the orexigenic action of ghrelin. *Cell Metab.*, 7: 389-399.
- Mauvoisin, D. and C. Mounier, 2011. Hormonal and nutritional regulation of SCD1 gene expression. *Biochimie*, 93: 78-86.
- Mitic-Culafic, D., B. Zegura, B. Nikolic, B. Vukovic-Gacic, J. Knezevic-Vukcevic and M. Filipic, 2009. Protective effect of linalool, myrcene and eucalyptol against t-butyl hydroperoxide induced genotoxicity in bacteria and cultured human cells. *Food Chem. Toxicol.*, 47: 260-266.
- Muccioli, G., N. Pons, C. Ghe, F. Catapano, R. Granata and E. Ghigo, 2004. Ghrelin and des-acyl ghrelin both inhibit isoproterenol-induced lipolysis in rat adipocytes via a non-type 1a growth hormone secretagogue receptor. *Eur. J. Pharmacol.*, 498: 27-35.
- Nygren, J., 2006. The metabolic effects of fasting and surgery. *Best Practice Res. Clin. Anaesthesiol.*, 20: 429-438.
- Pattnaik, S., V.R. Subramanyam, M. Bapaji and C.R. Kole, 1997. Antibacterial and antifungal activity of aromatic constituents of essential oils. *Microbios*, 89: 39-46.
- Rosenblum, E.R., J.S. Gavalier and D.H. Van Thiel, 1989. Lipid peroxidation: A mechanism for alcohol-induced testicular injury. *Free Radic. Biol. Med.*, 7: 569-577.
- Rothman, D.L., I. Magnusson, L.D. Katz, R.G. Shulman and G.I. Shulman, 1991. Quantitation of hepatic glycogenolysis and gluconeogenesis in fasting humans with ¹³C NMR. *Science*, 254: 573-576.
- Santos, A.M., M.R. Ferraz, C.V. Teixeira, F.J. Sampaio and C.D.F. Ramos, 2004. Effects of undernutrition on serum and testicular testosterone levels and sexual function in adult rats. *Horm. Metab. Res.*, 36: 27-33.
- Schneider, J.E. and A.G. Watts, 2009. Energy Partitioning, Ingestive Behavior and Reproductive Success. In: *Hormones, Brain and Behavior*, Pfaff, D.W. (Ed.). 2nd Edn., Vol. 1, Academic Press, USA., pp: 205-258.
- Selvakumar, E., C. Prahallathan, Y. Mythili and P. Varalakshmi, 2005. Beneficial effects of D1- α -lipoic acid on cyclophosphamide-induced oxidative stress in mitochondrial fractions of rat testis. *Chemico Biol. Interactions*, 152: 59-66.
- Sikka, S.C., 1996. Oxidative stress and role of antioxidants in normal and abnormal sperm function. *Frontiers Biosci.*, 1: 78-86.
- Sun, Y., L.W. Oberley and Y. Li, 1988. A simple method for clinical assay of superoxide dismutase. *Clin. Chem.*, 34: 497-500.

- Tasset, I., J. Pena, I. Jimena, M. Feijoo, M.D.C. Munoz, P. Montilla and I. Tunez, 2008. Effect of 17β -estradiol on olfactory bulbectomy-induced oxidative stress and behavioral changes in rats. *Neuropsychiatric Dis. Treatment*, 4: 441-449.
- Tepe, B., E. Donmez, M. Unlu, F. Candan and A. Sokmen *et al.*, 2004. Antimicrobial and antioxidative activities of the essential oils and methanol extracts of *Salvia cryptantha* (Montbret et Aucher ex Benth.) and *Salvia multicaulis* (Vahl). *Food Chem.*, 84: 519-525.
- Theander-Carrillo, C., P. Wiedmer, P. Cettour-Rose, R. Nogueiras and D. Perez-Tilve *et al.*, 2006. Ghrelin action in the brain controls adipocyte metabolism. *J. Clin. Invest.*, 116: 1983-1993.
- Wells, T., 2009. Ghrelin-defender of fat. *Progress Lipid Res.*, 48: 257-274.
- Zolfaghari, R. and A.C. Ross, 2003. Recent advances in molecular cloning of fatty acid desaturase genes and the regulation of their expression by dietary vitamin A and retinoic acid. *Prostaglandins, Leukotrienes Essential Fatty Acids*, 68: 171-179.