

Hypolipidemic and Hypocholesterolemic Effect of Medicinal Plant Combination in the Diet of Rats: Black Cumin Seed (*Nigella sativa*) and Turmeric (Curcumin)

M.H. Al-Nazawi and S.M. El-Bahr

Department of Physiology, Biochemistry and Pharmacology,
College of Veterinary Medicine and Animal Resources, King Faisal University, Al-Ahsa, Saudi Arabia

Abstract: A total of 24 rats were fed a diet either without (Group1) or with black cumin seed 2% (Group 2) or turmeric 2% (Group 3) or plant combination (1% each; Group 4) for 6 weeks. Body weight, feed intake and feed conversion ratio were recorded. At the end of the experiment, blood samples were collected from all groups. The whole blood and harvested sera were used for determination of hematological and biochemical parameters, respectively. The body weight gain and feed conversion ratio were comparable in all treated group when compared with the control. The Packed Cell Volume (PCV), Hemoglobin percentage (Hb%) and Mean Corpuscular Hemoglobin Concentration (MCHC) remained unchanged in all treated group when compared with the control. Spectrophotometric analysis revealed that serum total protein was significantly ($p < 0.05$) higher in black seed (Group 2) and turmeric (Group 3) treated groups when compared with the control. Globulin value was significantly ($p < 0.05$) higher in black cumin seed treated group than those of control and other groups. The hypolipidemic effect was recorded only in rats fed combination of medicinal plants. The hypocholesterolemia accompanied by lower level of Low Density Lipoprotein cholesterol (LDL-c) was recorded in rats fed both medicinal plants either alone or in combination. However, this effect was more pronounced in rats fed plants combination. The values of albumin, Alanine Transaminase (ALT), Aspartate Transaminase (AST), High Density Lipoprotein cholesterol (HDL-c), Very Low Density Lipoprotein cholesterol (VLDL-c), uric acid and creatinine in all treated groups remained comparable to those of the control group. The present study concluded that plant combination was safe to the animals as reflected on unchanged liver and kidney function biomarkers. Interestingly, black cumin seed and turmeric has hypocholesterolemic effect whenever administered alone or in combination (1% each) whereas the hypolipidemic effect was related to the combined administration of both plants only in the diet of rats.

Key words: Medicinal plants, blood, serum, lipid profile, biochemical parameters, rats

INTRODUCTION

Hyperlipidemia being an important risk factor for cardiovascular disease is a serious public health problem in the world. Its major role in the pathogenesis of atherosclerosis has been implicated by several clinical and epidemiological studies (Jaffar *et al.*, 2004). Regarding its treatment, now a days there is an increasing interest toward the potential health benefits of medicinal plants.

Black cumin seed (*Nigella sativa*) is a herbaceous plant which is a member of the Ranunculacea family. The seed contains >30% fixed oil and about 0.45% (w/w) of volatile oils (Abd El-Aal and Attia 1993; El-Dakhkhny *et al.*, 2000). The black cumin seed has antibacterial and antifungal properties (Rathee *et al.*, 1982; Gilani *et al.*, 2004; Hasan *et al.*, 2005; Abu-Al-Basal, 2011),

anthelmintic activity against tape worms (Agarwal *et al.*, 1979) and choleric action (Mahfouz *et al.*, 1962) and anticarcinogenic (Aggarwal *et al.*, 2005). Injection of black cumin seed extract to male rats (0.4 mL kg⁻¹ of body weight) decreased cholesterol levels but it increased the total serum protein and globulins levels meanwhile the values of serum albumin and urea were not significantly altered (Hedaya, 1995). Black cumin seeds have been reported to possess a favorable effect on serum lipid profile by decreasing the levels of total cholesterol, LDL-c, TAG and by elevating the HDL-c levels in rats (Badari *et al.*, 2000; Bashandy, 1996; Chaudary, 1996; El-Dakhkhny *et al.*, 2000; Hassanin and Hassan, 1996; Tayyab, 1995; Alsaif, 2008).

Turmeric (*Curcuma longa*) is a perennial herb that grows to a height of three to five feet and is cultivated

extensively in Asia and other countries with a tropical climate. Turmeric is a member of the *Curcuma* botanical group which is part of the ginger family of herbs, Zingiberaceae. Curcumin, the active ingredient from the spice turmeric is a potent antioxidant (El-Bahr *et al.*, 2007; Salama and El-Bahr, 2007; Sivabalan and Anuradha, 2010; Mehta *et al.*, 2012) and hepatoprotective properties (Pal *et al.*, 2001). The root and rhizome (underground stem) of the *Curcuma longa* L. plant is crushed and powdered into ground turmeric. Turmeric has hypocholesterolemic effect on hypercholesterolemic rats and decreased the raised level of liver cholesterol of cholesterol-fed rats (Reddy and Lokesh, 1994). The combined administration of Turmeric and black cumin seed was examined only in mugil cephalus fish (El-Bahr and Saad, 2008). Therefore, the present study aimed to investigate the effect of combined administration of black cumin seed and turmeric on some biochemical parameters with special references to lipid profiles in rats.

MATERIALS AND METHODS

Experimental animals: A total of twenty four adult albino rats weighing between 193±7 g (8 weeks age at the start) were obtained from the laboratory animal house of the College of Veterinary Medicine, King Faisal University, Saudi Arabia. They were maintained as performed by national guidelines and protocols, approved by the University Animal Ethics Committee. They were housed in clean and disinfected plastic cages. Commercial basal diet and water were provided *ad libitum*. Rats were subjected to natural photoperiod of 12 h light: dark cycle throughout the experimental period (6 weeks). All rats received basal diet for 2 weeks before the start of the experiment for adaptation and to ensure normal growth and behavior. They were maintained in their respective groups for 42 days, monitored closely every day and weighed every week. The rat food was weighed every day before and after feeding the animals to determine the daily food intake.

Plant materials: Turmeric (*curcuma longa*) and black cumin seed (*Nigella sativa*) were purchased from a local market in Al-Ahsa, Saudi Arabia and identified by botanists in the Faculty of Agriculture, King Faisal University, Saudi Arabia. Both plants were analyzed and their ingredients are shown in Table 1 (AOAC, 1995). The whole black cumin seeds and Turmeric were crushed in a blender and mixed with the diet (20 g kg⁻¹ diet; 2%) while plant combination was used to obtain (10 g kg⁻¹ diet; 1% for each plant).

Table 1: Proximate analysis of ingredients used in the diets of the experiment, dry matter basis

Ingredients	Percentage					
	Moisture	CP	EE	CF	NFE*	Ash
Black cumin seed	5.8	19.54	34.4	6.1	36.16	3.80
Turmeric	13.5	11.74	6.4	7.2	70.07	4.59

CP: Crude Protein; EE: Ether Extract; CF: Crude Fiber and *NFE: Nitrogen Free Extract is calculated by differences

Experimental design and sampling analysis: The rats were divided into five groups (6 rats each).

- Group 1: Rats fed basal diet (control rats)
- Group 2: Black cumin seed treated rats
- Group 3: Turmeric treated rats
- Group 4: Black cumin seed and turmeric plant combination treated rats

Daily measurements of body weight and food intake were recorded. Body weight, feed intake and feed conversion were determined at the start and at the end of the experiment (after 6 weeks). At the end of the experiment, blood samples were collected with and without anticoagulant for whole blood and serum analysis, respectively. Packed Cell Volume (PCV) was measured by a standard manual technique using microhematocrit capillary tubes and centrifuged at 2500 rpm for 5 min measurement of hemoglobin was performed using hemometer. Serum was separated by centrifugation for 10 min at 1200 g and was immediately frozen at -20°C until the time of analysis. The sera were used for spectrophotometric determination of the activities of Aspartate Transaminase (AST) and Alanine Transaminase (ALT) as directed by Reitman and Frankel (1957). In addition, total protein, albumin and globulin values were determined spectrophotometrically as implied by the methods of Doumas *et al.* (1981) and Reinhold (1953), respectively. Serum uric acid and creatinine were determined according to the method described by Tabacco *et al.* (1979). Furthermore, the obtained sera were used for spectrophotometric analysis of serum Triacylglycerol (TAG), total cholesterol by using of enzymatic method of commercial kits according to the methods of Gottfried and Rosenberg (1973) and Zak *et al.* (1954), respectively. Very Low Density Lipoprotein cholesterol (VLDL-c) was calculated by division of TAG by 5 while the LDL-c was calculated as total cholesterol (HDL-c + VLDL-c) = mg dL⁻¹ (Bauer, 1982).

Statistical analysis: All the grouped data were statistically evaluated and the significance of changes caused by various treatments was determined using Students t-test (Chou, 1975). The results have been

expressed as means±SD from seven rats in each group. A two ways ANOVA was also done wherever appropriate. The level of statistical significance was set at $p < 0.05$.

RESULTS AND DISCUSSION

Performance data: The present study showed that there was no significant changes ($p < 0.05$) in weight gain in all treated group when compared with the control (Table 2). The data shown in Table 3 demonstrated that feed intake was significantly higher ($p < 0.05$) in all treated group when compared with the control without any significant difference ($p < 0.05$) in feed conversion ratio. The increase in feed intake was more pronounced in rats fed plant combination (Group 4) than other treated groups (Group 2 and 3).

Hematological and biochemical parameters: Hematological findings (Table 4) indicated that PCV, Hb and MCHC were remained unchanged significantly ($p < 0.05$) in all treated group when compared with the control.

Table 2: Effect of oral administration of turmeric and/or black cumin seed for 6 weeks on body weight gain (g)

Groups	Body weight (g)		Gain in body weight (g)
	Initial	6th week	
I	228±5.1	262±6.2	34
II	228±7.2	258±7.1	30
III	230±4.2	260±3.1	30
IV	236±5.2	266±8.1	30

I (control), II (black cumin seed treated group), III (turmeric treated group), IV (black cumin seed and turmeric treated group) and values are mean±SD of 6 rats

Table 3: Effect of oral administration of turmeric and/or black cumin seed for 6 weeks on food intake (g day⁻¹) and feed conversion ratio (g diet/g weight) in rats

Groups	Food intake (g day ⁻¹)		Food conversion ratio	
	1st week	6th week	1st week	6th week
I	25.5±0.1	23.0±0.1	0.11±0.1	0.09±0.01
II	27.0±0.2*	26.2±0.5*	0.12±0.1	0.10±0.01
III	30.1±0.1*	25.0±0.6*	0.13±0.1	0.10±0.02
IV	31.2±0.2**	30.0±0.1**	0.13±0.1	0.11±0.01

I (control), II (black cumin seed treated group), III (turmeric treated group), IV (black cumin seed and turmeric treated group), values are mean±SD of 6 rats; *Means within the same column are statistically significant when compared to control (Group I) at $p < 0.05$; ** Means within the same column are statistically significant when compared to other group at $p < 0.05$

Table 4: Effect of oral administration of turmeric and/or black cumin seed for 6 weeks on hematological parameters in rats

Parameters (%)	I	II	III	IV
PCV	41.1±0.11	41.2±0.12	42.3±0.60	42.3±0.60
Hb	13.2±0.30	12.9±0.70	13.6±0.60	12.6±0.80
MCHC	32.1±1.22	31.3±0.91	32.1±0.10	29.81±0.1

I (control), II (black cumin seed treated group), III (turmeric treated group), IV (black cumin seed and turmeric treated group) and values are mean±SD of 6 rats

Serum total protein level was significantly ($p < 0.05$) higher in black seed (Group 2) and turmeric (Group 3) treated groups when compared with the control (Group 4). In addition, globulin value was significantly ($p < 0.05$) higher in black cumin seed treated group than those of the control and other groups (Group 4). Albumin, ALT, AST, HDL-c, VLDL-c, uric acid and creatinine levels were unchanged significantly ($p < 0.05$) in all treated group when compared with the control (Group 4). Spectrophotometric analysis revealed a significant ($p < 0.05$) lower levels of TAG in rats received plant combination (Group 4) when compared with the control whereas the level of this parameter was comparable to the control in other treated groups ($p < 0.05$). The present findings showed a significant ($p < 0.05$) lower levels of total cholesterol and LDL-c in rats received both medicinal plants either alone or in combination when compared with the control. However, the decrease in total cholesterol and LDL-c was more pronounced ($p < 0.05$) in rats received plant combination (Table 4).

Although, some publications have described different beneficial properties of black cumin seed as its extract, oil or active compound (s) such as thymoquinone, to date few have examined the effects of whole or crushed seed (Ibraheim, 2002; El-Bahr, 2007; Buriro and Tayyab, 2007; Pourghassem-Gargari *et al.*, 2009; Nasir and Grashorn, 2010; Tasawar *et al.*, 2011). In the contrary, many research articles demonstrated the beneficial effect of turmeric (Soliman, 2005; El-Bahr *et al.*, 2007; Salama and El-Bahr, 2007; Basavaraj *et al.*, 2011; Nouzarian *et al.*, 2011). To the researcher knowledge, the present study is the first study demonstrated the effect of combined administration of these medicinal plants on lipid profiles or rats serum. However, El-Bahr and Saad (2008) demonstrated the positive effect of combined administration of black cumin seed and turmeric in *Mugil cephalus* fish.

The data concerning the effect of black cumin seed or turmeric in different animal species was contradicted. In the present study, blood hematology picture, PCV, Hb and MCHC did not show significance difference between all treated groups when compared with the control. Similar results demonstrated that neither turmeric (Emadi *et al.*, 2007; Basavaraj *et al.*, 2011) nor black cumin seed (El-Bagir *et al.*, 2010) changed the hematological picture in animal and birds. In the other hand, some studies demonstrated that turmeric increased PCV and hemoglobin percentage (Sugiharto *et al.*, 2011) while other studies (Ekanem and Yusuf, 2008) reported a significant elevation of hematological parameters in rat fed lack cumin seed.

The present findings indicated that albumin, ALT, AST, HDL-c, VLDL-c, uric acid and creatinine were not affected by dietary inclusion of black cumin seed and/or turmeric powder in all treated rats when compared with the control. This indicated that both medicinal plants at the examined dose were safe to be included in rat diet as reflected on the above unchanged liver and kidney function biomarkers. In consistence of the results, El-Bagir *et al.* (2010) reported that inclusion of 15% black cumin seed in the diet of rats did not affect the concentration of albumin and globulins. In addition, Ekanem and Yusuf (2008) showed significant increase in serum liver enzyme activities in rats fed black seed oil. Moreover, Shewita and Taha (2011) demonstrated higher ALT and lower AST level in broiler chick fed black cumin seed. Also, Basavaraj *et al.* (2011) supplementation of Turmeric to broiler rabbits did not show significant changes in all above mentioned biochemical parameters (Table 5).

The previously reported hypolipidemic effect of Black cumin seed (Pourghassem-Gargari *et al.*, 2009; El-Bagir *et al.*, 2010) and turmeric (Hussain, 2002; Srinivasan, 2005; Soliman, 2005; Sugiharto *et al.*, 2011) was not similar to the findings. Similar hypolipidemic effect was observed in the present study only when both plants administered together as a combination. This might be indicated that half dose of each plant is enough to induce the hypolipidemic effect.

Cholesterol is transported via blood by lipoproteins. HDL-c (good cholesterol) transports it from tissues to liver and LDL-c (bad cholesterol) does it in the opposite direction. Therefore, decrease in serum LDL-c cholesterol is an indication of low rate of transportation of cholesterol from liver to tissues and subsequent transformation of cholesterol into bile acid by liver enzymes. HDL-c/LDL-c ratio consider as one of the most important parameters in

lipid metabolism. It gives an indication as to whether cholesterol is likely to be deposited in the arteries or not (Genest *et al.*, 1999). A ratio with lowered LDL-c as noticed in the present study considered beneficial as protecting from atherosclerosis. The hypocholesterolemia showed in the present study in rats treated with black cumin seed was in consistent with other findings in rats (Zaoui *et al.*, 2002; El-Dakhakhny *et al.*, 2000; Bamosa *et al.*, 1997; Kocyigit *et al.*, 2009), human (Dehkordi and Kamkhah, 2008) layers (Akhtar *et al.*, 2003) and rabbits (Nader *et al.*, 2010; Tousson *et al.*, 2011). In addition, the hypocholesterolemia showed in the present study in rats treated with turmeric was in accordance with other findings in rats (Hussain, 2002; Soliman, 2005; Kim and Kim, 2010). The pronounced hypocholesterolemia reported in the present study in rats administered the medicinal plant combination was not reported before. The hypocholesterolemia showed in the present study in rats treated with black cumin seed and/or turmeric might be attributed to the control of the microsomal 7 α -hydroxylation (the rate limiting enzyme in cholesterol catabolism). The pronounced effect of plant combination on cholesterol metabolism might be resulted from a synergistic effect between active ingredients of both plants. The hypolipidemic and hypocholesterolemic effect of both plants are probably related to decreased dietary cholesterol absorption increased primary bile acid synthesis and its fecal losses. This can be done also by the active ingredient of each plant which may act by making liver cells more efficient to remove LDL-C from blood through increasing LDL receptor densities in liver and by binding to apolipoprotein B (El-Beshbishy *et al.*, 2006; Weggemans and Trautwein, 2003).

CONCLUSION

The present study concluded that plant combination was safe to the animals as reflected on unchanged liver and kidney function biomarkers. Black cumin seed and Turmeric has hypocholesterolemic effect whenever administered alone or in combination (1% each) whereas the hypolipidemic effect was related to the combined administration of both plants only in the diet of rats.

ACKNOWLEDGEMENTS

The researchers thank Saudi Basic Industries Corporation (SABIC) and the Deanship of Scientific Research, King Faisal University for financial support (Grant No.: 116004).

Table 5: Effect of oral administration of turmeric and/or black cumin seed for 6 weeks on blood biochemistry in rats

Blood biochemistry	Group I	Group II	Group III	Group IV
Total protein (g L ⁻¹)	6.8±0.4	7.9±0.3*	7.4±0.1*	6.7±0.2
Albumin (g L ⁻¹)	3.1±0.2	3.3±0.3	3.5±0.1	3.0±0.2
Globulin (g L ⁻¹)	3.7±0.2	4.6±0.2*	3.9±0.2	3.7±0.2
A/G ratio	0.8±0.2	0.7±0.2	0.9±0.2	0.8±0.2
TAG (mg dL ⁻¹)	106.4±6.2	104.9±6.2	102.3±4.2	85.6±4.2*
ALT (U L ⁻¹)	17.9±0.2	18.2±0.2	18.6±0.2	19.1±0.2
AST (U L ⁻¹)	102.0±1.1	99.0±1.2	101.0±1.1	100.0±1.3
Total cholesterol (mg dL ⁻¹)	179.6±1.2	167.0±1.2*	170.0±2.2*	151.3±4.2**
HDL-c (mg dL ⁻¹)	54.2±2.9	55.0±3.7	55.0±3.2	57.6±5.6
LDL-c (mg dL ⁻¹)	104.1±2.1	91.5±1.5*	94.5±2.1*	73.7±0.9**
VLDL-c (mg dL ⁻¹)	21.3±1.2	21.0±1.2	20.5±1.2	20.0±1.2
Uric acid (mg dL ⁻¹)	1.2±0.1	1.3±0.1	1.1±0.1	1.1±0.2
Creatinine (mg dL ⁻¹)	0.3±0.1	0.3±0.1	0.3±0.1	0.3±0.1

I (control), II (black cumin seed treated group), III (turmeric treated group), IV (black cumin seed and turmeric treated group), values are mean±SD of 6 rats; *Means within the same column are statistically significant when compared to control (Group I) at p<0.05; **Means within the same column are statistically significant when compared to other group at p<0.05

REFERENCES

- AOAC, 1995. Official Methods of Analysis. 16th Edn., Association of Official Analytical Chemists, Washington, DC., USA.
- Abd El-Aal, E.S.M. and R.S. Attia, 1993. Characterization of black cumin (*Nigella sativa*): Chemical composition and lipid. Alex. Sci. Exch., 14: 467-467.
- Abu-Al-Basal, M.A., 2011. Influence of *Nigella sativa* fixed oil on some blood parameters and histopathology of skin in staphylococcal-infected BALB/c mice. Pak. J. Biol. Sci., 14: 1038-1046.
- Agarwal, P., M.D. Kharya and R. Shrivastava, 1979. Antimicrobial and anthelmintic activities of the essential oil of *Nigella sativa* L. Ind. J. Exp. Biol., 17: 1264-1275.
- Aggarwal, B.B., A. Kumar, M.S. Aggarwal and S. Shishodia, 2005. Curcumin Derived From Turmeric (*Curcuma longa*): A Spice for all Seasons. In: Phytopharmaceuticals in Cancer Chemoprevention, Preuss, H.G. (Eds). CRC Press Inc., Boca Raton, USA, pp: 349-387.
- Akhtar, M.S., Z. Nasir and A.R., Abid, 2003. Effect of feeding powdered *Nigella sativa* L. seeds on poultry egg production and their suitability for human consumption. Veterinarski Arch., 73: 181-190.
- Alsaif, M.A., 2008. Effect of *Nigella sativa* oil on metabolic responses to prolonged systemic injury in rats. J. Boil. Sci., 8: 974-983.
- Badari, O.A., A.B. Abdel-Naim, M.H. Abdel-Wahab and F.M. Hamada, 2000. The influence of thymoquinone on doxorubicin-induced hyperlipidemic nephropathy in rats. Toxicology, 143: 219-226.
- Bamosa, A.O., B.A. Ali and S.A. Sowayan, 1997. Effect of oral ingestion of *Nigella sativa* seeds on some blood parameters. Saudi Pharm. J., 5: 126-129.
- Basavaraj, M., V. Nagabhushana, N. Prakash, M.M. Appannavar, P. Waggmare and S. Mallikarjunappa, 2011. Effect of dietary supplementation of *Curcuma longa* on the biochemical profile and meat characteristics of broiler rabbits under summer stress. Vet. World, 4: 15-18.
- Bashandy, S.A.E., 1996. Effect of *Nigella sativa* oil on liver and kidney functions of adult and senile rats. Egypt. J. Pharm. Sci., 37: 313-327.
- Bauer, J.D., 1982. Clinical Laboratory Methods. 9th Edn., The C.V. Company II 1830, Westline Industrial, Missouri.
- Buriro, M.A. and M. Tayyab, 2007. Effect of *Nigella sativa* on lipid profile in albino rats. Gomal J. Med. Sci., 5: 28-31.
- Chaudary, S.A., 1996. Serum lipid profile in albino rats fed on atherogenic supplemented palm oil diet and *Nigella sativa*. Ph.D. Thesis, Department of Pathology, Postgraduate Medical Institute, Lahore, Pakistan.
- Chou Y.H., 1975. Experimental Design and the Analysis of Variance, Statistical Analysis-I. Holt, Reinhart and Winston Publication, New York, pp: 340-352.
- Dehkordi, F.R. and A.F. Kamkhah, 2008. Antihypertensive effect of *Nigella sativa* seed extract in patients with mild hypertension. Fundam. Clin. Pharmacol., 22: 447-452.
- Doumas, B.T., D.D. Bayson, R.J. Carter, T. Peters and R. Schaffer, 1981. Estimation of total serum protein. Clin. Chem., 27: 1642-1643.
- Ekanem, J.T. and O.K. Yusuf, 2008. Some biochemical and haematological effects of black seed (*Nigella sativa*) oil on *T. brucei*-infected rats. Afr. J. Biomed. Res., 11: 79-85.
- El-Bagir, N.M., I.T.O. Farah, A. Alhaidary, H.E. Mohamed and A.C. Beynen, 2010. Clinical laboratory serum values in rabbits fed diets containing black cumin seed. J. Anim. Vet. Adv., 9: 2532-2536.
- El-Bahr, S.M. and T.T. Saad, 2008. Effect of black cumin seeds (*Nigella sativa*) and/or Turmeric (*Curcumin*) on hematological, biochemical and immunological parameters of *Mugil cephalus* fish vaccinated with *Aeromonas hydrophila* bacterin. Proceedings of the 13th Scientific Congress, November 25-28, 2008, Faculty of Veterinary Medicine, Assuit University, pp: 365-388.
- El-Bahr, S.M., 2007. Effect of black cumin seeds (*Nigella sativa*) on the profile of serum lipids, lipoproteins and fatty acids in pekin ducklings. Int. J. Applied Chem., 3: 221-230.
- El-Bahr, S.M., M.A. Korshom, A.E.A. Mandour, A.A. El-Bessomy and M.A. Lebdah, 2007. The protective effect of Turmeric on iron overload in albino rats. Egypt. J. Biochem. Mol. Biol., 25: 94-113.
- El-Beshbishy, H.A., AN.B. Singab, J. Sinkkonen and L. Pihlaja, 2006. Hypolipidemic and antioxidant effects of *Morus alba* L. (Egyptian mulberry) root bark fractions supplementations in cholesterol fed rats. Life Sci., 78: 2724-2733.
- El-Dakhkhny, M., N.I. Mady and M.A. Halim, 2000. *Nigella sativa* L. oil protects against induced hepatotoxicity and improves serum lipid profile in rats. Arzneimittelforschung, 50: 832-836.
- Emadi, M., H. Kermanshahi and E. Maroufyan, 2007. Effect of varying levels of turmeric rhizome powder on some blood parameters of broiler chickens fed corn soybean meal based diets. Int. J. Poultr. Sci., 6: 345-348.

- Genest, J.J., M. Marcil, M. Denis and L. Yu, 1999. High density lipoproteins in health and in disease. J. Investig. Med., 47: 31-42.
- Gilani, A.H., Q.Jabeen and M.A.U. Khan, 2004. A review of medicinal uses and pharmacological activities of *nigella sativa*. Pak. J. Biol. Sci., 7: 441-451.
- Gottfried, S.P. and B. Rosenberg, 1973. Improved manual spectrophotometric procedure for determination of serum triglycerides. Clin. Chem., 19: 1077-1078.
- Hasan, M.M., S.P. Chowdhury, Shahidul Alam, B. Hossain and M.S. Alam, 2005. Antifungal effects of plant extracts on seed-borne fungi of wheat seed regarding seed germination, Seedling health and vigour index. Pak. J. Biol. Sci., 8: 1284-1289.
- Hassanin, N.I. and F.M. Hassan, 1996. A preliminary study on the effect of *Nigella sativa* L. seeds on hypoglycemia. Vet. Med. J. Giza, 44: 699-708.
- Hedayat, S.A., 1995. Effect of *Nigella sativa* (black seeds) extract on some haematological and biochemical parameters in rats. Alex. J. Vet. Sci., 11: 95-99.
- Hussain, H.E.M.A., 2002. Hypoglycemic, hypolipidemic and antioxidant properties of combination of curcumin from *curcuma longa* Linn. and partially purified product from *Abroma augusta*, Linn. in streptozotocin induced diabetes. Indian J. Clin. Biochem., 17: 33-43.
- Ibraheim, Z.Z., 2002. Effect of *Nigella sativa* seeds and total oil on some blood parameters in female doxorubicin volunteers. Saudi Pharm. J., 10: 54-59.
- Jaffar, A.R., J. Babb and A. Movahed, 2004. Optimal management of hyperlipidemia in primary prevention of cardiovascular disease. Int. J. Cardiol., 97: 355-366.
- Kim, M. and Y. Kim, 2010. Hypocholesterolemic effects of curcumin via up-regulation of cholesterol 7 α -hydroxylase in rats fed a high fat diet. Nutr. Res. Pract., 4: 191-195.
- Kocyyigit, Y., Y. Atamer and E. Uysal, 2009. The effect of dietary supplementation of *Nigella sativa* L. on serum lipid profile in rats. Saudi Med. J., 30: 893-896.
- Mahfouz, M., M. El-Dakhakhny, A. Gemel and H. Moussa, 1962. Choleric action of *Nigella sativa* seed oil. Egypt. Pharm. Bull., 44: 225-229.
- Mehta, A., G. Kaur and M. Chintamaneni, 2012. Piperine and quercetin enhances antioxidant and hepatoprotective effect of curcumin in paracetamol induced oxidative stress. Int. J. Pharmacol., 8: 101-107.
- Nader, M.A., D.S. El-Agamy and G.M. Suddek, 2010. Protective effects of propolis and thymoquinone on development of atherosclerosis in cholesterol-fed rabbits. Arch. Pharm. Res., 33: 637-643.
- Nasir, Z. and M.A. Grashorn, 2010. Effects of *Echinacea purpurea* and *Nigella sativa* supplementation on broiler performance, carcass and meat quality. J. Anim. Feed Sci., 19: 94-104.
- Nouzarian, R., S.A. Tabeidian, M. Toghyani, G. Ghalamkari and M. Toghyani, 2011. Effect of turmeric powder on performance, carcass traits, humoral immune responses and serum metabolites in broiler chickens. J. Anim. Feed Sci., 20: 389-400.
- Pal, S., T. Choudhuri, S. Chattopadhyay, A., Bhattacharya, G.K. Datta, T. Das and G. Sa, 2001. Mechanisms of curcumin induced apoptosis of Ehrlich's ascites carcinoma cells. Biochem. Biophys. Res. Commun., 288: 658-665.
- Pourghassem-Gargari, B., V. Ebrahimzadeh-Attary, M. Rafrat and A. Gorbani, 2009. Effect of dietary supplementation with *Nigella sativa* L. on serum lipid profile, lipid peroxidation and antioxidant defense system in hyperlipidemic rabbits. J. Med. Plants Res., 3: 815-821.
- Rathee, P.S., S.H. Mishra and R. Kaushal, 1982. Antimicrobial activity of essential oil, fixed oil and unsaponifiable matter of *Nigella sativa* L. Indian J. Pharm. Sci., 44: 8-10.
- Reddy, A.C.P. and B.R. Lokesh, 1994. Alterations in lipid peroxidation in rat liver by dietary n-3 fatty acids: Modulation of anti-oxidant enzymes by curcumin, eugenol and vitamin E. J. Nutr. Biochem., 5: 181-188.
- Reinhold, R.R., 1953. Determination of serum albumin. Clin. Chem., 21: 1370-1372.
- Reitman, S. and S. Frankel, 1957. A colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminase. Am. J. Clin. Pathol., 28: 56-63.
- Salama, A.F. and S.M. El-Bahr, 2007. Effect of curcumin on cadmium-induced oxidative testicular damage in rats. J. Med. Res. Inst., 28: 167-173.
- Shewita, R.S. and A.E. Taha, 2011. Effect of dietary supplementation of different levels of black seed (*Nigella sativa* L.) on growth performance, immunological, hematological and carcass parameters of broiler chicks. World Acad. Sci. Eng. Technol., 77: 788-794.
- Sivabalan, S. and C.V. Anuradha, 2010. A comparative study on the antioxidant and glucose-lowering effects of curcumin and bisdemethoxycurcumin analog through *in vitro* assays. Int. J. Pharmacol., 6: 664-669.
- Soliman, G.Z.A., 2005. Effect of curcumin, mixture of curcumin and piperine and curcumin (Turmeric) on lipid profile of normal and hyperlipidemic rats. Egypt. J. Hosp. Med., 21: 145-161.

- Srinivasan, K., 2005. Role of spices beyond food flavoring: Nutraceuticals with multiple health effects. *Food Rev. Int.*, 21: 167-188.
- Sugiharto, I., E. Widiastuti and N.S. Prabowo, 2011. Effect of turmeric on blood parameters, feed efficiency and abdominal fat content in broilers. *J. Indonesian Trop. Anim. Agric.*, 36: 21-26.
- Tabacco, A., F. Meiathini, E. Moda and P. Tarli, 1979. Simplified enzymic/colorimetric serum urea nitrogen determination. *Clin. Chem.*, 25: 336-337.
- Tasawar, Z., Z. Siraj, N., Ahmad and M.H. Lashari, 2011. The effects of *Nigella sativa* (Kalonji) on lipid profile in patients with stable coronary artery disease in Multan, Pakistan. *Pak. J. Nutr.*, 10: 162-167.
- Tayyab, M., 1995. Effect of consumption of dietary *elaesis guineensis* (*Palm kernel*) seeds on serum lipid profile in albino rats. Ph.D. Thesis, Department of Pathology, Postgraduate Medical Institute, Lahore, Pakistan.
- Tousson, E., M., El-moghazy and E. El-Atrsh, 2011. The possible effect of diets containing *Nigella sativa* and *Thymus vulgaris* on blood parameters and some organs structure in rabbit. *Toxicol. Ind. Health*, 27: 107-116.
- Weggemans, R.M. and E.A. Trautwein, 2003. Relation between soyassociated isoflavones and LDL and HDL cholesterol concentrations in humans: A meta analyses. *Eur. J. Clin. Nutr.*, 57: 940-946.
- Zak, B., R.C. Dickenman, E.G. White, H. Burnett and P.J. Cherney, 1954. Rapid estimation of free and total cholesterol. *Am. J. Clin. Pathol.*, 24: 1307-1315.
- Zaoui, A., Y. Cherrah, K. Alaoui, N. Mahassine, H. Amarouch and M. Hassar, 2002. Effects of *Nigella sativa* fixed oil on blood homeostasis in rat. *J. Ethnopharmacol.*, 79: 23-26.