

## Influence of Selenium on Hepatic Antioxidant Capacity in Ducklings Intoxicated with Aflatoxin B<sub>1</sub>

<sup>1</sup>D. Y. Shi, <sup>2</sup>S. Q. Liao, <sup>1</sup>R. S. Su, <sup>1</sup>R. Q. Lin, <sup>1</sup>Y. C. Lin, <sup>1</sup>Z. X. Tang and <sup>1</sup>S. N. Guo  
<sup>1</sup>College of Veterinary Medicine, South China Agricultural University,  
<sup>2</sup>Guangdong Academy of Agricultural Sciences, Institute of Veterinary Medicine,  
510640 Guangzhou, Guangdong Province, China

**Abstract:** To investigate the influence of selenium on hepatic antioxidant capacity in ducklings administrated with Aflatoxin B<sub>1</sub> (AFB<sub>1</sub>), ninety 7 days old ducklings were randomly divided into three groups (groups I-III). Group I was used as a blank control. Group II was administered with AFB<sub>1</sub> (0.1 mg kg<sup>-1</sup> body weight). Group III was administered with AFB<sub>1</sub> (0.1 mg kg<sup>-1</sup> body weight) plus selenium (sodium selenite, 1 mg kg<sup>-1</sup> body weight). All treatments were given once daily for 21 days. The results showed that the activity of hepatic Superoxide Dismutase (SOD), Catalase (CAT), Glutathione Peroxidase (GSH-Px) and Glutathione Reductase (GR) in group II ducklings significantly decreased when compared with group I (p<0.01). Furthermore, the content of hepatic Malondialdehyde (MDA) significantly increased (p<0.01). However, the activities of hepatic SOD, CAT, GSH-Px and GR in group III ducklings significantly increased when compared with group II (p<0.05). In addition, the content of hepatic MDA significantly decreased (p<0.01). These results revealed that AFB<sub>1</sub> significantly induced hepatic antioxidant function dysfunction. However, selenium could significantly alleviate the negative effect induced by AFB<sub>1</sub>.

**Key words:** Selenium, hepatic antioxidant capacity, aflatoxin B<sub>1</sub>, activity, ducklings, dysfunction

---

### INTRODUCTION

Aflatoxin B<sub>1</sub> is produced by toxigenic fungi belonging to the genus *Aspergillus* and has a long history of association with illness in domesticated animals and in humans (Wogan, 1999; Bondy and Pestka, 2000). The liver is especially sensitive to AFB<sub>1</sub>.

In a number of studies, the AFB<sub>1</sub> is firstly a hepatotoxin causing an excessive build-up of hepatic lipids with enlargement of the liver, proliferation of the biliary ducts (Adav and Godinwar, 1997) and hepatocellular carcinoma (Hamilton, 1978). Recently, the direct evidence of the involvement of free radicals in AFB<sub>1</sub> toxicity was demonstrated by Towner *et al.* (2003) who identified free radicals *in vivo* in rat bile following AFB<sub>1</sub> administration.

It was reported that AFB<sub>1</sub>-mediated hepatic lipid peroxidation were reduced by the pretreatment of rats with antioxidants, selenium and vitamin E (Shen *et al.*, 1996). The aim of the present study was designed to investigate the effect of selenium on hepatic antioxidant capacity in ducklings intoxicated with AFB<sub>1</sub>.

### MATERIALS AND METHODS

**Drugs and chemicals:** Aflatoxin B<sub>1</sub> and dimethyl sulfoxide were purchased from Sigma. Detection kit which includes superoxide dismutase, catalase, glutathione peroxidase, glutathione reductase and malondialdehyde were purchased from Nanjing Jiancheng Bioengineering Institute. All other chemicals were of analytical grade.

**Animals and treatments:** All experimental protocols were approved by the Ethics Committee of South China Agricultural University. This study was carried out on 90 Guangdong white ducklings weighing 180-200 g body weight. About 7 days old ducklings were obtained from the Research Center of Experimental Animals at South China Agricultural University.

The animals were randomly divided into three equal groups containing 30 ducklings with an equal number of male and female ducklings. Group I was used as control and intragastrically administered with Dimethyl Sulfoxide (DMSO). Group II was intragastrically administered with AFB<sub>1</sub> (0.1 mg kg<sup>-1</sup> body weight). Group III was intragastrically administered with AFB<sub>1</sub> (0.1 mg kg<sup>-1</sup>

body weight) plus selenium (sodium selenite, 1 mg kg<sup>-1</sup> body weight). These treatments were administrated once daily for a period of 21 days under the same condition. AFB<sub>1</sub> was diluted with DMSO for experimental ducklings. All ducklings had free access to water and food at room temperature during the study.

**Liver sample preparation:** On 7th, 14th and 21st days after treatment in every experimental group, five ducklings were randomly taken out to obtain, respectively about 1 g of liver and prepare liver homogenate in physiological saline. The supernatant fluid was taken out to detect after liver homogenate was centrifuged at 1,000×g for 5 min. All the steps are strictly operated on ice to guarantee the isolation of high-quality liver homogenate supernatant fluid. The protein concentration of the final liver homogenate supernatant fluid was determined with the Bradford's assay using bovine serum albumin as a protein standard (Bradford and Dodd, 1977).

**Assessment of hepatic antioxidant capacity:** The activities of hepatic superoxide dismutase, catalase, glutathione peroxidase and glutathione reductase were detected according to instruction in the detection kit. Similarly, the content of hepatic malondialdehyde was detected. The activities of hepatic superoxide dismutase, catalase, glutathione peroxidase and glutathione reductase are expressed as international units/mg hepatic protein. The content of malondialdehyde is expressed as nmol/mg hepatic protein.

**Statistical analysis:** The statistical significance of differences between groups in these studies was determined using a one-way analysis of variance and the results were presented as the mean±SE. The significance level was p<0.05.

**RESULTS AND DISCUSSION**

As shown in Table 1, the activity of hepatic Superoxide Dismutase (SOD) was significantly affected after ducklings were intragastrically administered with AFB<sub>1</sub> for 7, 14 and 21 days. The activity of SOD decreased by 17.19, 43.73 and 43.27%, respectively (p<0.01). However, in group III ducklings intragastrically administered with AFB<sub>1</sub> plus selenium, the activity of SOD increased by 13.46, 60.86 and 62.85%, respectively, compared with group II (p<0.01).

As shown in Table 2, the activity of hepatic Catalase (CAT) was significantly affected after ducklings were

intragastrically administered with AFB<sub>1</sub> for 7, 14 and 21 days. The activity of CAT decreased by 31.91, 37.29 and 55.43%, respectively (p<0.01). However, in group III ducklings intragastrically administered with AFB<sub>1</sub> plus selenium, the activity of CAT increased by 36.88, 39.36 and 84.77%, respectively compared with group II (p<0.01).

As shown in Table 3, the activity of hepatic Glutathione Peroxidase (GSH-Px) was significantly affected after ducklings were intragastrically administered with AFB<sub>1</sub> for 7, 14 and 21 days. The activity of GSH-Px decreased by 32.42, 34.71 and 39.88%, respectively (p<0.01). However, in group III ducklings intragastrically administered with AFB<sub>1</sub> plus selenium, the activity of GSH-Px increased by 27.92, 46.14 and 53.98%, respectively compared with group II (p<0.05).

As shown in Table 4, the activity of hepatic Glutathione Reductase (GR) was significantly affected after ducklings were intragastrically administered with AFB<sub>1</sub> for 7, 14 and 21 days. The activity of GR decreased by 27.49, 33.09 and 38.67%, respectively (p<0.01).

Table 1: The activity of hepatic Superoxide Dismutase (SOD) for 7, 14 and 21 days after ducklings administrated with AFB<sub>1</sub>

Three different times after treatment (days)			
Groups	7	14	21
I	282.40±21.26	267.53±19.30	229.31±18.98
II	232.16±16.42 <sup>a</sup>	150.54±12.86 <sup>ab</sup>	130.08±9.710 <sup>ab</sup>
III	263.42±19.58 <sup>b</sup>	242.16±17.69 <sup>bb</sup>	211.83±11.42 <sup>bb</sup>

<sup>a</sup>p<0.05, <sup>ab</sup>p<0.01, significantly different, group II compared with group I. <sup>b</sup>p<0.05, <sup>bb</sup>p<0.01, significantly different, group III compared with group II. (the same following)

Table 2: The activity of hepatic Catalase (CAT) for 7, 14 and 21 days after ducklings administrated with AFB<sub>1</sub>

Three different times after treatment (days)			
Groups	7	14	21
I	26.64±2.32	24.27±2.22	22.10±2.240
II	18.14±1.20 <sup>aa</sup>	15.22±1.33 <sup>aa</sup>	9.85±0.890 <sup>aa</sup>
III	24.83±2.30 <sup>bb</sup>	21.21±2.09 <sup>bb</sup>	18.20±1.670 <sup>bb</sup>

Table 3: The activity of hepatic Glutathione Peroxidase (GSH-Px) for 7, 14 and 21 days after ducklings administrated with AFB<sub>1</sub>

Three different times after treatment (days)			
Groups	7	14	21
I	466.22±32.70	409.22±36.48	347.79±30.36
II	315.08±28.01 <sup>aa</sup>	267.17±21.28 <sup>aa</sup>	209.09±18.92 <sup>aa</sup>
III	403.06±40.24 <sup>bb</sup>	390.43±27.79 <sup>bb</sup>	321.95±22.08 <sup>bb</sup>

Table 4: The activity of hepatic Glutathione Reductase (GR) for 7, 14 and 21 days after ducklings administrated with AFB<sub>1</sub>

Three different times after treatment (days)			
Groups	7	14	21
I	22.95±2.25	20.94±1.76	18.00±1.84
II	16.64±1.33 <sup>aa</sup>	14.01±1.17 <sup>aa</sup>	11.04±1.26 <sup>aa</sup>
III	19.53±2.20 <sup>b</sup>	17.95±1.47 <sup>bb</sup>	15.83±1.76 <sup>bb</sup>

Table 5: The content of hepatic Malondialdehyde (MDA) for 7, 14 and 21 days after ducklings administrated with AFB<sub>1</sub>

Groups	Three different times after treatment (days)		
	7	14	21
I	0.65±0.08	0.75±0.06	0.81±0.06
II	1.04±0.04 <sup>aa</sup>	1.38±0.09 <sup>aa</sup>	1.66±0.06 <sup>aa</sup>
III	0.75±0.03 <sup>bb</sup>	0.89±0.06 <sup>bb</sup>	1.02±0.05 <sup>bb</sup>

However, in group III ducklings intragastrically administered with AFB<sub>1</sub> plus selenium, the activity of GR increased by 17.37, 28.12 and 43.39%, respectively compared with group II (p<0.05).

As shown in Table 5, the content of hepatic Malondialdehyde (MDA) was significantly affected after ducklings were intragastrically administered with AFB<sub>1</sub> for 7, 14 and 21 days. The content of MDA increased by 60.00, 84.00 and 104.94%, respectively (p<0.01). However, in group III ducklings intragastrically administered with AFB<sub>1</sub> plus selenium, the content of MDA decreased by 27.88, 35.51 and 38.55%, respectively compared with group II (p<0.01).

It is known that AFB<sub>1</sub> may induce the production of free radicals and/or the reduction of antioxidant defenses (Shen *et al.*, 1996; Leal *et al.*, 1999). Reactive Oxygen Metabolites (ROM) include intracellular thiols (SH), MDA. The MDA production is recognized as an important factor in determining alteration of membrane fluidity (Chen and Yu, 1994; Ferrante *et al.*, 2002) and increase of membrane fragility accompanying final cell death (Halliwell and Chirico, 1993). A few researchers (Abado-Becognee *et al.*, 1998; Yin *et al.*, 1998; Abel and Gelderblom, 1998) reported that some mycotoxins can cause cell membrane damage through the increase of lipid peroxidation. In continuation of these studies, we investigated the effect of AFB<sub>1</sub> on ducklings' hepatic antioxidant capacity in this research. For 7, 14 and 21 days after ducklings administrated with AFB<sub>1</sub>, respectively it showed that the activities of hepatic SOD, CAT, GSH-Px and GR in group II ducklings (administered with AFB<sub>1</sub>) significantly decreased when compared with group I (p<0.01). Furthermore, the content of hepatic MDA in group II significantly increased (p<0.01).

Although, the mechanism underlying the hepatotoxicity of aflatoxins is not fully understood, several reports suggest that toxicity may ensue through the generation of intracellular Reactive Oxygen Species (ROS) like superoxide anion, hydroxyl radical and hydrogen peroxide during the metabolic processing of AFB<sub>1</sub> by cytochrome P450 in the liver (Sohn *et al.*, 2003). These species may attack soluble cell compounds as well as membranes, eventually leading to the impairment of cell functioning and cytolysis (Blaszczyk *et al.*, 2010; Berg *et al.*, 2004). But peroxidative damages induced in the cell are encountered by elaborate defense mechanisms including enzymic and nonenzymic antioxidants. AFB<sub>1</sub>

mediated hepatic lipid peroxidation were reduced by the pretreatment of rats with antioxidants, selenium and Vitamin E. Defense mechanisms are reinforced by increasing dietary intake of antioxidants and micronutrients such as vitamins and Selenium (Se). The importance of Se is characterized by its role as a constituent of several key antioxidants as well as the unique redox characteristics of selenocysteine and its use in antioxidant enzymes such as thioredoxin reductase. The reduction of reactive oxygen metabolites by glutathione peroxidases helps to maintain membrane integrity (Brenneisen *et al.*, 2005). In this study, we assessed the effect of selenium on hepatic antioxidant function in ducklings administrated with AFB<sub>1</sub>. For 7, 14 and 21 days after ducklings administrated with AFB<sub>1</sub>, respectively it showed that the activities of hepatic SOD, CAT, GSH-Px and GR in group III ducklings (administered with AFB<sub>1</sub> plus sodium selenite) significantly increased when compared with group II (p<0.05). In addition, the content of hepatic MDA in group III significantly decreased (p<0.01).

## CONCLUSION

AFB<sub>1</sub> significantly induced hepatic antioxidant function dysfunction. The result showed that the activity of hepatic Superoxide Dismutase (SOD), Catalase (CAT), Glutathione Peroxidase (GSH-Px) and Glutathione Reductase (GR) significantly decreased when compared. Furthermore, the content of hepatic Malondialdehyde (MDA) significantly increased. However, selenium could significantly alleviate the negative effect induced by AFB<sub>1</sub>.

## ACKNOWLEDGEMENTS

This research was supported by the National Natural Science Foundation of China (Grant No. 38071900). D.Y. Shi, S.Q. Liao and R.S. Su contributed equally to this research.

## REFERENCES

- Abado-Becognee, K., T.A. Mobio, R. Ennamany, F. Fleurat-Lessard, W.T. Shier, F. Badria and E.E. Creppy, 1998. Cytotoxicity of fumonisin B1: Implication of lipid peroxidation and inhibition of protein and DNA syntheses. *Arch. Toxicol.*, 72: 233-236.
- Abel, S. and W.C. Gelderblom, 1998. Oxidative damage and fumonisin B<sub>1</sub>-induced toxicity in primary rat hepatocytes and rat liver *in vivo*. *Toxicology*, 131: 121-131.

- Aday, S.S. and S.P. Godinwar, 1997. Effects of aflatoxins B<sub>1</sub> on liver microsomal enzymes in different strains of chickens. *Comp. Biochem. Physiol. C Pharmacol. Environm.*, 118: 185-189.
- Blaszczyk, I., E. Grucka-Mamczar, S. Kasperczyk and E. Birkner, 2010. Influence of methionine upon the activity of antioxidative enzymes in the kidney of rats exposed to sodium fluoride. *Biol. Trace Elem. Res.*, 133: 60-70.
- Berg, D., M.B. Youdim and P. Riederer, 2004. Redox imbalance. *Cell Tissue Res.*, 318: 201-213.
- Bondy, G.S. and J.J. Pestka, 2000. Immunomodulation by fungal toxins. *J. Toxicol. Environ. Health B Crit. Rev.*, 3: 109-143.
- Bradford, H.F. and P.R. Dodd, 1977. Convulsions and activation of epileptic foci induced by monosodium glutamate and related compounds. *Biochem. Pharmacol.*, 26: 253-254.
- Brenneisen, P., H. Steinbrenner and H. Sies, 2005. Selenium, oxidative stress and health aspects. *Mol. Aspects Med.*, 26: 256-267.
- Chen, J.J. and B.P. Yu, 1994. Alterations in mitochondrial membrane fluidity by lipid peroxidation products. *Free Radic. Biol. Med.*, 117: 411-418.
- Ferrante, M.C., R. Meli, G.M. Raso, E. Esposito, L. Severino, G. Di Carlo and A. Lucisano, 2002. Effect of fumonisin B<sub>1</sub> on structure and function of macrophage plasma membrane. *Toxicol. Lett.*, 129: 181-187.
- Halliwell, B. and S. Chirico, 1993. Lipid peroxidation: Its mechanism, measurement and significance. *Am. J. Clin. Nutr.*, 57: 715S-724S.
- Hamilton, P.B., 1978. Fallacies in our understanding of mycotoxins. *J. Food Prot.*, 41: 404-408.
- Leal, M., A. Shimada, F. Ruiz and E.G. de Mejia, 1999. Effect of lycopene on lipid peroxidation and glutathione-dependent enzymes induced by T-2 toxin *in vivo*. *Toxicol. Lett.*, 109: 1-10.
- Shen, H.M., C.Y. Shi, Y. Shen and C.N. Ong, 1996. Detection of elevated reactive oxygen species level in cultured rat hepatocytes treated with aflatoxin B<sub>1</sub>. *Free. Radic. Biol. Med.*, 21: 139-146.
- Sohn, D.H., Y.C. Kim, S.H. Oh, E.J. Park, X. Li and B.H. Lee, 2003. Hepatoprotective and free radical scavenging effects of *Nelumbo nucifera*. *Phytomedicine*, 10: 165-169.
- Towner, R.A., S.Y. Qian, M.B. Kadiiska and R.P. Mason, 2003. *In vivo* identification of aflatoxin-induced free radicals in rat bile. *Free Radic. Biol. Med.*, 35: 1330-1340.
- Wogan, G.N., 1999. Aflatoxin as a human carcinogen. *Hepatology*, 30: 573-575.
- Yin, J.J., M.J. Smith, R.M. Eppley, S.W. Page and J.A. Sphon, 1998. Effects of fumonisin B<sub>1</sub> on lipid peroxidation in membranes. *Biochim. Biophys. Acta*, 1371: 134-142.