

Isolation, Identification and Antioxidant Properties of *Xylaria escharoidea* Associated with Termite Nest

¹Shuangfen Yue, ²Saifei Li, ²Huaan Wen and ¹Li Fan

¹College of Life Science, Capital Normal University, Beijing, China

²State Key Laboratory of Mycology, Institute of Microbiology,
Chinese Academy of Sciences, Beijing, China

Abstract: The strain of *Xylaria* Y-4 was isolated from an active termite nest and identified as *Xylaria escharoidea* based on ITS1-5.8S-ITS2 sequences. Intracellular Polysaccharide (IPS) and Extracellular Polysaccharide (EPS) were extracted from *Xylaria escharoidea* (strain Y-4) in submerged culture. Antioxidant potency was investigated by assays of various established antioxidant *in vitro* systems such as hydroxyl ($\cdot\text{OH}$) and DPPH radical scavenging ability, reducing power and ferrous ion chelating ability. Results revealed that the antioxidant activities of IPS extracts were generally more potent than EPS extracts in *in vitro* assays. IPS exhibited stronger scavenging activity towards DPPH radicals with EC_{50} of 0.75 mg mL^{-1} .

Key words: Free radicals, polysaccharide, reducing power, IPS, *in vitro*

INTRODUCTION

Reactive Oxygen Species (ROS) including hydrogen, peroxidesuperoxide anion, hydroxyl radicals and singlet oxygen is generated by aerobic metabolism of the body (Lee *et al.*, 2004). They can cause protein and lipid oxidation, DNA strand break and base modification, resulting in various diseases and disorders such as cancer, cardiovascular diseases, atherosclerosis and aging (Wu and Hansen, 2008; Xu *et al.*, 2009). Almost all organisms are well protected against free radical damage by oxidative enzymes such as superoxide dismutase, catalase and chemical compounds (Niki *et al.*, 1994). However, these systems are frequently insufficient to totally prevent the damage, resulting in diseases and accelerated ageing (Ames *et al.*, 1993). Some synthetic antioxidants such as Butylated Hydroxy-Anisole (BHA) and Butylated Hydroxytoluene (BHT) have been used in preservation of foods but several studies suggest that they could promote tumour formation (Cheung *et al.*, 2003). Therefore, considerable interest has arisen in looking for natural antioxidants.

Fungus combs are the fecal-based fungus gardens built by macrotermitine termites in their nests where they cultivate mycelia of the genus *Termitomyces*. Some species of *Xylaria* are often found in the fungus combs abandoned by termites and *Xylaria nigripes* (Koltz.) Sacc., a chinese traditional medicine was studied widely in China. *Xylaria nigripes*, also known as Wu Ling Shen, is known to enhance immunity and hematopoiesis (Xu, 1997). It is also used for treating insomnia, trauma

and as a diuretic and nerve tonic (Dai and Yang, 2008; Ma *et al.*, 1999; Xu, 1997). Moreover, other studies have shown that it possesses good antioxidant and hepatoprotective activities (Ko *et al.*, 2009; Song *et al.*, 2011; Wu, 2001). Another species *Xylaria gracillima* (Fr.) Fr., also found in abandoned nest of termites was reported to have anti-tumor activities (Dong, 1998) and antioxidant activities (Li and Wen, 2008).

Xylaria escharoidea (Berk.) Fr. is a species undoubtedly associated with termite nest. So far, there has been no official report on the antioxidant activities of *X. escharoidea*. The aim of this research was to study the antioxidant properties of extracts from *X. escharoidea* (Y-4) in submerged culture.

MATERIALS AND METHODS

Chemicals and reagents: Ferrozine, potassium ferricyanide, ferric trichloride, absolute ethyl alcohol, 1,1-Diphenyl-2-Picryl-Hydrazyl (DPPH), sodium salicylate, ferrous sulfate, Hydrogen peroxide (H_2O_2), ascorbic acid (VC), monosodium phosphate, disodium hydrogen phosphate, ethanoic acid, sodium acetate. All chemicals and solvents used in this study were of analytical grade available commercially.

Fungal isolation and identification: The fungus comb sample was collected from an active termite nest in Changsha, Hunan Province, China, Sep. 2011. When back to the laboratory, fungus comb was incubated at room temperature in a moist chamber where filter paper soaked

in sterile demineralized water was added to keep moist environment. After 5-7 days, *Xylaria* appeared and was inoculate on Potato Dextrose Agar (PDA) media to obtain pure cultures. The strain of *Xylaria* Y-4 was isolated, purified and preserved on PDA slant at 4°C. Cultures and anamorph of *Xylaria* Y-4 researchers obtained in this study had colony and anamorphic features much like those *X. escharoidea* strains described by Ju and Rogers (1999).

Phylogenetic analyses: Total genomic DNA was extracted from fungal mycelia using the Cetyl Trimethyl Ammonium Bromide (CTAB) Method (O'Donnell *et al.* 1997). Primers ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') (White *et al.*, 1990) were used in amplifying the regions of ITS1-5.8S-ITS2. DNA fragment was amplified and sequenced using the method described by Qian *et al.* (2011). The other sequence data of ITS rDNA included in this study were downloaded from GenBank.

DNA sequences used in this study were aligned using the alignment program ClustalX Version 1.81 (Thompson *et al.*, 1997). The alignment was manually adjusted with Se-Al v.2.03a (Rambaut, 2000). The aligned dataset was analyzed with Maximum Parsimony (MP) using PAUP×4.0b10 (Swofford, 2002). Maximum parsimony analysis was conducted using heuristic searches with 1000 replicates of random-addition sequence, Tree Bisection Reconnection (TBR) branch swapping algorithm. All characters were equally weighted and unordered. Gaps were treated as missing data to minimize homology assumptions. A Bootstrap (BS) analysis was performed with 1000 replicates, each with 10 random taxon addition sequences. TBR branch swapping was employed.

Fungal cultivation: Strain Y-4 was cultured on a PDA agar plate for 5-6 days at 25°C. Four active mycelial agar discs (diameter 0.4 cm) were inoculated into a 250 mL erlenmeyer flask containing 100 mL submerged medium of basic medium (pH 6.0) including 2% glucose, 1% peptone, 0.1% KH₂PO₄, 0.1% MgSO₄ and then incubated at 25°C, 150 r min⁻¹ for 1 week on a rotary shaker.

Extraction of water soluble polysaccharides: Mycelia of the cultured fungus were filtered through a filter paper and then washed with distilled water. After lyophilization, the dried mycelia were boiled with distilled water at 100°C, 2 h for 3 times and the supernatant was condensed and was mixed with ethanol (95%, v/v) at 4°C overnight for 3 times, to obtained Intracellular Polysaccharide (IPS). The yield of IPS from dried mycelia was 13.68 mg g⁻¹. To extract Extracellular Polysaccharide (EPS), the residual

fermentation broth was condensed by evaporator and mixed with four volumes of ethanol (95%, v/v) at 4°C, 18 h for 3 times. Then, supernatant was discarded after centrifugation at 3000 rpm for 10 min. The yield of EPS from fermentation broth was 6.51 g L⁻¹. IPS and EPS were lyophilized to powder. The two kinds of crude polysaccharide were redissolved in distilled water for determination of antioxidant activity.

Hydroxyl radical scavenging assay: Hydroxyl radical scavenging activity was measured by a modified Smirnov and Cumbes (1989)'s Method. The reaction mixture, containing 0.3 mL sodium salicylate (20 mmol L⁻¹), 1 mL ferrous sulfate (1.5 mmol L⁻¹), 0.1 mL different samples (10-90 mg mL⁻¹) and 0.7 mL H₂O₂ (6 mmol L⁻¹) was incubated in a water bath at 37°C for 1 h. The absorbance was measured at 510 nm using Microplate Reader and VC was used for comparison. The capability of scavenging hydroxyl radical was calculated using the equation:

$$\text{Scavenging effect (\%)} = \frac{A_0 - A_1}{A_0} \times 100$$

Where:

A₀ = Absorbance of the control (blank, without samples)

A₁ = Absorbance of the samples

Reducing power assay: The reducing power of the extracts was determined according to the method described earlier by Yen and Chen (1995) with slight modifications. About 1 mL of each extract (10-90 mg mL⁻¹) in deionised water was mixed with 2.5 mL of sodium phosphate buffer (200 mmol L⁻¹, pH 6.6) and 2.5 mL of 1% potassium ferricyanide. The mixture was incubated at 50°C for 20 min and then rapidly cooled. After 2.5 mL of 10% trichloroacetic acid were added, the mixture was centrifuged at 4000 r min⁻¹ for 10 min, 2.5 mL of the supernatant was mixed with 5 mL of distilled water and 0.5 mL of ferric chloride (0.1%) for 10 min. The absorbance was measured at 700 nm and VC was used for positive control. A higher absorbance indicates a higher reducing power.

DPPH radical scavenging assay: The effect of extracts on DPPH radical was estimated according to the method of Blois (1958). The 0.1 mL sample was mixed with 2.0 mL of acetic acid buffer solution (0.05 mol L⁻¹, pH 5.5), 1.9 mL of anhydrous ethanol and 1 mL DPPH solution (0.5 mmol L⁻¹) and the mixture was left to stand in dark at room temperature for 30 min. The absorbance was measured at 517 nm. VC and distilled water were used as positive control and blank control, respectively. The scavenging ability was calculated by using the equation:

$$\text{Scavenging ability (\%)} = \frac{[A_0 - (A_1 - A_2)]}{A_0} \times 100\%$$

Where:

A_0 = Absorbance of the control (blank, without samples)

A_1 = Absorbance of the samples

A_2 = Absorbance of the samples and the other solvent (without DPPH)

Ferrous ion chelating activity assay: The chelating activity of ferrous ions by the extracts was estimated by the methods of Decker and Welch (1990). Briefly, 1 mL sample (10-90 mg mL⁻¹) was added to 3.7 mL distilled water and 0.1 mL FeCl₂ (2.0 mmol L⁻¹). The reaction was initiated by the addition of 0.2 mL ferrozine (5 mmol L⁻¹) and the mixture was shaken vigorously and left standing at room temperature for 10 min. The absorbance of the resulting solution was then measured spectrophotometrically at 562 nm. Distilled water was used as blank control and VC was used as positive control. The chelating activity of the samples for ferrous ion was calculated as:

$$\text{Chelating rate (\%)} = \frac{A_0 - A_1}{A_0} \times 100\%$$

Where:

A_0 = Absorbance of the control (blank, without samples)

A_1 = Absorbance of the presence of samples

Statistical analysis: All data were expressed as Mean±(SD) for at least triplicate analyses on the same sample. Analysis of variance was performed by ANOVA procedures using SPSS 11.5 (SPSS Inc., Chicago, IL, USA). Significant differences between means were determined by Duncan's multiple range tests (p<0.05).

RESULTS

Phylogenetic analysis: Strain Y-4 was isolated from the fungus comb with the method of *in situ* cultivation and exhibited 99% similarity with *Xylaria escharoidea* (EU179864) (Ju and Hsieh, 2007) in GenBank database based on ITS1-5.8S-ITS2 sequences. Phylogenetic analysis (Fig. 1) based on ITS1-5.8S-ITS2 regions shows




Fig. 1: Phylogenetic tree constructed with the maximum parsimony method according to *rDNA-ITS* (ITS1-5.8S-ITS2) gene evolutionary distance among the strain of Y-4 and other termite-associated *Xylaria* species. *Annulohypoxylon cohaerens* (EF026140), *Biscogniauxia arima* (EF026150) and *Biscogniauxia mediterranea* (AF326479) were used as out-groups. GenBank accession numbers are given in parentheses. The confidence values from 1,000 replicate bootstrap samplings were shown at each node




Fig. 2: Hydroxyl radical scavenging activity of IPS and EPS from *X. escharoidea* (strain Y-4) in submerged culture. Each value is the Mean±SD of triplicate measurements

the sequence of Y-4 also grouped in one clade with the epitype sequence of *X. escharoidea* (EU179864) with a strong support (BP = 100). So, strain Y-4 was identified as *X. escharoidea*.

Hydroxyl radical scavenging activity: Hydroxyl radical is known as the strongest reactive oxygen radical and can induce severely damage to the adjacent biomolecules like DNA, lipids and proteins (Spencer *et al.*, 1994). Figure 2 shows that IPS and EPS from *X. escharoidea* (strain Y-4) exhibited excellent and concentration dependent scavenging abilities toward hydroxyl radicals. At a concentration of 3 mg mL⁻¹, the scavenging ability of IPS and EPS against hydroxyl radical reached 78.54 and 79.42%, respectively. In general, IPS had almost the same power with EPS at all tested concentrations. However, based on EC₅₀ values, IPS (0.70 mg mL⁻¹) was more effective than EPS (1.13 mg mL⁻¹) on scavenging hydroxyl radical. The antioxidant properties are inversely correlated with their EC₅₀ values and <10 mg mL⁻¹ are indicative of the effective antioxidant activity (Lee *et al.*, 2007).

Reducing power: The reducing capacities are generally associated with the presence of reductants (antioxidants). The presence of reductants in the samples would result in the reducing of Fe³⁺ to Fe²⁺ by donating an electron. Amount of Fe²⁺ complex can be then be monitored by measuring the formation of Perl's Prussian blue at 700 nm. Increasing absorbance at 700 nm indicates an increase in reductive ability. The reducing power of ISP and ESP extracted from *X. escharoidea* (strain Y-4) were shown in Fig. 3. Higher absorbance value means stronger reducing power of samples. Apparently, the reducing power of all samples was moderate. However, it is worthwhile to note




Fig. 3: Reducing power of IPS and EPS from *X. escharoidea* (strain Y-4) in submerged culture. Each value is the Mean±SD of triplicate measurements




Fig. 4: DPPH radical scavenging activity of IPS and EPS from the *X. escharoidea* (strain Y-4) in submerged culture. Each value is the Mean±SD of triplicate measurements

that IPS and EPS were much stronger than that of ascorbic acid when concentration above 30 mg mL⁻¹. The EC₅₀ values of IPS and EPS were 1.36 and 1.23 mg mL⁻¹, respectively which reveals that the reducing power of EPS are higher than that of IPS (Table 1).

DPPH radical scavenging activity: The model of scavenging DPPH radical is based on the reduction of the absorbance of methanolic DPPH solution at 517 nm in the presence of a proton-donating substance. The method is widely used to evaluate the free radical-scavenging capacities of samples (Soare *et al.*, 1997; Naik *et al.*, 2003). The scavenging abilities of water-soluble polysaccharides from the *X. escharoidea* on DPPH correlated well with the increasing of concentrations (Fig. 4). IPS and EPS had almost the same activities on scavenging DPPH radical.

Table 1: EC₅₀ values obtained in the different antioxidant activity assays of IPS and EPS from *X. escharoidea* (strain Y-4) (mg/mL)

Samples	Hydroxyl radicals	Reducing power	DPPH radicals	Ferrous ions
IPS	0.75±0.021 ^b	1.36±0.027 ^a	25.88±1.034 ^c	1.29±0.042 ^a
EPS	1.13±0.036 ^c	1.23±0.045 ^b	27.01±1.22 ^b	2.01±0.108 ^b
VC	0.70±0.005 ^a	<0.2	1.21±0.087 ^a	ND
EDTA	ND	ND	ND	<0.10

EC₅₀ value = The effective concentration at which hydroxyl radicals were scavenged by 50% DPPH radicals were scavenged by 50%, the absorbance was 0.5 and ferrous ions were chelated by 50%, respectively; IPS = Intracellular Polysaccharide; EPS = Extracellular Polysaccharide. Each value is expressed as Mean±Standard Deviation (n = 3). Means with different letters within a column are significantly different (p<0.05); ND = Not Done




Fig. 5: Ferrous ion chelating activity of IPS and EPS from the *X. escharoidea* (strain Y-4) in submerged culture. Each value is the mean±SD of triplicate measurements

EC₅₀ values of IPS and EPS were 25.88 and 27.01 mg mL⁻¹, respectively. At concentration of 90 mg mL⁻¹, the highest inhibition rate of IPS and EPS reached 82.68 and 84.90%, respectively.

Ferrous ion chelating ability: Ferrozine can quantitatively form complexes with ferrous ion. However in the presence of chelating agents, the complex formation is disrupted with the result that the red colour of the complex is decreased. Therefore, measurement of colour reduction can estimate the chelating activity of the coexisting chelator. As shown in Fig. 5, IPS and EPS exhibited obvious chelating activity on ferrous ion in a concentration dependent manner. At concentration of 9 mg mL⁻¹, the chelating effects IPS and EPS were 87.16 and 77.20%. The EC₅₀ values of IPS and EPS were 1.29 and 2.01 mg mL⁻¹. Obviously, the chelating activity of EPS was lower than that of IPS.

DISCUSSION

The present study provides the first analyses for the antioxidant activities of *X. Escharoidea* (strain Y-4)

associated with fungus comb using *in vitro* assays. The water-soluble polysaccharide IPS and EPS of *X. escharoidea* (strain Y-4) exhibited excellent hydroxyl radical scavenging activity and ferrous ion chelating activity. The antioxidant activities of IPS were generally more potent than EPS as antioxidants. In extremely high concentrations, the reducing power of IPS and EPS even was much higher than positive control VC. According to Li and Wen (2008), IPS of *X. gracillima* founded on waste termite nest was more effective than EPS on chelating ferrous ion whereas which was weaker than EPS on scavenging hydroxyl radical.

In ferrous ion chelating assay, the potency of IPS of *X. escharoidea* (strain Y-4) was weaker than that of *X. gracillima* at the same concentration (Li and Wen, 2008). While in hydroxyl radicals scavenging assay, from 0.125-16 mg mL⁻¹, the highest activity of IPS extract from *X. gracillima* was 54.9% (Li and Wen, 2008) which was lower than IPS from *X. escharoidea* (strain Y-4) (58.91%, 1 mg mL⁻¹). Based on EC₅₀ values, the mycelial extract from *Xylaria nigripes* (Koltz.) Sacc. (EC₅₀ 73.49 µg mL⁻¹) (Ko *et al.*, 2009) exhibited stronger scavenging activity of DPPH radical than that from *X. escharoidea* (strain Y-4) (EC₅₀ 25.88 mg mL⁻¹). In addition, the antioxidant and anti-tumor activity of nine compounds from mycelium of *X. nigripes* were studied by Gong *et al.* (2008). The results showed that (2, 6-dihydroxyphenyl)-3-hydroxybutan-one had strong DPPH radical scavenging ability and reducing capacity (Gong *et al.*, 2008).

CONCLUSION

There are few reports about antioxidant activities of EPS from *Xylaria* associated with termite nest. According to Li and Wen (2008), the components extracted from *X. gracillima* exhibited a dose-dependent scavenging activity towards hydroxyl radicals. At concentration of 2 mg mL⁻¹, the DPPH radical scavenging activity of EPS from *X. gracillima* was about 18.79% which was weaker than that of *X. escharoidea* (strain Y-4) (74.15%) at the same concentration. But the chelating ferrous ion activity of EPS from *X. gracillima* was stronger than that of *X. escharoidea* (strain Y-4) at concentration 2 mg mL⁻¹.

ACKNOWLEDGEMENTS

This study was supported by National Natural Science Foundation of China (NNSFC) (No. 31070017). Researchers would like to thank Prof. Zuohong Chen (College of Life Sciences, Hunan Normal University) for his helpful assistance in sampling of termite nest. Huan Wen and Li Fan contributed equally in this research.

REFERENCES

- Ames, B.N., M.K. Shigenaga and T.M. Hagen, 1993. Oxidants, antioxidants and the degenerative diseases of aging. *Proc. Natl. Acad. Sci.*, 90: 7915-7922.
- Blois, M.S., 1958. Antioxidant determinations by the use of a stable free radical. *Nature*, 181: 1199-1200.
- Cheung, L.M., P.C.K. Cheung and V.E.C. Ooi, 2003. Antioxidant activity and total phenolics of edible mushroom extracts. *Food Chem.*, 81: 249-255.
- Dai, Y.C. and Z.L. Yang, 2008. A revised checklist of medicinal fungi in China. *Mycosystema*, 27: 801-824.
- Decker, E.A. and B. Welch, 1990. Role of ferritin as a lipid oxidation catalyst in muscle food. *J. Agric. Food Chem.*, 38: 674-677.
- Dong, Z.L., 1998. Study on the clinic effect for terminal cancer cure of compound capsule from *Xylaria gracillima*. *Sci. Technol. Termites*, 15: 32-33.
- Gong, Q.F., S.H. Wu, N.H. Tan and Z.H. Chen, 2008. Study on compounds with antioxidant and antitumor activity from fermental mycelium of *Xylaria nigripes*. *Food Sci. Technol.*, 33: 28-31.
- Ju, Y.M. and H.M. Hsieh, 2007. *Xylaria* species associated with nests of *Odontotermes formosanus* in Taiwan. *Mycologia*, 99: 936-957.
- Ju, Y.M. and J.D. Rogers, 1999. The xylariaceae of Taiwan (excluding anthostomella). *Mycotaxon*, 73: 343-440.
- Ko, H.J., A. Song, M.N. Lai and L.T. Ng, 2009. Antioxidant and antiradical activities of Wu Ling Shen in a cell free system. *Am. J. Chin. Med.*, 37: 815-828.
- Lee, J., N. Koo and D.B. Min, 2004. Reactive oxygen species, aging and antioxidative nutraceuticals. *Compreh. Rev. Food Sci.*, 3: 21-33.
- Lee, Y.L., M.T. Yen and J.L. Mau, 2007. Antioxidant properties of various extracts from *Hypsizygus marmoreus*. *Food Chem.*, 104: 1-9.
- Li, S.F. and H.A. Wen, 2008. Antioxidant activities of bioactive components from *Xylaria gracillima* in submerged culture. *Mycosystema*, 4: 587-593.
- Ma, Z.Z., P.P. Zuo, W.R. Chen, L.J. Tang, Q. Shen, X.H. Sun and Y. Shen, 1999. Studies on the sedative and sleeping effects of Wuling mycelia and its pharmacological mechanism. *Chin. Pharm. J.*, 34: 374-377.
- Naik, G.H., K.I. Priyadarshini, J.G. Satav, M.M. Banavalikar, D.P. Sohoni, M.K. Biyani and H. Mohan, 2003. Comparative antioxidant activity of individual herbal components used in ayurvedic medicine. *Phytochemistry*, 63: 97-104.
- Niki, E., H. Shimaski and M. Mino, 1994. Antioxidantism-free radical and biological defence. Gakkai Syuppan Center, Tokyo, pp: 3-16.
- O'Donnell, K., E. Cigelnik, N.S. Weber and J.M. Trappe, 1997. Phylogenetic relationships among ascomycetous truffles and the true and false morels inferred from 18S and 28S ribosomal DNA sequence analysis. *Mycologia*, 89: 48-65.
- Qian, Q., S.F. Li and H.A. Wen, 2011. Fungal diversity of fungus comb in termite nests. *Mycosystema*, 30: 556-565.
- Rambaut, A., 2000. Estimating the rate of molecular evolution: Incorporating non-contemporaneous sequences into maximum likelihood phylogenies. *Bioinformatics*, 16: 395-399.
- Smirnoff, N. and Q.J. Cumbes, 1989. Hydroxyl radical scavenging activity of compatible solutes. *Phytochemistry*, 28: 1057-1060.
- Soare, J.R., T.C.P. Dinis, A.P. Cunha and L. Almeida, 1997. Antioxidant activity of some extracts of *Thymus zygis*. *Free Radic. Res.*, 26: 469-478.
- Song, A., H.J. Ko, M.N. Lai and L.T. Ng, 2011. Protective effects of Wu-Ling-Shen (*Xylaria nigripes*) on carbon tetrachloride-induced hepatotoxicity in mice. *Immunopharmacol. Immunotoxicol.*, 33: 454-460.
- Spencer, J.P.E., A. Jenner, O.I. Aruoma, P.J. Evans and H. Kaur *et al.*, 1994. Intense oxidative DNA damage promoted by L-DOPA and its metabolites implications for neurodegenerative disease. *FEBS Lett.*, 353: 246-250.
- Swofford, D.L., 2002. *Phylogenetic Analysis Using Parsimony (and other Methods)*. 1st Edn., Sinauer Associates, Sunderland.
- Thompson, J.D., T.J. Gibson, F. Plewniak, F. Jeanmougi and D.G. Higgins, 1997. The CLUSTAL_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.*, 25: 4876-4882.
- White, T.J., T. Bruns, S. Lee and J. Taylor, 1990. Amplification and Direct Sequencing of Fungal Ribosomal RNA Genes for Phylogenetics. In: *PCR Protocols: A Guide to Methods and Applications*, Innis, M.A., D.H. Gelfand, J.J. Sninsky and T.J. White (Eds.). Academic Press, New York, USA., pp: 315-322.
- Wu, G.F., 2001. A study on DPPH free-radical scavengers from *Xylaria nigripes*. *Acta Microbiol. Sin.*, 41: 363-366.

- Wu, X.J. and C. Hansen, 2008. Antioxidant capacity, phenolic content and polysaccharide content of *Lentinus edodes* grown in whey permeate-based submerged culture. *J. Food Sci.*, 73: M1-M8.
- Xu, J.T., 1997. *Chinese Medicinal Mycology*. Beijing Medical University and Peking Union Medical College Press, Beijing, China, pp: 664-675.
- Xu, W., F. Zhang, Y. Luo, L. Ma, X. Kou and K. Huang, 2009. Antioxidant activity of a water-soluble polysaccharide purified from *Pteridium aquilinum*. *Carbohydr. Res.*, 344: 217-222.
- Yen, G.C. and H.Y. Chen, 1995. Antioxidant activity of various tea extracts in relation to their antimutagenicity. *J. Agric. Food Chem.*, 43: 27-32.