

Bilateral Contracted Flexor Tendon of the Carpal Joint and Congenital Ankylosis of the Humero-Radial Joint in a 72 h Old Heifer Calf: A Case Report

¹G.E. Ochube, ²B.Y. Kaltungo and ³U.B. Abubakar

¹Veterinary Hospital, Gombe, Nigeria

²Veterinary Teaching Hospital, ³Department of Veterinary Medicine,
Faculty of Veterinary Medicine, Ahmadu Bello University, Zaria, Nigeria

Abstract: Congenital anomaly by autosomal recessive genes has been implicated as the main etiology of contracted flexor tendons in musculoskeletal deformities of new borne calves. Surgical intervention of this case revealed that ankylosis was caused by wrong origination and insertion of malformed ulnaris lateralis muscle. A stay in the form of splints was placed throughout the limb. Plasters of Paris were then applied to the whole length of the limb to enhance stability. The surgical site was fenestrated to allow aeration and dressing, the overall result of the procedure was encouraging. After 7 weeks, a second surgery was performed to correct the bilateral contracted flexor tendon of the carpal joint. This was achieved by deep digital flexor tendon tenotomy of the carpal joint and placing a fiber glass cast on both joints.

Key words: Congenital ankylosis, flexor tendon, ulnaris lateralis muscle, plasters of Paris, calf

INTRODUCTION

Congenital and inherited anomalies can result in the birth of diseased or deformed neonates. Congenital disorders can be due to viral infections of the fetus or to ingestion of teratogenic agents by the dam at certain stages of gestation. The Musculoskeletal System can also be affected by certain neurological disorders. Contracted flexor tendons are probably the most prevalent abnormality of the Musculoskeletal System of the new borne calves. The condition is caused by an autosomal recessive gene.

At birth, the pastern and fetlock of the fore limbs and sometimes the carpal joints are flexed to varying degrees due to shortening of the deep and superficial digital flexors and associated muscles. Slightly affected animals bear weight on the soles of the feet and walk on their toes/hooves. More severely affected animals walk on the dorsal surfaces of the joint which become damaged and suppurative arthritis develop. Rupture of the common digital extensor can occur as a sequela (Horney and Amstutz, 1980).

Literature classified the muscles of the ante brachium into extensors and flexors. The extensors are located at the cranio-lateral aspect of the fore arm. The flexors are on the caudo-medial aspect of the fore arm (Riegall and Susan, 2000). Cranio-laterally, the extensors are: extensor carpi radialis, extensor digitorum longus.

Extensor carpi radialis is the most cranial muscles to the lateral surface and largest. It originates from the epicondylar crest of the humerus and inserts on the metacarpal tuberosity. It extends the carpal joint and is innervated by the radial nerve (Chibuzo and Sivachelvan, 1989).

While ulnaris lateralis which is a flexor muscle is a long fusiform muscle which lies between the lateral digital extensor cranially and the ulnar head of the deep digital flexor caudally. The ulnaris lateralis arises from the lateral epicondyle of the humerus. It inserts by two tendons on the accessory carpal bone. It flexes the carpal joint, it is the only flexor of the carpus that is innervated by the radial nerve which usually innervates the extensors of the carpus and digits (Chibuzo and Sivachelvan, 1989).

CASE REPORT

A 72 h old female calf weighing 45 kg belonging to a client was presented to the Veterinary Hospital, Gombe with a twisted left fore limb that appears to be congenitally malformed (Fig. 1). On the physical examination, animal appeared alert and full of energy. Vital parameters such as temperature (39.5°C), pulse rate (37.0 beat min⁻¹), respiratory rate (12 cycles min⁻¹) were taken and these were within normal range. The animal was apparently normal and suckles well.



Fig. 1: Calf at presentation

PLAN OF ACTION

Surgical correction of the ankylosed elbow joint and carpal joint tenotomy at a later date.

SURGICAL PROCEDURE

First surgery: Proper shaving and disinfection of the area was carried out. Acepromazine/Xylazine was given as a muscle relaxant/sedative at a dose of 0.02-0.04 and 0.05-0.08 mg kg⁻¹ i/m, respectively. A cruciate incision was made on the lateral aspect of the elbow joint and extends to the cranial part of the extensor carpi radialis (Von-Veet and Valentine, 2007).

After exposing the joint, it was noticed that the ankylosis was caused by wrong origination and insertion of malformed ulnar lateral muscle. After removal of the rudimentary muscle that was responsible for the ankylosis, the extensor carpi radialis muscle was inverted and closed to give strength to the area after healing. Skin was closed routinely using size 1 nylon. A stay in form of splint was placed throughout the limb (from the distal aspect of the elbow joint to the metacarpal joint). Plaster of Paris (POP) was then applied to the whole length of the limb to enhance stability (Fig. 2). The surgical site was fenestrated for dressing. After 7 weeks first surgery and following good prognosis from post operative follow ups, the second surgery was performed to correct the bilateral contracted flexor tendon.

Second surgery: The calf was anaesthetized and prepared for surgery. A 2.5 cm longitudinal midline incision was made over the palmar aspect of the right pastern just proximal to the bulb of the heels (Wagner *et al.*, 1982). The distal synovial sheath was opened and the deep



Fig. 2: Calf 20 min after first surgery



Fig. 3: The 2 months after second surgery



Fig. 4: Calf ambulating 4 months after second surgery

digital flexor tendon was elevated and incised transversely. The ends of the severed tendons retracted approximately 1 cm and sutured together using size 1 nylon. Fiber glass cast was applied from the metacarpal joint to the pastern joint leaving the surgical site fenestrated for dressing and post operative follow ups. The same procedure was performed for the other limb

(Fig. 3). The 4 months following second surgery, the calf was ambulating well and appeared to have a normal foot axis (Fig. 4).

DISCUSSION

The etiology of this case was probably due to genetic malformation that led to wrong origination and insertion of this muscle leading to ankylosis of the metacarpal joint. The faulty origination emanated from the medial part of the epicondylar crest of the humerus and inserts proximally to the accessory carpal bone therefore shortening the axis/angle of the elevation of the joint. This is in line with research carried out by McIlwraith and Fessler (1978).

As the result of the complexity of the above surgical procedure, it became necessary to do the surgery in two phases. Because of the nature of the ankylosed elbow joint, researchers opted for that as the first procedure. This enhanced the straightening of the limb.

After this, the limb was fixed using splints and plaster of Paris. The animal was left to recover fully and a tentative date (7 weeks after first surgery) was agreed upon for the second surgery which was also a success. After 6 weeks following first surgery and good prognosis noticed, the cast was removed in preparatory for second surgery. Following the second surgery, the animal showed signs of pain and discomfort, this was managed with oral administration of phenylbutazone (4.4 mg kg⁻¹, orally bid). After 3 months of second surgery, the calf was ambulating well and there was absence of pain when digital pressure was applied to the joint. The fiber cast was removed a month later.

In majority of affected horses/cattle, deep digital flexor tendon deformities can be corrected by non surgical treatment or by inferior check ligament desmotomy (Sonnichsen, 1982). However, these procedures may not produce a normal hoof axis when the dorsal surface of the hoof wall is flexed past vertical (McIlwraith and Fessler, 1978). According to Fackelman *et al.* (1983), deep digital flexor tenotomy has been reported in horses as a treatment for severe inter-phalangeal joint flexural deformity.

CONCLUSION

Surgical intervention has proved very useful in managing this case although cumbersome. The post operative care is very intensive and requires slinging the animal to enable it to stand most of the time. Deep digital flexor tendon is a relatively simple surgical procedure but post operative care of uttermost importance in the

outcome. Genetically influenced deformities like this can be prevented with proper breeding methods in place where suspected animals with bad traits are culled (Leipold *et al.*, 1993).

ACKNOWLEDGEMENTS

Researchers sincerely appreciate the contributions of the following Dr's Tosin Oluwadare, Chinyere Nbanefor, staff and management of Gombe State Veterinary Hospital, the Government and people of Gombe State for providing the enabling environment that made this research possible. Finally, researchers are grateful to Professor Oladele Sunday blessing of Department of Veterinary Pathology, Faculty of Veterinary Medicine, Ahmadu Bello University, Zaria for reading and editing this manuscript.

REFERENCES

- Chibuzo, G.O. and M.A. Sivachelvan, 1989. Dissection Manual of Small Ruminants. Savannah Press, Nigeria, pp: 243-250.
- Fackelman, G.E., J.A. Auer, J. Orsini and S.B. Von, 1983. Surgical treatment of severe flexural deformity of the distal inter phalangeal joint in young horses. J. Am. Vet. Med. Assoc., 187: 1351-1353.
- Horney, F.D. and H.E. Amstutz, 1980. Musculoskeletal System. American Veterinary Publications, Santa Barbara, pp: 863-885.
- Leipold, W.L., T. Hinaga and S.M. Dennis, 1993. Congenital defects of the bovine Musculoskeletal System and joints. Vet. Clin. North Am. Food Anim. Pract. J., 9: 93-104.
- McIlwraith, C.W. and J.F. Fessler, 1978. Evaluation of inferior check ligament desmotomy for treatment of acquired flexor tendon contracture in the horse. J. Am. Vet. Med. Assoc., 172: 293-298.
- Riegall, R.J. and E.H. Susan, 2000. Illustrate Atlas of Clinical Equine Anatomy and Common Disorders of the Horse. Vol. 2, Equistar Publications Limited, Marysville, Ohio.
- Sonnichsen, H.V., 1982. Subcarpal check ligament desmotomy for the treatment of contracted deep flexor tendon in foals. Equine Vet. J., 14: 256-257.
- Von-Veet, J.F. and B.A. Valentine, 2007. Pathology of Domestic Animals. 5th Edn., Elsevier, Edinburg, Texas, pp: 204-207.
- Wagner, P.C., S.M. Reed and G.A. Hegreberg, 1982. Contracted tendons (flexural deformities) in the young horse. Compendium Continuous Educ. Pract. Vet. J., 4: 5101-5108.