

## A Study of Fitting an Alpine Winter Pasture Evapotranspiration to a Model Based on the Penman-Monteith Equation

<sup>1</sup>Jie Li, <sup>1,2</sup>Bin Wang, <sup>1</sup>Xiaoqin Liu, <sup>1</sup>Lulu Feng, <sup>1</sup>Rongrong Ma, <sup>1</sup>Ben Niu,  
<sup>3</sup>Liang Zhao, <sup>3</sup>Yingnian Li, <sup>4</sup>Yanhong Tang, <sup>5</sup>Mingyuan Du and <sup>1</sup>Song Gu

<sup>1</sup>College of Life Science, Nankai University, Tianjin, China

<sup>2</sup>College of Life Science and Agriculture and Forestry, Qiqihar University, Qiqihar, China

<sup>3</sup>Chinese Academy of Sciences, Northwest Plateau Institute of Biology, Xining, China

<sup>4</sup>National Institute for Environmental Studies, Tsukuba, Japan

<sup>5</sup>National Institute for Agro-Environmental Sciences, Tsukuba, Japan

---

**Abstract:** To accurately estimate the magnitude and seasonal dynamics of Evapotranspiration (ET) over an important alpine winter pasture in the Northeastern Qinghai-Tibetan Plateau, researchers employed the Food and Agriculture Organization (FAO) Penman Monteith (P-M) Model. The model was also used to investigate changes in the crop coefficient ( $k_c$ ) which was calculated as the ratio of the measured actual ET ( $ET_a$  from the Eddy Covariance (EC) System) to the reference ET ( $ET_0$  from the P-M Model). The results indicated a reference ET of 900 mm year<sup>-1</sup> from the alpine meadow pasture which was significantly higher than the actual ET (426 mm year<sup>-1</sup>). In addition, the seasonal dynamics of these two ET values differed. The reference ET peaked from April to July while the actual ET was primarily in growing season. The value of  $k_c$  exhibited significant seasonal variations within the range 0.3-1.0 with a mean  $k_c$  of 0.55 during the growing season. The correlation analysis of the relationship between daily  $k_c$  and its primary environmental factors indicated that the Vapor Pressure Deficit (VPD), net Radiation ( $R_n$ ) and air Temperature ( $T_a$ ) were the major influencing factors of the daily  $k_c$ . The daily  $k_c$  showed a linear increase with  $R_n$  and  $T_a$  and a linear decrease with the VPD. With respect to the biotic factors, the biomass exhibited a significant positive correlation with  $k_c$ . Thus, a daily  $k_c$  Model is developed as a function of the VPD,  $R_n$ ,  $T_a$  and biomass. This ET Model was validated using 2005 data and showed a satisfactory consistency between the simulated and measured ET.

**Key words:** Evapotranspiration, reference evapotranspiration, crop coefficient,  $k_c$ ,  $T_a$

---

### INTRODUCTION

The hydrologic balance of terrestrial ecosystems are important determinants of ecosystem structure, function and productivity (Burman and Pochop, 1994). Evapotranspiration (ET) which is the second-largest water flux in the terrestrial hydrologic cycle, plays an important role in the maintenance of the water and energy balance of the ground surface (Oki and Kanae, 2006).

In addition, ET processes are closely related to vegetation conditions, the ecophysiological processes of plants, soil environments and micrometeorological characteristics. Thus, ET is a pivotal water exchange process in the Soil Plant Atmosphere Continuum (SPAC). Therefore, accurate ET estimation is not only significant to the regulation and management of hydrologic cycles in ecosystems but also to scientific decisions regarding local ecological construction and production activities in agriculture and animal husbandry (Tyagi *et al.*, 2000;

Gong *et al.*, 2006; Torriani *et al.*, 2007). Domestic and international researchers have investigated land surface ET for >200 years and have developed a number of fitting models and observational methods for ecosystem evapotranspiration (Priestley and Taylor, 1972; Monteith and Unsworth, 1990; Frank, 2003; Li *et al.*, 2005). Currently, the Eddy Covariance (EC) System is the most extensively used and sophisticated micrometeorological approach. Ecosystem ET may be continuously monitored with the EC System without damage to the soil or vegetation. The Penman equation which utilizes conventional meteorological data is the most influential fitting equation for predicting ET. This equation was proposed in 1948 when Penman proposed an formula for calculating ET from a water surface which considers radiant energy, air saturation deficit, wind speed and other ET-influencing factors. Based on prior research, Monteith modified the Penman equation and proposed the Penman-Monteith (P-M) Model which is a canopy ET

model that has been extensively applied (Monteith and Unsworth, 1990). In 1990s, the Food and Agriculture Organization (FAO) of the United Nations amended the P-M Model for the estimation of actual ET from farmland and grassland and recommended this modified model as a standard method for calculating ET (Allen *et al.*, 1998). In the P-M Model recommended by the FAO (the FAO-P-M Model), a crop coefficient ( $k_c$ ) has been introduced to correct reference ET values and to accurately estimate actual ET values in specific ecosystems. Studies have reported that  $k_c$  is related to various biotic factors such as crop type and growth stage (Williams and Ayars, 2005). For homogeneous vegetation which consists of plants in a particular growth stage,  $k_c$  may be regarded as a constant. For example in the FAO-56 Method,  $k_c$  values are assigned based on a pasture's growth stage thus  $k_c$  values for the initial, mid-season and late-season growth stages are 0.4, 1.05 and 0.85, respectively (Allen *et al.*, 1998). Recent studies have demonstrated that  $k_c$  exhibits specific variations and is affected by radiant energy, moisture levels and other environmental factors (Lockwood, 1999; Yang and Zhou, 2011). The  $k_c$  value is the key to using the FAO-P-M Model to accurately estimate the actual ET of an ecosystem. Thus, the study of the patterns and characteristics of changes in  $k_c$  in natural pastureland can improve the accuracy and simplicity of the calculation of the ecological water demand of pasturelands and provide a theoretical foundation for the grazing production of these lands.

The Qinghai-Tibetan plateau which is referred to as "Asia's water tower", plays a pivotal role in the global hydrologic cycle and the global water balance. Alpine meadow which is one of the most extensively distributed types of vegetation in the Qinghai-Tibetan plateau, covers an area of approximately 1,200,000 km<sup>2</sup> (Zheng *et al.*, 2000). The alpine pasture in this study is an alpine meadow in the northeast of the Plateau which is also an unique natural landscape and one of the most important grassland resource for grazing (Cui and Graf, 2009). Few studies have explored the ET characteristics of meadow ecosystems (Li *et al.*, 2013) as a result, studies on applicable models for accurately and conveniently evaluating ET in the alpine pasture meadow ecosystem also remain scarce.

This study aimed to achieve the following objectives: use the FAO-P-M Model and Eddy Covariance System to explore the dynamics of the actual ET and reference ET changes in the alpine meadow ecosystem of the Qinghai-Tibetan Plateau and derive a suitable  $k_c$ -based Model of the alpine meadow ecosystem by determining how  $k_c$  values vary with changes in meadow climate and

vegetation and by establishing the relationships between  $k_c$  and the factors (including both the environmental and biotic factors).

## MATERIALS AND METHODS

**Study site:** The study area is located in the Northeastern of the Qinghai-Tibetan plateau (37°36'N, 101°18'E; 3250 m). The climate in this area is characterized by long, cold and dry winters and short, cool and moist Summers. The annual mean temperature in the study area is -1.7°C; January is the coldest month (-15°C) and July is the warmest month (10°C). The average annual duration of sunshine is 2462.7 h and the total radiation ( $R_s$ ) received per year ranges from 6000-7000 MJ m<sup>-2</sup>. The alpine meadow features high vegetation cover composed of short and densely distributed plants. The vegetation primarily consists of meadow species such as *Kobresia humilis*, *K. pygmaea*, *K. Tibetica* and *Stipa aliena*. The meadow land which is primarily used for Winter pasture (January to May and September to December) serves as an important grazing resource in the plateau region. The aboveground biomass of the alpine meadow increases in early May and peaks in August with a multi-year average biomass of approximately 350 g m<sup>-2</sup>. The study area receives a yearly precipitation of approximately 600 mm. This precipitation is primarily concentrated between May and September and changes in the Soil Water Content (SWC) are subject to precipitation-related effects.

### Measurements

**Flux measurements:** EC observation system was placed in the center of the study area. This area features flat terrain which enables unobscured observation and provides a sufficiently large "fetch" to satisfy the required physical conditions for eddy and meteorological observations. ET was measured using a H<sub>2</sub>O/CO<sub>2</sub> infrared gas analyzer (Li-7500, Li-Cor, USA) and a three-dimensional sonic anemometer (CSAT3, CSI, USA) with a sensor placed 2.2 m above the ground. A data-logger (CR5000, Campbell Inc.) was employed to continuously record water vapor flux data and output average values at 15 min intervals. The actual ET (ET<sub>a</sub>) was measured by the EC System in this study.

**Meteorological measurements:** Environmental variables were continuously measured at this site. Air temperature (T<sub>a</sub>), humidity and actual vapor pressures were measured at 110 and 220 cm (HMP45C, CSI); soil temperature was measured at 0, 5, 10, 20, 30 and 50 cm depths (thermocouple); solar Radiation (R<sub>s</sub>) and net Radiation (R<sub>n</sub>) were measured at 150 cm (CNR-1, Kipp and Zonen,

Netherlands); a horizontal wind-speed sensor (014A and 034A-L, CSI) was attached at 110 and 220 cm to measure horizontal wind speeds; precipitation was measured at 70 cm (TE525MM, CSI); SWC was measured at 5, 20 and 50 cm depths (TDR soil moisture sensor; CS615, CSI) and the soil heat flux was set at a depth of 2 cm (HFT-3, CSI). The signals were sampled at 10 Hz and 15 min mean data were logged by the data-logger (CR23X, CSI).

**Observations of vegetation data:** Aboveground biomass in the study area was measured semimonthly during the growing season. A harvesting method was adopted to obtain these measurements. In particular, five 0.5×0.5 m quadrants were randomly selected; within each quadrant, plants were harvested by cutting the plants at ground level and loading the plants into sampling bags. Each sample was numbered, rapidly transported to a laboratory and dried at 65°C until a constant weight was obtained for each sample. These weights were subsequently converted into g/m<sup>2</sup>.

**The FAO-P-M Model**

**The calculation of reference ET:** The reference crop comprises a well-managed short grass of uniform height (8-15 cm) that completely covers the ground and grows lushly across open ground without experiencing water stress. The reference crop ET refers to the ET of the reference crop under these conditions (Allen *et al.*, 1998). In this study, the standardized FAO-P-M Model was used to calculate the actual ET in the lpine meadow ecosystem. This model includes two components reference ET and k<sub>c</sub> as indicated in the following Eq. 1:

$$ET_{P\_M} = k_c ET_0 \tag{1}$$

Where:

ET<sub>P\_M</sub> = The actual ET of an ecosystem in the FAO-P-M Model

k<sub>c</sub> = The crop coefficient

ET<sub>0</sub> = The reference ET

The reference surface consists of well-watered grass with a height of 12 cm and a fixed surface resistance of 70 sec m<sup>-1</sup>. The applicable equation is expressed as:

$$ET_0 = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T+273} u(e_s - e_a)}{\Delta + \gamma(1+0.34u)} \tag{2}$$

Where:

Δ = The slope of the saturation vapor pressure curve at the examined temperature

R<sub>n</sub> = The net radiation

G = The soil heat flux

γ = The psychrometric constant

T = The mean daily temperature at a height of 2 m

u = The wind speed at a height of 2 m

e<sub>s</sub> and e<sub>a</sub> = The saturation vapor pressure and actual vapor pressure, respectively

Δ and γ = Calculated as:

$$\Delta = \frac{4098 \times \{0.6108 \exp[17.27T/(T+237.3)]\}}{(T+237.3)^2} \tag{3}$$

$$\gamma = \frac{c_p \times P}{0.622\lambda} \tag{4}$$

$$\lambda = 2.501 - (2.361 \times 10^{-3}) \times T \tag{5}$$

Where:

C<sub>p</sub> = The specific heat at constant pressure which has a fixed value of 1.013×10<sup>-3</sup> MJ/kg°C

P = The atmospheric pressure

0.662 = The ratio of the molecular weight of water vapor to the molecular weight of dry air

λ = The latent heat of vaporization which represents the energy per unit volume of water required to convert water into steam under ambient temperature and pressure conditions

The study meadow area is located on a plateau zone at an average elevation of 3250 m. Using Eq. 6, the atmospheric pressure (P) can be calculated from this average elevation (H = 3250 m):

$$P = 1013 - 0.1093H \tag{6}$$

**The crop coefficient (k<sub>c</sub>):** The crop coefficient k<sub>c</sub> was obtained from the ratio of the reference ET in 2004 to the ET obtained from EC System observations in 2004 and fitted with environmental factors to obtain an empirical equation which was eventually validated by 2005 data. In the FAO-56 approach, the growth of pasture plants was divided into different growth stages (Allen *et al.*, 1998). The study area of the current investigation features an alpine climate in this area, plant growth typically begins in late April and the growing season extends from May to September.

**Evaluation methods:** In this study, the slope, linear correlation coefficient (R<sup>2</sup>), Relative Root-Mean-Squared Error (RRMSE), Index of Agreement (IA) and Coefficient of Determination (CD) were utilized to examine the extent of the differences between the observed and calculated values of ET and to statistically analyze the accuracy of the adopted fitting approach from multiple perspectives. The slope and R<sup>2</sup> were calculated by Origin 8.0. RRMSE, IA and CD values were computed as (Alexandris and Kerkides, 2003; Yang and Zhou, 2011).

$$RRMSE = \sqrt{\frac{\sum_{i=1}^n (O_i - E_i)^2}{n}} \cdot \frac{1}{O^2} \quad (7)$$

$$IA = 1 - \frac{\sum_{i=1}^n (E_i - O_i)^2}{\sum_{i=1}^n (|E_i - \bar{O}| + |O_i - \bar{O}|)^2} \quad (8)$$

$$CD = \frac{\sum_{i=1}^n (O_i - \bar{O})^2}{\sum_{i=1}^n (E_i - O)^2} \quad (9)$$

Where:

$O_i$  = *i*th observed value

$E_i$  = *i*th simulated or estimated value

*n* = The number of samples

$\bar{O}$  = The mean observed value

## RESULTS AND DISCUSSION

### Seasonal variations in reference ET ( $ET_0$ ) and actual ET:

As indicated in Fig. 1, the reference ET derived from the P-M Model was significantly higher than the actual ET measured by the EC System. The reference ET ( $ET_0$ ) refers to the reference capacity of moisture diffusion from the ecosystem to the atmosphere under specific environmental conditions thus  $ET_0$  represents the maximum ET. The annual reference ET of the examined alpine meadow was 900.2 mm which was significantly higher than the actual ET of 425.8 mm. These parameters also exhibited significant seasonal variations.  $ET_0$  began to significantly increase in March and attained peak levels from April to July however, it began to decline in August. The total  $ET_0$  during the growing season (from May to September) was 487.1 mm which represented 54% of the annual cumulative  $ET_0$ . Conversely, seasonal variations in




Fig. 1: Annual variations in reference ET ( $ET_0$ ) and actual ET ( $ET_a$ )

actual ET ( $ET_a$ ) in the alpine meadow lagged behind variations in reference ET. The  $ET_a$  was concentrated in the growing season between May and September the  $ET_a$  was 337.1 mm which represented 79% of the annual cumulative  $ET_a$ .

According to its definition, the reference ET ( $ET_0$ ) was only affected by climatic parameters (Allen *et al.*, 1998). As shown in Eq. 2, the net radiation ( $R_n$ ), Vapor Pressure Deficit (VPD,  $e_s - e_a$ ), air temperature ( $T_a$ ) and wind speed (*u*) determined the reference ET. The annual variation of these environmental factors were shown in Fig. 2.

As the maximum of  $R_n$ , VPD and *u* occurred during April to June (Fig. 2),  $ET_0$  reached its maximum value at the similar period (March to June). However, with the canopy development in growing season, the effect of biotic factors will exert increasing influence on the actual ET. Therefore, with the  $ET_a$  rapidly increase in growing season, the difference between  $ET_0$  and  $ET_a$  became smaller (Fig. 1).

Moreover, the reference ET in growing season (487.1 mm) was lower than the value of precipitation (579.7 mm). The ratio of  $ET_0$  to P was 84% during growing season. This result was similar to a nonirrigated pasture (Sumner and Jacobs, 2005) however, it was much lower than some grassland ecosystems (Wever *et al.*, 2002; Li *et al.*, 2005). In these grassland, the reference ET was significantly higher than the precipitation in growing season.

Comparing to these grassland (Wever *et al.*, 2002; Li *et al.*, 2005), the lower  $ET_0$  may be caused by the unique climate in this plateau. The net radiation ( $R_n$ ) in the Qinghai-Tibetan Plateau was much lower than that for lowland grasslands (Hammerle *et al.*, 2008), despite the high incident solar radiation ( $R_s$ ), due to the fact that the net long-wave radiation in alpine region is much higher than that for lowland regions (Zhang *et al.*, 2010). In this alpine pasture, the  $R_n$  was much lower than the  $R_s$  and the average  $R_n$  and  $R_s$  values were 7.7 and 16.6 mol/( $m^2$  day), respectively (Fig. 2a). The low energy available for water evaporation may limit the reference ET in some extents. Moreover, VPD and  $T_a$  were meteorological variables for controlling on water vapour exchange between the atmosphere and vegetation (Baldocchi and Meyers, 1998). Gu *et al.* (2008) indicated that low VPD and  $T_a$  due to the frequent precipitation and the altitude were characteristics of the climate in the alpine meadow in Northeast Qinghai-Tibetan plateau. Figure 2b showed the variation of VPD and  $T_a$  in this study site. The maximum value of daily mean VPD and  $T_a$  were only 1.5 kPa and 14.5°C which were greatly lower than many other grassland with




Fig. 2: Temporal variations of a) net radiation ( $R_n$ ) and solar radiation ( $R_s$ ); b) daily mean air temperature ( $T_a$ ) and Vapor Pressure Deficit (VPD); c) daily Precipitation (P) and soil water content at 5 cm depth (SWC) and d) wind speed (u) in the alpine meadow during 2004

the maximum VPD ranged from about 2-5 kPa and with the maximum  $T_a$  ranged from about 20-30°C (Kellner, 2001; Hao *et al.*, 2007; Dong *et al.*, 2011). The low VPD and  $T_a$  might imply the weak driving power and are considered as the factors to restrict the reference ET in growing season.

**The seasonal variation in  $k_c$ :** Due to the difference between reference ET and actual ET, the correct




Fig. 3: Seasonal variation in the  $k_c$

determination of  $k_c$  which is the ratio of daily ET to  $ET_0$  is important for the accurate estimation of the actual ET. In this study,  $k_c$  exhibited a gradual rise from mid-April to mid-June and remained at a high level in July and August (Fig. 3). The value of  $k_c$  rapidly declined in September and was  $<0.2$  by the end of October. During the growing season which extended from May-September,  $k_c$  exhibited a mean value of 0.55 and fluctuated within the range 0.30-0.92. This range is consistent with the range of  $k_c$  values reported in FAO-56 (0.30-1.05) (Allen *et al.*, 1998) but is significantly higher than the range of  $k_c$  values observed for a typical steppe (0.32-0.68) (Miao *et al.*, 2009) and temperate desert steppe (0.02-0.50) (Yang and Zhou, 2011) of Inner Mongolia. These results indicated that the  $k_c$  for the alpine pasture features more adequate moisture and better plant growth conditions than arid and semi-arid steppe ecosystems.

**Factors that impact  $k_c$**

**The effect of environmental factors on  $k_c$ :** Previous studies have focused on  $k_c$  values related to local climate conditions (Lockwood, 1999; Yang and Zhou, 2011). In this meadow, the Pearson product-moment correlation coefficients for the relationships between  $k_c$  and its main environmental factors were computed on a daily scale (Table 1). This computation showed that the VPD and  $R_n$  were critical for controlling  $k_c$ .  $T_a$  also showed a significant relation to  $k_c$ , whereas u and SWC at a depth of 5 cm were not significant at the 95% confidence level.

The responses of  $k_c$  to changes in VPD,  $R_n$  and  $T_a$  are illustrated in Fig. 4. To reduce or offset the errors associated with the  $k_c$  values, VPD,  $R_n$  and  $T_a$  were grouped into bins with the following criterion: 0.2 kPa for VPD, 1 MJ/m<sup>2</sup> d for  $R_n$  and 2°C for  $T_a$ .  $k_c$  increased linearly in response to an increase in  $R_n$  and  $T_a$  and in response to a decrease in VPD (Fig. 4). Zhou and Zhou (2009) also obtained a similar result from a study of a reed marsh.

Table 1: Pearson product-moment correlation coefficients for the relationships between daily  $k_c$  and daily average values for environmental variables: SWC at a depth of 5 cm, atmospheric VPD, wind speed ( $u$ ),  $R_n$  and  $T_a$  over the alpine meadow on a daily basis for days without rain during the 2004 growing season ( $n = 106$ )

	$k_c$	SWC	VPD	$u$	$R_n$	$T_a$
$k_c$	1.000					
SWC	-0.168	1.000				
VPD	-0.509**	-0.324*	1.000			
$u$	-0.175	0.339*	-0.114	1.000		
$R_n$	0.442**	-0.440**	0.164	-0.309*	1.000	
$T_a$	0.402*	-0.590**	0.400*	-0.518**	0.429**	1.000

demonstrated that the Soil Water Content (SWC) was the most important factor for determining  $k_c$ . In this study, the root-layer (0-10 cm) SWC fluctuated within the range 0.2-0.6  $m^3/m^3$  (Fig. 1c) which was much better than the SWC at desert steppe in inner Mongolia (0.05-0.15  $m^3/m^3$ ). The higher soil moist in this alpine pasture might weaken the SWC impact on  $k_c$ .

Daily  $k_c$  for the alpine pasture can be expressed by an empirical equation from the correlation and regression analyses of the relationships between  $k_c$  and its statistically significant environmental variables:

$$k_c = 0.597 - 0.801VPD + 0.026R_n + 0.040T_a \quad (10)$$

In this case, based on daily meteorological data and reference  $ET_0$  data during the growing season, the daily actual ET can be calculated by:

$$ET = (0.597 - 0.801VPD + 0.026R_n + 0.040T_a) \times ET_0 \quad (11)$$

**The effects of biotic factors on  $k_c$ :** In addition to environmental factors, the plant species composition, growth conditions and other biotic factors can significantly affect  $k_c$  (Tyagi *et al.*, 2000; Williams *et al.*, 2003; De Medeiros *et al.*, 2005). Biomass directly reflects the growth status of a biological community. In this study, the sampled plants began to grow at the end of April which caused an increase in the biomass that primarily began in early May. The biomass rapidly increased during June and August and attained a peak in late August (with a maximum value of 353.1  $g/m^2$  in the study year). After August, the biomass rapidly declined as plants gradually died (Fig. 5a).

The linear regression analysis of the relationship between  $k_c$  and the biomass demonstrated a significant positive correlation between  $k_c$  and the biomass, however the significance of this correlation changed during different growth stages. As indicated in Fig. 5b, the biomass increased and  $k_c$  rapidly increased from 0.45-0.93 from May to middle August. After biomass attained its peak,  $k_c$  began to decline and decreased to 0.39 in late October. While no significant decrease in  $k_c$  was observed from September to October because  $k_c$  only decreased from 0.54-0.39 during this period. The following linear equations describe how  $k_c$  increases with biomass. May-Middle August:

$$k_c = 0.0019 \times \text{biomass} + 0.4521 \quad (12)$$

Middle August-October:

$$k_c = 0.0008 \times \text{biomass} + 0.2514 \quad (13)$$




Fig. 4: Linear response of  $k_c$  to a) VPD; b)  $R_n$  and c)  $T_a$ . The daily  $k_c$  data were averaged with a bin width of 0.2 kPa for VPD, 1  $MJ/(m^2 \text{ day})$  for  $R_n$  and  $2^\circ C$  for  $T_a$ , respectively. Bars indicate  $\pm SD$

However, the result was different from a study in a temperate desert steppe (Yang and Zhou, 2011) which




Fig. 5: a) The seasonal variation of biomass in this alpine meadow pasture in 2004 and b) The response of  $k_c$  to biomass. The  $k_c$  was averaged with a bin width of 50  $g/m^2$  for biomass. Bars indicate  $\pm$ SD

Considering both biotic and environmental factors, daily actual ET can be calculated as follows. May-Middle August:

$$ET = (0.596 - 0.801VPD + 0.026R_n + 0.040T_a) (0.0019 \text{ biomass} + 0.4521) \times ET_0 \quad (14)$$

Middle August-October:

$$ET = (0.596 - 0.801VPD + 0.026R_n + 0.040T_a) (0.0008 \text{ biomass} + 0.2514) \times ET_0 \quad (15)$$

**Test for modeling evapotranspiration:** Equation 15 was validated using the EC data collected during the 2005 growing season. The model performed well during the growing season. The simulated daily ET ( $ET_{mod}$ ) followed the seasonal trend of the EC measured ET ( $ET_a$ ) (Fig. 6). The slope (= 0.95) of the regression line between the measured and simulated ET passes through the origin close to 1 with  $R^2 = 0.76$ , a RRMSE of 0.23  $mm \text{ day}^{-1}$  and an IA of 0.92.

In this study, the abiotic and biotic factors enabled a better estimation of evapotranspiration. RRMSE is a




Fig. 6: Seasonal variation in the measured and simulated ET ( $ET_a$  and  $ET_{mod}$ ) for days without rain in 2005 using Eq. 15

Table 2: The validation statistics of the model simulated by the meteorological factors versus the model simulated by the meteorological factors and biotic factors

Variables	Slope	R <sup>2</sup>	RRMSE	IA	CD
$ET_{mod1}$	0.89	0.68	0.28	0.87	0.76
$ET_{mod2}$	0.95	0.76	0.23	0.93	0.87

$ET_{mod1}$  is simulated by the meteorological factors (VPD,  $R_n$  and  $T_a$ );  $ET_{mod2}$  is simulated by these meteorological factors and biomass

measure of the relative magnitude of residuals; smaller values of RRMSE indicate better model calculation results. The IA can be used to evaluate the correlation between the observed values and the simulated values; the closer IA is to 1, the more closely the fitted values match the observed values. The CD can be used to measure the dispersion of the simulated values from the mean observed values; a CD greater than 0.8 indicates satisfactory simulation results (Cai *et al.*, 2007). In Table 2, the slope,  $R^2$ , IA and CD of  $ET_{mod1}$  are greater and the RRMSE is smaller than those of  $ET_{mod2}$ . These results indicated that relative to the fitting method that accounts for meteorological variables, the fitting method that comprehensively considers meteorological factors and vegetation conditions can be used to derive simulated ET results that are closer to the actual observed ET results (Table 2).

## CONCLUSION

The FAO-Penman-Monteith Model was used to simulate ET over the alpine meadow on the Qinghai-Tibetan plateau. The reference ET was estimated by the FAO-P-M Model and showed difference from the actual ET measured by the EC System. The crop coefficients ( $k_c$ ) was used to estimate  $ET_a$ . The average value of  $k_c$  during growing season was 0.55 (ranging from

0.30-0.92). VPD,  $R_n$  and  $T_a$  explained the majority of the daily variation in  $k_c$  which showed a linear increase with an increase in  $R_n$  and  $T_a$  and a linear decrease with an increase in VPD. Researchers also considered the impact of biomass on  $k_c$  which showed a significant positive effect on  $k_c$ . A daily empirical  $k_c$  model driven by VPD,  $R_n$ ,  $T_a$  and biomass was developed to estimate daily actual ET using the FAO-56  $k_c$  approach and the P-M Model for reference ET. This ET model was validated against 2005 growing season data for this meadow and demonstrated a suitable and consistent performance between the simulated and measured ET.

#### ACKNOWLEDGEMENTS

This study was supported by the National Natural Science Foundation of China (31070433) and the Japan-China Research Cooperative Program (2010DFA31290).

#### REFERENCES

- Alexandris, S. and P. Kerkides, 2003. New empirical formula for hourly estimations of reference evapotranspiration. *Agric. Water Manage.*, 60: 157-180.
- Allen, R.G., L.S. Pereira, D. Raes and M. Smith, 1998. FAO irrigation and drainage Paper, No. 56, crop evapotranspiration, guidelines for computing crop water requirements. FAO, Rome.
- Baldocchi, D. and T. Meyers, 1998. On using eco-physiological, micrometeorological and biogeochemical theory to evaluate carbon dioxide, water vapor and trace gas fluxes over vegetation: A perspective. *Agric. Forest Meteorol.*, 90: 1-25.
- Burman, R.D. and L.O. Pochop, 1994. Evaporation, Evapotranspiration and Climatic Data. Vol. 22, Elsevier Science Ltd., USA., ISBN: 9780444819406, pp: 1-5.
- Cai, J., Y. Liu, T. Lei and L.S. Pereira, 2007. Estimating reference evapotranspiration with the FAO Penman-Monteith equation using daily weather forecast messages. *Agric. Forest Meteorol.*, 145: 22-35.
- Cui, X. and H.F. Graf, 2009. Recent land cover changes on the Tibetan Plateau: A review. *Climat. Change*, 94: 47-61.
- De Medeiros, G.A., F.B. Arruda and E. Sakai, 2005. Crop coefficient for irrigated beans derived using three reference evaporation methods. *Agric. Forest Meteorol.*, 135: 135-143.
- Dong, G., J. Guo, J. Chen, G. Sun, S. Gao, L. Hu and Y. Wang, 2011. Effects of spring drought on carbon sequestration, evapotranspiration and water use efficiency in the Songnen meadow steppe in northeast China. *Ecophysiology*, 4: 211-224.
- Frank, A.B., 2003. Evapotranspiration from northern semiarid grasslands. *Agron. J.*, 95: 1504-1509.
- Gong, L., C.Y. Xu, D. Chen, S. Halldin and Y.D. Chen, 2006. Sensitivity of the Penman-Monteith reference evapotranspiration to key climatic variables in the Changjiang (Yangtze River) basin. *J. Hydrol.*, 329: 620-629.
- Gu, S., Y. Tang, X. Cui, M. Du and L. Zhao *et al.*, 2008. Characterizing evapotranspiration over a meadow ecosystem on the Qinghai-Tibetan plateau. *J. Geophys. Res.: Atmos.*, Vol. 113. 10.1029/2007JD009173.
- Hammerle, A., A. Haslwanter, U. Tappeiner, A. Cernusca and G. Wohlfahrt, 2008. Leaf area controls on energy partitioning of a temperate mountain grassland. *Biogeosciences*, 5: 421-431.
- Hao, Y., Y. Wang, X. Huang, X. Cui and X. Zhou *et al.*, 2007. Seasonal and interannual variation in water vapor and energy exchange over a typical steppe in inner Mongolia, China. *Agric. Forest Meteorol.*, 146: 57-69.
- Kellner, E., 2001. Surface energy fluxes and control of evapotranspiration from a Swedish *Sphagnum* mire. *Agric. Forest Meteorol.*, 27: 101-123.
- Li, J., S. Jiang, B. Wang, W.W. Jiang, Y.H. Tang, M.Y. Du and S. Gu, 2013. Evapotranspiration and its energy exchange in alpine meadow ecosystem on the Qinghai-Tibetan plateau. *J. Integr. Agric.*, 12: 1396-1401.
- Li, S.G., C.T. Lai, G. Lee, S. Shimoda, T. Yokoyama, A. Higuchi and T. Oikawa, 2005. Evapotranspiration from a wet temperate grassland and its sensitivity to microenvironmental variables. *Hydrol. Process.*, 19: 517-532.
- Lockwood, J.G., 1999. Is potential evapotranspiration and its relationship with actual evapotranspiration sensitive to elevated atmospheric CO<sub>2</sub> levels? *Clim. Change*, 41: 193-212.
- Miao, H., S. Chen, J. Chen, W. Zhang and P. Zhang *et al.*, 2009. Cultivation and grazing altered evapotranspiration and dynamics in inner Mongolia steppes. *Agric. Forest Meteorol.*, 149: 1810-1819.
- Monteith, J.L. and M.H. Unsworth, 1990. Principles of Environmental Physics. 2nd Edn., Edward Arnold, New York.
- Oku, T. and S. Kanae, 2006. Global hydrological cycles and world water resources. *Science*, 313: 1068-1072.
- Priestley, C.H.B. and R.J. Taylor, 1972. On the assessment of surface heat flux and evaporation using large-scale parameters. *Mon. Weather Rev.*, 100: 81-92.
- Sumner, D.M. and J.M. Jacobs, 2005. Utility of penman-monteith, priestley-taylor, reference evapotranspiration and pan evaporation methods to estimate pasture evapotranspiration. *J. Hydrol.*, 308: 81-104.



- Torriani, D., P.L. Calanca, S. Schmid, M. Beniston and J. Fuhrer, 2007. Potential effects of changes in mean climate and climate variability on the yield of winter and spring crops in Switzerland. *Clim. Res.*, 34: 59-69.
- Tyagi, N.K., D.K. Sharma and S.K. Luthra, 2000. Determination of evapotranspiration and crop coefficients of rice and sunflower with lysimeter. *Agric. Water Manage.*, 45: 41-54.
- Wever, L.A., L.B. Flanagan and P.J. Carlson, 2002. Seasonal and interannual variation in evapotranspiration, energy balance and surface conductance in a northern temperate grassland. *Agric. Forest Meteorol.*, 112: 31-49.
- Williams, L.E. and J.E. Ayars, 2005. Grapevine water use and the crop coefficient are linear functions of the shaded area measured beneath the canopy. *Agric. Forest Meteorol.*, 132: 201-211.
- Williams, L.E., C.J. Phene, D.W. Grimes and T.J. Trout, 2003. Water use of mature Thompson Seedless grapevines in California. *Irrigation Sci.*, 22: 11-18.
- Yang, F. and G. Zhou, 2011. Characteristics and modeling of evapotranspiration over a temperate desert steppe in inner Mongolia, China. *J. Hydrol.*, 396: 139-147.
- Zhang, X., S. Gu, X. Zhao, X. Cui and L. Zhao *et al.*, 2010. Radiation partitioning and its relation to environmental factors above a meadow ecosystem on the Qinghai-Tibetan plateau. *J. Geophys. Res.: Atmos.*, Vol. 115. 10.1029/2009JD012373.
- Zheng, D., Q.S. Zhang and S.H. Wu, 2000. Mountain Geocology and Sustainable Development of the Tibetan Plateau. Springer, New York, ISBN: 9780792366881, Pages: 393.
- Zhou, L. and G. Zhou, 2009. Measurement and modelling of evapotranspiration over a reed (*Phragmites australis*) marsh in Northeast China. *J. Hydrol.*, 372: 41-47.