

In Vitro* Callus Induction from Seed, Leaf and Protocorm Explants of *Paphiopedilum rothschildianum

¹Makdi Masnoddin, ²Rimi Repin and ³Zaleha Abd Aziz

¹Preparatory Centre for Science and Technology, Universiti Malaysia Sabah (UMS),
Jalan, 88400 Kota Kinabalu, Sabah, Malaysia

²Orchid Conservation Centre, Sabah Park, P.O. Box 10626, 88806 Kota Kinabalu, Sabah, Malaysia

³Faculty of Science and Natural Resources, University Malaysia Sabah (UMS),
Jalan, 88400 Kota Kinabalu, Sabah, Malaysia

Abstract: This research was conducted to investigate the effects of different types of explants on the formation of *Paphiopedilum rothschildianum* callus. Seed, Leaf Segments (LS), Seed-Derived Protocorms (SDP) and Secondary Protocorms (SP) were cultured on half-strength (MS) Media Supplemented with 5 mg/L 2, 4-Dichlorophenoacetic acid (2,4-D) and 1 mg/L 1-phenyl-3-(1-23-Thiadiazol-5-yl)-urea (TDZ). Lower percentage of seed 73.5±14.3 and SP 65.0±22.4% formed callus as compared to SDP 87.5±16.9%. However, the mass of callus induced was highest on SP as compared to SDP and seed explants. Leaf explants failed to form callus and died after 40 days of culture. Callus started to form on seed, SDP and SP after 30 days of culture. Further, study on SDP shows that 3 months old SDP cultured on medium with 1 mg/L TDZ and 4 mg/L 2,4-D was the best explant for callus induction with 82.5%±16.9 explant forming callus recorded as compared to 81.3±12.5 and 72.0±11.0 for 1 and 6 months old SDP, respectively. Calli produced from the present study slowly regenerates into Protocorm Like Bodies (PLBs) and develop shoots and leaflets that further grow into plantlets after 120 days of culture on development medium.

Key words: Seed-derived protocorms, secondary protocorms, Protocorm Like Bodies (PLBs), 2, 4-dichlorophenoacetic acid, 1 mg/L 1-phenyl-3-(1-23-Thiadiazol-5-yl)-urea (TDZ), callus

INTRODUCTION

Paphiopedilum rothschildianum is a rare slipper orchid endemic to the area around Mount Kinabalu, Sabah, Malaysia (Cribb, 1998). Being rare, the species is classified as endangered, the Convention on International Trade in Endangered Species (CITES) is discussed in this study. As with other *Paphiopedilum*, the species has a very slow growth rate, consequently propagating the orchid through the natural ways takes several years. Hence, multiplying the orchid through tissue culture techniques is seen as the best alternative. Micropropagation has been applied to propagate many orchid species in large scales, the technique produced high mass of plants in a considerably short time and with a lower cost. Micropropagation through callus culture is now widely used to mass propagate many orchids, these include *Cymbidium ensifolium* var. *misericors* (Chang and Chang, 1998); *Phaelonopsis* (Ishii *et al.*, 1998; Chen *et al.*, 2000; Tokuhara and Mii, 2001); *Pleione formosana* Hayata (Lu, 2004); *Dendrobium fimbriatum*

(Roy and Banerjee, 2003). Only a few research on callus culture of slipper orchids had been reported; Lin *et al.* (2000) and Hong *et al.* (2008) successfully induced callus of *Paphiopedilum* hybrids using seed-derived protocorm and seed respectively as explants while Lee and Lee (2003) established callus culture of *Cypripedium formosanum* using seed-derived protocorm. To date, there is no reports on callus culture for *P. rothschildianum*. Hence, the objective of this research was to evaluate the potential of different explants to produce callus.

MATERIALS AND METHODS

Plant material: *Paphiopedilum rothschildianum* capsules were collected from Poring and Kinabalu National Park Orchid Nursery. The capsules were surface sterilised by immersing in 10% (v/v) chlorox followed by rinsing three time with sterile distilled water. Sterilised capsules were then dissected longitudinally and some of the seeds were used as explants. To obtain protocorms, a portion of the seeds were germinated on half strength by

Murashige and Skooge, medium (MS) and the resulting protocorms were used as explants designated as Seed-Derived Protocorms (SDP). Subsequently, some of the protocorms formed from these seeds were cultured on a multiplication medium (modified RE medium) and new protocorms or secondary protocorms formed on the protocorm explants were designated as Secondary Protocorms (SP). Leaves from in vitro plantlets of *P. rothschildianum* were cut into pieces of 1×1 cm and used as explants designated as Leaf Segments (LS). For SDP explants, 1, 3 and 6 months old explants were also tested on MS Medium Supplemented with 1 mg/L TDZ and 4 or 5 mg/L 2, 4-D.

Callus induction: To induce callus, medium based on Hong *et al.* (2008) was employed. The medium was comprised of half strength MS basal medium supplemented with full strength MS Vitamin, 2 g/L peptone, 170 mg/L NaH_2PO_4 , 20 g/L sucrose, 5 mg/L 2, 4-D with 1 mg/L TDZ and 2.2 g/L gelrite. The study on the effect of age of SDP explants were conducted using two different combinations of PGRs; 1 mg/L TDZ with 4 and 1 mg/L TDZ with 5 mg/L 2, 4-D. The pH of medium was adjusted to 5.2 and autoclaved at 121°C and 15 psi for 20 min. The medium was dispersed in 9 cm diameter petri dishes and four explants of seed, SDP, SP and LS were cultured on each petri dish. Each treatment was replicated 5 times. All cultures were maintained at $25 \pm 2^\circ\text{C}$ in darkness. Observations were made every 10 days for 150 days and explants were subcultured onto fresh medium at 4 weeks intervals.

Data collection and statistical analysis: All data were analyzed using SPSS (Statistical Package for Social Science) Version 22.0 and subjected to Analysis of Variance (ANOVA). Duncan's multiple range tests were conducted for mean comparisons of data collected using $p < 0.05$.

RESULTS AND DISCUSSION

Combination of 2, 4-D (5 mg/L) and 1 mg/L TDZ did successfully induce and stimulate the growth of callus on seeds, SP, SDP explants. Callus became visible after 30 days of culture and slowly increases in mass through the 150 days period (Plate 1 A-H). All explants produced friable callus with creamy colour. Similar results were reported by Lin *et al.* (2000), Lee and Lee (2003) and Hong *et al.* (2008) with 5, 2, 4-D and 1 mg/L TDZ gave the highest percentages of explants forming callus. Unfortunately, LS failed to produce callus and died after 40 days of culture (Plate 1I-J). Stems, root tips and leaves

Table 1: Formation of callus by different types of *Paphiopedilum rothschildianum* explants on $\frac{1}{2}$ (MS) Medium Supplemented with 1 mg/L TDZ+5 mg/L 2, 4-D in darkness, $25 \pm 2^\circ\text{C}$

Explants	Percentage of explant forming callus at different days (mean%±SD)		
	30	90	150
Seed	7.5±4.3 ^b	NA	73.5±14.3 ^a
SDP	56.7±24.5 ^a	85.4±22.5 ^a	87.5±16.9 ^a
SP	41.7±21.2 ^a	50.0±35.4 ^a	65.0±22.4 ^a
Leaf	0	0	0

^a, ^bData were taken from 5 replicates with the same latters are not significantly different at $p < 0.05$ using Duncan's multiple range test, SD = Standard Deviation

Table 2: The sizes of callus induced on *Paphiopedilum rothschildianum* explants on $\frac{1}{2}$ MS Medium Supplemented with 1 mg/L TDZ+5 mg/L 2, 4-D in darkness, $25 \pm 2^\circ\text{C}$ based on the cultured days

Explants	Callus size at different days		
	30	60	90
Seed	+	++	++
SDP	+	++	++
SP	++	+++	+++

Callus mass were cataloged as large (+++); moderate (++); small (+)

of *Paphiopedilum* hybrids failed to produce callus when cultured on media containing combination of TDZ and either 2, 4-D and PBOA as well as medium containing 2, 4-D in combination of TDZ, BA or 2ip as reported by Lin *et al.* (2000) (Fig. 1).

The highest percentages of explants forming callus after 150 days of culture was SDP (87.5%±16.9) followed by seeds (73.5%±14.3) and SP (65.0%±22.4) (Table 1). These results are in contrast with the report made by Lin *et al.* (2000) which showed seed explants formed better callus than SDP. Although, SDP gave the highest percentage of explant forming callus, it was SP explants that showed higher mass of callus formed on explants (Table 2). Better formation of callus by SP explants might due to the carry over effect as SP explants originated from multiplication medium which contained PGRs.

Makara *et al.* (2010) conclude that PGR like TDZ has a carry-over effect from his study on banana. They found that the proliferation rates of shoots originating from basal cycle medium with various TDZ concentrations were significantly higher than those from 5 mg/L BAP, thus, suggested that TDZ had a high carry over effect, enabling the shoots to continue proliferating on the hormone free medium (Makara *et al.*, 2010).

Further study on the age of SDP explants showed that the percentage of callus formation on 3 months old SDP was significantly higher (70.0%±19.7, 65.0%±22.4) as compared to 1 month old SDP (43.8%±12.5, 40.0%±22.5) for every treatments with 4 mg/L and 5 mg/L 2, 4-D, respectively, after 30 days of culture (Table 3 and 4; Fig. 2). After 90 days of culture, the percentage of explants forming callus on 3 months old SDP increased to




Fig. 1: Callus induction of *Paphiopedilum rothschildianum* on $\frac{1}{2}$ MS medium with 1 mg/L TDZ and 5 mg/L 2, 4-D in darkness, $25\pm 2^\circ\text{C}$; a) SDP after 30 days of culture (bar = 0.18 cm); b) SDP after 60 days of culture (bar = 0.19 cm); c) SDP after 90 days of culture (bar = 0.19 cm); d) SP after 30 days of culture (bar = 0.19 cm); e) SP after 60 days of culture (bar = 0.19 cm); f) SP after 90 days of culture (bar = 0.19 cm); g) Seed after 30 days of culture (bar = 0.05 cm); h) Seeds after 80 days of culture (bar = 0.21 cm); i) LS after 10 days of culture (bar = 0.22 cm) and j) LS after 40 days of culture (bar = 0.21 cm)




Fig. 2: Callus induction of *Paphiopedilum rothschildianum* on $\frac{1}{2}$ MS medium with 1 mg/L TDZ+4 mg/L 2, 4-D in darkness, $25\pm 2^\circ\text{C}$ (bar = 2.5 mm) Callus formed on 1-month-old SDP explants; a) After 30 days of culture, b) Slowly increase in size after 60 days of culture; c) Proliferate further after 90 days of culture; callus formation was faster on 3 months old SDP; d) After 30 days of culture, e) Proliferated well after 60 days of culture; f) The mass of callus further increase after 90 days of culture. Lower mass of callus was observed on 6 months old SDP; g) After 30 days of culture; h) The mass of callus slowly increase after 60 days of culture and i) and the size of callus further increase after 90 days of culture

Table 3: The effect of age of Seed-Derived Protocorm (SDP) on callus induction of *Paphiopedilum rothschildianum* on ½ (MS) Medium Supplemented with 1 mg/L TDZ and different concentration of 2, 4-D in darkness, 25±2°C

Age of explants (months)	2, 4-D (mg/L)	Percentage of explant forming callus at different days (mean±SD)		
		30	60	90
1	4	43.8±12.5 ^{ab}	68.8±12.5 ^{ab}	81.3±12.5 ^a
	5	40.0±22.5 ^b	65.0±22.4 ^{ab}	70.0±32.6 ^{ab}
3	4	70.0±19.7 ^a	82.5±16.9 ^a	82.5±16.9 ^a
	5	65.0±22.4 ^{ab}	80.0±20.9 ^a	80.0±20.9 ^a
6	4	16.0±8.9 ^c	68.0±11.0 ^{ab}	72.0±11.0 ^{ab}
	5	16.0±16.7 ^c	48.0±11.0 ^c	48.0±11.0 ^b

^{a, b}Data were taken from 5 replicates with the same letters are not significantly different at p<0.05 using Duncan's multiple range test, SD = Standard Deviation

Table 4: The sizes of callus induced on *Paphiopedilum rothschildianum* 1 and 3 months old explants on ½ (MS) Medium Supplemented with 1 mg/L TDZ and different concentration of 2, 4-D in darkness, 25±2°C

Age of explants (months)	Concentration of 2, 4-D (mg/L)	Sizes of callus at different days		
		30	60	90
1	4	+	++	+++
	5	+	++	+++
3	4	++	++	+++
	5	++	++	+++
6	4	+	++	+++
	5	+	++	+++

Callus mass were cataloged as large (+++); moderate (++); small (+)

82.5%±16.9 for medium 4 mg/L 2, 4-D and 80.0±20.9 for medium with 5 mg/L 2, 4-D. In addition, after 30 days of culture, the size of callus formed on 3 months old SDP (Table 4, Fig. 2d) was bigger as compared to 1 month old SDP (Fig. 2a) and 6 months old SDP (Fig. 2g). These results indicates that the formation of callus was faster on 3 months old SDP. This is supported by Chen *et al.* (2000) in their study that shows 2 months old protocorm explants of *Phalaenopsis* was better for callus induction than 1 month old protocorm. The maturity of explants known to have critical impact on their growth and development in in vitro culture. The results on Table 3, also shows that 6 months old SDP give the lowest percentage of callus formation on both treatment of 4 mg/L 2, 4-D (72.0±11.0%) and 5 mg/L 2, 4-D (48.0±11.0%). This shows that callus induction also affected by the age of SDP explants. Younger explants consists if more responsive and actively dividing cells. Feng and Chen (2014) suggest that 2 months old seedlings of *Phalaenopsis aphrodite* were more suitable for inducing embryogenesis than 4 months old seedlings. The present study shows that, although, younger explants give better response to callus formation on SDP explants, the maturity of explants also plays an important role in their growth and development. Explant aged 1 months old was young but




Fig. 3: Growth and development of Protocorm Like Bodies (PLBs) regenerated from callus induced on 3 months old SDP (Bar = 1 mm); a) Callus induced was transferred to regeneration medium and slowly regenerated into PLBs; b) PLBs then multiplied and transferred to development medium where it undergoes growth and development after 14 days of culture; c) The formation of shoots and leaflets was observable after 50 days of culture; d) The shoots and leaflets slowly developed into plantlets after 120 days of culture

not matured while 6 months old SDP was too matured. Therefore, the study conclude that 3 months old SDP was the best explant for callus induction because of its young and matured cells.

The results also indicate that lower concentration of 2, 4-D was better for callus induction on all SDP explants with different age (Table 3). The highest percentage of explant forming callus (82.5%±16.9) was on the medium with 1 mg/L TDZ+4 mg/L 2, 4-D. This indicates that SDP explants contain sufficient endogenous auxin for callus induction. The addition of 2, 4-D may limit the capacity of the explant to produce callus. The effect of endogenous auxin in combination with cytokinin were demonstrated by Khosravi *et al.* (2008) on *Dendrobium*. Their study showed that the combination of exogenous cytokinin and endogenous auxin significantly increased the regeneration response of *Dendrobium calli*. However, increasing the concentration of auxin may limit the explant response. This was shown in the study by Chen and Chang (2001) in which the presence of endogenous auxin reduced the rate of direct embryogenesis from leaf explants of *Oncidium* "Grower Ramsey".

Calli originated from 3 months old SDP was transferred to regeneration medium and slowly regenerates into Protocorm Like Bodies (PLBs) formation (Fig. 3a). The PLBs then transferred

to development medium and slowly develop shoots and leaflets that further grow into plantlets after 120 days of culture (Fig. 3b-d).

CONCLUSION

This research proved that callus can be induced on *P. rothschildianum* explants within 90 days and although, SDP explants formed the highest percentage of callus it was SP explant that produced the largest biomass.

RECOMMENDATIONS

Further study on SDP explants showed that 3 months old SDP was the best explant source for callus induction. This is mainly due to the level of maturity of the explant. The calli obtained from the present work was able to regenerate into plantlets after 120 days of culture on the development medium.

ACKNOWLEDGEMENTS

The researchers wish to thank Taman Sabah for supplying *P. rothschildianum* pod and Universiti Malaysia Sabah for funding the research under the Fundamental Research Grant Scheme (GPS0009-SG-1/2009).

REFERENCES

Chang, C. and W.C. Chang, 1998. Plant regeneration from callus culture of *Cymbidium ensifolium*, var. *misericors*. *Plant Cell Rep.*, 17: 251-255.
 Chen, J.T. and W.C. Chang, 2001. Effects of auxins and cytokinins on direct somatic embryogenesis on leaf explants of *Oncidium 'Gower Ramsey'*. *Plant Growth Regul.*, 34: 229-232.
 Chen, Y.C., C. Chang and W.C. Chang, 2000. A reliable protocol for plant regeneration from callus culture of *Phalaenopsis*. *Vitro Cell. Dev. Biol. Plant*, 36: 420-423.

Cribb, P., 1998. The Genus *Paphiopedilum*. 2nd Edn., Natural History Publication (Borneo), Kota Kinabalu, Malaysia, ISBN-13: 9789838120234, Pages: 427.
 Feng, J.H. and J.T. Chen, 2014. A novel in vitro protocol for inducing direct somatic embryogenesis in *Phalaenopsis aphrodite* without taking explants. *Sci. World J.*, 2014: 1-7.
 Hong, P.I., J.T. Chen and W.C. Chang, 2008. Plant regeneration via protocorm-like body formation and shoot multiplication from seed-derived callus of a *maudiae* type slipper orchid. *Acta Physiologiae Plantarum*, 30: 755-759.
 Ishii, Y., T. Takamura, M. Goi and M. Tanaka, 1998. Callus induction and somatic embryogenesis of *Phalaenopsis*. *Plant Cell Rep.*, 17: 446-450.
 Khosravi, A.R., M.A. Kadir, S.B. Kazemin, F.Q. Zaman and A.E.D. Silva, 2008. Establishment of a plant regeneration system from callus of *Dendrobium* cv. *Serdang beauty*. *Afr. J. Biotechnol.*, 7: 4093-4099.
 Lee, Y.I. and N. Lee, 2003. Plant regeneration from protocorm-derived callus of *Cypripedium formosanum*. *In vitro Cell. Dev. Biol. Plant*, 39: 475-479.
 Lin, Y.H., C. Chang and W.C. Chang, 2000. Plant regeneration from callus culture of a *Paphiopedilum* hybrid. *Plant Cell Tissue Organ Cult.*, 62: 21-25.
 Lu, M.C., 2004. High frequency plant regeneration from callus culture of *Pleione* for *Mosana Hayata*. *Plant Cell Tissue Org. Cult.*, 78: 93-96.
 Makara, A.M., P.R. Rubaihayo and J.S. Magambo, 2010. Carry-over effect of Thidiazuron on banana in vitro proliferation at different culture cycles and light incubation conditions. *Afr. J. Biotechnol.*, 21: 3079-3085.
 Roy, J. and N. Banerjee, 2003. Induction of callus and plant regeneration from shoot-tip explants of *Dendrobium fimbriatum* Lindl. var. *oculatum* Hk. f. *Sci. Hortic.*, 97: 333-340.
 Tokuhara, K. and M. Mii, 2001. Induction of embryogenic callus and cell suspension culture from shoot tips excised from flower stalk buds of *Phalaenopsis* (Orchidaceae). *In vitro Cell. Dev. Biol. Plant*, 37: 457-461.