

A Pseudo B-Ideal, Pseudo H-Ideal and a Pseudo Essence of a Pseudo BH-Algebra

Husein Hadi Abbass and Adel Hashem Nouri

Department of Mathematics, Faculty of Computer Science and Mathematics,
 University of Kufa, Kufa, Iraq

Abstract: In this study, we define the notion of a pseudo B-ideal, a pseudo H-ideal and a pseudo essence of a pseudo BH-algebra. Also, we study some properties and relationship between them.

Key words: BH-algebra, ideal of BH-algebra, pseudo BH-algebra, pseudo ideal of a pseudo BH-algebra, pseudo B-ideal, pseudo H-ideal, pseudo essence, pseudo 0-commutative, pseudo G-part, pseudo BCA-part, pseudo closed ideal

INTRODUCTION

Jun *et al.* (1998) introduced the notion of BH-algebra which is a generalization of BCH-algebra and the notion of ideal of a BH-algebra. Kim and Ahn (2011) introduced the notion of essence of BH-algebra. Abbass and Dahham (2012) introduced the notion of completely closed ideal of a BH-algebra. By Abbass and Mahdi (2014) introduced the notion of a closed ideal, p-ideal, q-ideal to a BH-algebra and BCA-part. Jun and Kim (2015) introduced the notion of a pseudo BH-algebra.

MATERIALS AND METHODS

In this study, some basic concepts about a BH-algebra, ideal of BH-algebra, essence BH-algebra, 0-commutative BH-algebra, G-part of BH-algebra, BCA part of BH-algebra pseudo, BH-algebra, pseudo subalgebra of a pseudo BH-algebra and pseudo ideal of a pseudo BH-algebra are given.

Definition 1; Jun *et al.* (1998): A BH-algebra is a nonempty set \mathfrak{X} with constant 0 and a binary operation “*” satisfying the following conditions:

- $x*x = 0, \forall x \in \mathfrak{X}$
- $x*0 = x, \forall x \in \mathfrak{X}$
- $x*y = 0$ and $y*x = 0 \Rightarrow x = y, \forall x, y \in \mathfrak{X}$

Definition 2; Abbass and Dahham (2012): A nonempty subset S of a BH-algebra \mathfrak{X} is called a subalgebra of \mathfrak{X} , if for any $x, y \in S$, we have $x*y \in S$.

Definition 3; Abbass and Dahham (2012): A BH-algebra \mathfrak{X} is said a 0-commutative if : $x*(0*y) = y*(0*x)$. For all $x, y, z \in \mathfrak{X}$.

Definition 4; Abbass and Mhadi (2014): Let \mathfrak{X} be a BH-algebra. Then the set $G(\mathfrak{X}) = \{x \in \mathfrak{X} : 0*x = x\}$ is called G-part.

Definition 5; Abbass and Mahdi (2014): Let \mathfrak{X} be a BH-algebra. Then the set $\mathfrak{X}_+ = \{x \in \mathfrak{X} : 0*x = 0\}$ is called the BCA-part of \mathfrak{X} .

Definition 6; Kim and Ahn (2011): Let, \mathfrak{X} be a BH-algebra. For any subsets G and H of \mathfrak{X} , we define $G*H = \{x*y : x \in G, y \in H\}$.

Theorem 1; Kim and Ahn (2011): Let a subsets A, B and E of a BH-algebra, we have:

- $A \subseteq B \Rightarrow A*E \subseteq B*E$ and $E*A \subseteq E*B$
- $(A \cap B)*E \subseteq (A*E) \cap (B*E)$
- $E*(A \cap B) \subseteq (E*A) \cap (E*B)$
- $(A \cup B)*E = (A*E) \cup (B*E)$
- $E*(A \cup B) = (E*A) \cup (E*B)$

Definition 7; Kim and Ahn (2011): If A is a nonempty subset of a BH-algebra \mathfrak{X} satisfies $A*\mathfrak{X} = A$, then A is called essence of \mathfrak{X} .

Theorem 2; Kim and Ahn (2011): Let \mathfrak{X} be a BH-algebra. Then every a essence of \mathfrak{X} is a subalgebra of \mathfrak{X} .

Theorem 3; Kim and Ahn (2011): Let \mathfrak{X} be a BH-algebra. Then, every essence of \mathfrak{X} contains the zero element 0.

Definition 8; Jun et al. (1998): Let, \mathfrak{X} be a BH-algebra and $I(\neq \emptyset) \subseteq \mathfrak{X}$. Then, I is called an ideal of \mathfrak{X} if it satisfies:

- $0 \in I$
- If $x * y \in I$ and $y \in I \Rightarrow x \in I$, for all $x \in \mathfrak{X}$

Definition 9: An ideal I of a BCH-algebra is called a closed ideal of \mathfrak{X} if for every $x \in I$, we have $0 * x \in I$. We generalize the concept of an ideal to a BH-algebra.

Definition 10: An ideal I of a BH-algebra \mathfrak{X} is called a closed ideal of \mathfrak{X} if: $0 * x \in I$, for all $x \in I$.

Definition 11; Abbass and Dahham (2012): Let \mathfrak{X} be a BH-algebra and I be a subset of \mathfrak{X} . Then I is called a BH-ideal of \mathfrak{X} if it satisfies the following conditions:

- $0 \in I$
- $x * y \in I$ and $y \in I$ imply $x \in I$
- $x \in I$ and $y \in \mathfrak{X}$ imply $x * y \in I$, $I * \mathfrak{X} \subseteq I$

Definition 12; Jun and Kim (2015): A pseudo BH-algebra is a nonempty set \mathfrak{X} with a constant 0 and two binary operations “ $*$ ” and “ $\#$ ” satisfying the following condition:

- $x * x = x \# x = 0$
- $x * 0 = x \# 0 = x$
- $x * y = y \# x = 0 \Rightarrow x = y$, $\forall x, y \in \mathfrak{X}$

Definition 13; Jun and Kim (2015): Let $(\mathfrak{X}, *, \#, 0)$ be a pseudo BH-algebra, then a nonempty subset S of a pseudo BH-algebra \mathfrak{X} is called a pseudo subalgebra of \mathfrak{X} , if for any $x, y \in S$, we have $x * y, x \# y \in S$.

Definition 14; Jun and Kim (2015): Let $(\mathfrak{X}, *, \#, 0)$ be a pseudo BH-algebra, then I is called pseudo ideal of \mathfrak{X} , if it satisfies:

- $0 \in I$
- $x * y, x \# y \in I, y \in I \Rightarrow x \in I, \forall x, y \in \mathfrak{X}$

Definition 15; Jun and Kim (2015): A pseudo ideal I of a pseudo BH-algebra \mathfrak{X} is called a pseudo closed ideal of \mathfrak{X} , if for every $x \in I$, we have $0 * x, 0 \# x \in I$.

RESULTS AND DISCUSSION

In this study, we define a new types of a pseudo ideals, a pseudo essence subset and a pseudo essence ideal of a pseudo BH-algebra. Also, we study some propositions to other some types of a pseudo ideals of a pseudo BH-algebra.

Definition 1: A pseudo BH-algebra \mathfrak{X} is said a pseudo 0-commutative if:

- $x * (0 \# y) = y * (0 \# x)$
- $x \# (0 * y) = y \# (0 * x)$. For all $x, y, z \in \mathfrak{X}$

Example 1: Let $\mathfrak{X} = \{0, 1, 2\}$ be a set with the following Cayley Table 1. Then \mathfrak{X} is a pseudo BH-algebra.

Table 1: Pseudo 0-commutative

*	0	1	2
0	0	1	2
1	1	0	1
2	2	1	0

#	0	1	2
0	0	1	2
1	1	0	2
2	2	2	0

Definition 2: Let \mathfrak{X} be a pseudo BH-algebra. Then the set $G(\mathfrak{X}) = \{x \in \mathfrak{X} : 0 * x = 0 \# x = x\}$ is called a pseudo G-part of \mathfrak{X} .

Example 2: Let $\mathfrak{X} = \{0, 1, 2, 3\}$ be a set with the following Cayley Table 2.

Table 2: Pseudo G-part of \mathfrak{X}

*	0	1	2	3
0	0	1	2	3
1	1	0	2	3
2	2	1	0	1
3	3	3	3	0

#	0	1	2	3
0	0	1	2	3
1	1	0	2	3
2	2	2	0	2
3	3	3	3	1

Definition 3: Let \mathfrak{X} be a pseudo BH-algebra. Then the set $\mathfrak{X}_* = \{x \in \mathfrak{X} : 0 * x = 0 \# x = 0\}$ is called a BCA-part of \mathfrak{X} .

Example 3: Let $\mathfrak{X} = \{0, 1, 2, 3\}$ be a set with the following Cayley Table 3.

Table 3: Pseudo BCA-part of \mathfrak{X}

*	0	1	2	3
0	0	0	0	0
1	1	1	0	2
2	2	2	1	0
3	3	3	3	0

#	0	1	2	3
0	0	0	0	0
1	1	1	0	2
2	2	2	2	0
3	3	3	3	1

Definition 4: A nonempty subset I of a pseudo BH-algebra \mathfrak{X} is called a pseudo B-ideal of \mathfrak{X} if it satisfies:

- $0 \in I$
- $x * (z \# (0 * y)), y \in I$ imply $x * z \in I$
- $x \# (z - (0 \# y)), y \in I$ imply $x \# z \in I$

Example 4: Let $\mathfrak{X} = \{0, 1, 2, 3\}$ be a set with the following cayley (Table 4). Then, \mathfrak{X} is a pseudo BH-algebra and let $I = \{0, 1\}$ be a subset of \mathfrak{X} , then, it is a pseudo B-ideal of \mathfrak{X} .

Table 4: Pseudo B-ideal of \mathfrak{X}

*	0	1	2
0	0	0	0
1	1	0	2
2	2	1	0

Proposition 1: Let, \mathfrak{X} be a pseudo BH-algebra such that $\mathfrak{X} = \mathfrak{X}_+$, then, every pseudo ideal of \mathfrak{X} is a B-ideal of \mathfrak{X} .

Proof: Let, I be a pseudo ideal of \mathfrak{X} and $x^*(z \# (0*y))$, $y \in I$. For all $x, y, z \in \mathfrak{X}$. Since, $\mathfrak{X} = \mathfrak{X}_+ \Rightarrow x^*(z \# 0) \in I$. Since, \mathfrak{X} is a pseudo BH-algebra $\Rightarrow x^*z \in I$. Thus, $x^*z \in I$. Similarly, $x \# (z^*(0*y))$, $y \in I$ imply $x \# z \in I$. Hence, I is a pseudo B-ideal of \mathfrak{X} .

Proposition 2: Let \mathfrak{X} be a pseudo BH-algebra. If a pseudo B-ideal of \mathfrak{X} is a pseudo G-part of \mathfrak{X} then, it is a pseudo ideal of \mathfrak{X} .

Proof: Let, \mathfrak{X} be a pseudo ideal of \mathfrak{X} and let $x*y, x \# y \in I$, $y \in I$. For all $x, y, z \in \mathfrak{X}$. Since, I is pseudo G-part of $\mathfrak{X} \Rightarrow x^*0 \in I \Rightarrow x^*(0 \# (0*y)) \in I$ and $y \in I$. Since, I is a pseudo B-ideal of $\mathfrak{X} \Rightarrow x^*0 \in I$. Since, \mathfrak{X} is a pseudo BH-algebra $\Rightarrow x \in I$. Similarly, $x \# y \in I$, $y \in I \Rightarrow x \in I$. Hence, I is a pseudo ideal of \mathfrak{X} .

Proposition 3: Let \mathfrak{X} be a pseudo BH-algebra such that $y = z \# (0*y)$ and $y = z^*(0 \# y)$, then every pseudo B-ideal of \mathfrak{X} is a pseudo ideal of \mathfrak{X} .

Proof: Let I be a pseudo B-ideal of \mathfrak{X} and $x^*(z \# (0*y))$, $y \in I$. For all $x, y, z \in \mathfrak{X}$. Since, $y = z \# (0*y)$ then $x*y, y \in I$ imply $x \in I$. Similarly, $x \# (z^*(0 \# y))$, $y \in I$ imply $x \in I$. Hence, I is a pseudo ideal of \mathfrak{X} .

Definition 5: A non empty subset I of a pseudo BH-algebra \mathfrak{X} is called a pseudo H-ideal of \mathfrak{X} if it satisfies:

- $0 \in I$
- $(x*y) \# (x*z) \in I$ and $y \in I \Rightarrow x \in I$
- $(x \# y)^*(x \# z) \in I$ and $y \in I \Rightarrow x \in I$. For all $x, y, z \in \mathfrak{X}$

Example 5: Let, $\mathfrak{X} = \{0, 1, 2, 3\}$ be a set with the following Cayley Table 5. Then, \mathfrak{X} is a pseudo BH-algebra and let $I = \{0, 1\}$ be a subset of \mathfrak{X} , then it is a pseudo H-ideal of \mathfrak{X} .

Table 5: Pseudo H-ideal of \mathfrak{X}

*	0	1	2	3
0	0	1	2	3
1	1	0	2	3
2	2	1	0	2
3	3	0	0	0

Proposition 4: Every pseudo H-ideal of a pseudo BH-algebra \mathfrak{X} is a pseudo ideal of \mathfrak{X} .

Proof: Let, I be a pseudo H*-ideal of \mathfrak{X} and let $x*y, x \# y \in I$ and $y \in I$. For all $x, y, z \in \mathfrak{X}$. Since, \mathfrak{X} is a pseudo BH-algebra $\Rightarrow (x*y) \in I \Rightarrow ((x*y) \# 0) \in I \Rightarrow (x*y) \# (x*x) \in I$ and $y \in I$. Since, I is a pseudo H-ideal $\Rightarrow x \in I$. Similarly, $x \# y \in I$ and $y \in I \Rightarrow x \in I$. Hence, I is a pseudo ideal of \mathfrak{X} .

Definition 6: Let, \mathfrak{X} be a pseudo BH-algebra. For a subsets A and B of \mathfrak{X} , then $A*B$ and $A \# B$ are defined as follows:

- $A*B = \{x*y: x \in A, y \in B\}$
- $A \# B = \{x \# y: x \in A, y \in B\}$

Proposition 5: Let, \mathfrak{X} be a pseudo BH-algebra.

- If $0 \in B \subseteq \mathfrak{X}$. Then $\forall B \subseteq \mathfrak{X}$, we have $B \subseteq A*B$ and $B \subseteq A \# B$
- If $0 \in A \subseteq \mathfrak{X}$. Then $\forall B \subseteq \mathfrak{X}$, we have $B \subseteq A*B$ and $B \subseteq A \# B$

Proof: Let, $x \in A$. Since, \mathfrak{X} is a pseudo BH-algebra, then, $x = x^*0 \in A*B$ and $x = x \# 0 \in A \# B$. Hence, $(A \subseteq A*B)$ $(A \subseteq A \# B)$. Similarly of (1).

Definition 7: If A is a nonempty subset of a pseudo BH-algebra \mathfrak{X} satisfies $A*\mathfrak{X} = A$ and $A \# \mathfrak{X} = A$, then A is called a pseudo essence subset of \mathfrak{X} . If A is a pseudo ideal of \mathfrak{X} , then it is called a pseudo essence ideal of \mathfrak{X} .

Example 6: Let, $\mathfrak{X} = \{0, 1, 2, 3\}$ be a set with the following Cayley Table 6.

Table 6: Pseudo essence of \mathfrak{X}

*	0	1	2	3
0	0	0	0	0
1	1	1	0	2
2	2	2	1	0
3	3	3	3	0

Then \mathfrak{X} is a pseudo BH-algebra. Let, $A = \{0, 1\}$, $B = \{0, 2\}$ and $C = \{0, 1, 2\}$ then A, B and C are a pseudo

essence subset of \mathfrak{X} . But $D = \{0, 3\}$ is not a pseudo essence subset of \mathfrak{X} , since, $3*2 = 1 \notin D$ and $3\#2 = 2 \notin D$.

Proposition 6: Let, \mathfrak{X} be a pseudo BH-algebra. Then every pseudo essence ideal of \mathfrak{X} is a pseudo essence subset of \mathfrak{X} .

Proof: Let, A be a pseudo ideal of \mathfrak{X} and let $x, y \in A$. Since, $x*y \in A \subseteq A^*A \subseteq A^*\mathfrak{X} = A$ and $x\#y \in A \subseteq A\#A \subseteq A\#\mathfrak{X} = A$. Hence, A is a pseudo essence subset of \mathfrak{X} .

Remark 1: The converse of proposition (6) may be not true in general as follows in example (1), since, A is a pseudo essence $1*3 = 0 \in A$ and $1 \in A$ but $3 \notin A$ and $1\#3 = 1 \in A$ and $1 \in A$ but $3 \notin A$.

Proposition 7: Let, \mathfrak{X} be a pseudo BH-algebra. Then, every pseudo essence ideal of \mathfrak{X} is a pseudo closed ideal of \mathfrak{X} .

Proof: Let, A be a pseudo essence ideal of \mathfrak{X} , then, $0 \in A$. Let, $x \in A$, then, $0*x \in A^*A \subseteq A^*\mathfrak{X} = A$. Thus, $0*x \in A$, similarly, $0\#x \in A$. Hence, A is a pseudo essence closed of \mathfrak{X} .

Definition 8: A nonempty subset I of a pseudo BH-algebra \mathfrak{X} . Then, I is called pseudo BH-ideal of \mathfrak{X} if it satisfies:

- $0 \in I$
- $x*y, x\#y \in I$ and $y \in I$ imply $x \in I$
- $x \in I$ and $y \in I$ and imply $x*y, x\#y \in I, I*\mathfrak{X}, I\#\mathfrak{X} \subseteq I$. For all $x, y \in \mathfrak{X}$

Proposition 8: Let, \mathfrak{X} be a pseudo BH-algebra. Then every a pseudo essence ideal of \mathfrak{X} is a pseudo BH-ideal of \mathfrak{X} .

Proof: Let, A be pseudo essence ideal of \mathfrak{X} . Since, $A^*\mathfrak{X} = A$, then $A^*\mathfrak{X} \subseteq A$ and $A\#\mathfrak{X} = A$, then $A\#\mathfrak{X} \subseteq A$. Thus, A is a pseudo essence ideal of \mathfrak{X} and $A^*\mathfrak{X}, A\#\mathfrak{X} \subseteq A$. Hence, A is a pseudo pseudo BH-ideal of \mathfrak{X} .

CONCLUSION

In this study, the notions of pseudo B-ideal, pseudo H-ideal and pseudo essence of a pseudo BH-algebra are introduced. Furthermore, the results are examined in terms of the relationship between pseudo B-idea, pseudo H-ideal and pseudo essence of a pseudo BH-algebra.

REFERENCES

- Abbass, H.H. and A.A. Hamza, 2017. On U-BG-filter of a U-BG-BH-algebra. Appl. Math. Sci., 11: 1297-1305.
- Abbass, H.H. and H.A. Dahham, 2012. Some types of fuzzy ideals with respect to an element of a BG-algebra. MSc Thesis, University of Kufa, Kufa, Iraq.
- Abbass, H.H. and L.S. Mahdi, 2014. A new class of BH-algebra. MSc. Thesis, University of Kufa, Kufa, Iraq.
- Jun, Y.B. and S.S. Kim, 2015. On pseudo BH-algebra. Honam Math. J., 37: 207-219.
- Jun, Y.B., E.H. Roh and H.S. Kim, 1998. On BH-algebras. Sci. Math., 1: 347-534.
- Kim, E.M. and S.S. Ahn, 2011. An application of complicationness to BH-algebra. J. Korea Soc. Math. Educ., 4: 293-304.