

Asymptotic Behavior of Eigenvalue and Eigenfunction of a Six Order Boundary Value Problem

Karwan H.F. Jwamer and Khelan H. Qadr
 Department of Mathematics, College of Science, University of Sulaimani,
 Kurdistan Region, Sulaymaniyah, Iraq

Abstract: We consider differential operators with separate boundary conditions in this study. And we find a new expression and their derivative of six linearly independent solution. And it will also prove the existence of uniqueness for separate boundary conditions. We obtain asymptotic formulas for the individual values and functions of these problems with boundary value where the potential $q(x)$ is an arbitrary complex function valued in $[a, b]$.

Key words: Differential operator, eigenvalue, eigenfunctions, spectral parameter, arbitrary complex function, potential

INTRODUCTION

It is well known that many researchers have investigated the spectral properties of the Sturm-Liouville operator generated by the separate boundary condition (Aigunov, 1996) and many researchers have found asymptotic formula for the Sturm-Liouville operator's eigenvalues and functions in the case of periodic and anti-periodic boundary conditions (Menken, 2010; Naimark, 1967; Moller and Zinsou, 2012; Jwamer and Aigounv, 2010; Aigounov and Tamila, 2009; Aigunov, 1996 and Tamarkin, 1928). Many researchers have been interested in the ongoing Sturm-Liouville issue in recent years as we see N.B. Kermov, H. Menken by Menken (2010), Karwan and Rando (2017) found upper bound with smooth coefficients for the proper functions of the fourth boundary value issue. In this study, we consider the differential operator:

$$y^{(6)}(x) + q(x)y(x) = \lambda^6 y(x)$$

$$U_i(y) = \sum_{j=0}^5 a_{ij} y^{(j)}(b, \lambda), i = 0, 1, 2$$

where, a_{ij} are real numbers:

$$U_3(y) = \sum_{i=1}^6 (\lambda)^{i-1} y^{(6-i)}(b, \lambda)$$

$$U_4(y) = \sum_{i=1}^6 (-\lambda)^{i-1} y^{(6-i)}(b, \lambda)$$

$$U_5(y) = \sum_{i=1}^6 \left(\frac{\lambda}{\left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)} \right)^{i-1} y^{(6-i)}(b, \lambda)$$

Where:

λ : A spectral parameter

$q(x)$: An arbitrary complex valued function

FUNDAMENTAL DIFFERENTIAL EQUATION SOLUTIONS SYSTEMS

The expressions of six linearly independent solutions and their derivatives can be found in this study.

Theorem 1: The fundamental system of solutions of linear differential equation:

$$y^{(6)}(x) + q(x)y(x) = \lambda^6 y(x) \quad (1)$$

Are $y_0(x, \lambda), y_1(x, \lambda), y_2(x, \lambda), y_3(x, \lambda), y_4(x, \lambda), y_5(x, \lambda)$ that satisfy the initial conditions:

$$y_i^{(n)}(0, \lambda) = \begin{cases} 1 & \text{if } i = n \\ 0 & \text{if } i \neq n \end{cases} \quad (2)$$

Where:

$$\begin{aligned} y_0 &= \frac{1}{3} \left[\cosh \lambda x + \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x + \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x \right] + \frac{1}{3\lambda^5} \\ &\int_0^x \left[\sinh \lambda(x-\xi) + \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-\xi) + \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-\xi) \right] \\ &q(\xi) y_0(\xi) d\xi \end{aligned}$$

$$y_1 = \left[\frac{1}{3\lambda} \sinh \lambda x + \frac{1}{6\lambda} \left(1 + \frac{1}{32} i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x + \frac{1}{6\lambda} \left(-1 + \frac{1}{32} i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x \right] + \frac{1}{3\lambda^5}$$

$$\int_a^b \left[\sinh \lambda(x-\xi) + \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) + \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) \right] q(\xi) y_0(\xi) d\xi$$

$$y_2 = \left[\frac{1}{3\lambda^2} \cosh \lambda x + \frac{1}{6\lambda^2} \left(-1 + \frac{1}{32} i \right) \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x + \frac{1}{6\lambda^2} \left(1 - \frac{1}{32} i \right) \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x \right] + \frac{1}{3\lambda^5}$$

$$\int_0^x \left[\sinh \lambda(x-\xi) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) \right] q(\xi) y_0(\xi) d\xi$$

$$y_3 = \frac{1}{3\lambda^3} \left[\sinh \lambda x - \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x + \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x \right] + \frac{1}{3\lambda^5}$$

$$\int_0^x \left[\sinh \lambda(x-\xi) + \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) + \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) \right] q(\xi) y_0(\xi) d\xi$$

$$y_4 = \left[\frac{1}{3\lambda^4} \cosh \lambda x + \frac{1}{6\lambda^4} \left(1 - \frac{1}{32} i \right) \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x + \frac{1}{6\lambda^4} \left(-1 + \frac{1}{32} i \right) \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x \right] + \frac{1}{3\lambda^5}$$

$$\int_0^x \left[\sinh \lambda(x-\xi) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) \right] q(\xi) y_0(\xi) d\xi$$

$$y_5 = \left[\frac{1}{3\lambda^5} \sinh \lambda x - \frac{1}{6\lambda^5} \left(-1 + \frac{1}{32} i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x - \frac{1}{6\lambda^5} \left(1 + \frac{1}{32} i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x \right] + \frac{1}{3\lambda^5}$$

$$\int_0^x \left[\sinh \lambda(x-\xi) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) \right] q(\xi) y_0(\xi) d\xi$$

Proof: Consider the linear differential operator:

$$I(y) = -y^{(6)}(x) + q(x)y(x) \quad (3)$$

$$w_0 = 1, w_1 = \frac{1}{2} + i \frac{\sqrt{3}}{2}, w_2 = -\frac{1}{2} + i \frac{\sqrt{3}}{2}, \\ w_3 = -1, w_4 = -\frac{1}{2} - i \frac{\sqrt{3}}{2}, w_5 = \frac{1}{2} - i \frac{\sqrt{3}}{2}$$

We want to find a solution that is not zero:

$$I(y) - \lambda^6 y(x) = 0$$

Which satisfy the initial conditions Eq. 2. First, we reduce Eq. 3 to an integro-differential equations:

$$y^{(6)}(x) + \lambda^6 y(x) = q(x)y(x) \\ m(y) = y^{(6)}(x) + \lambda^6 y(x), m(y) = q(x)y(x) \quad (4)$$

The homogeneous linear differential equation $y^{(6)}(x) + \lambda^6 y(x) = 0$ has for $\lambda \neq 0$ the solutions:

$$e^{\lambda w_0 x}, e^{\lambda w_1 x}, e^{\lambda w_2 x}, e^{\lambda w_3 x}, e^{\lambda w_4 x}, e^{\lambda w_5 x}$$

Where:

Then by using the method of variation of parameters we can express the solutions of Eq. 4 for $k = 0, 1, 2, 3, 4, 5$ as:

$$y_k = c_0 e^{\lambda x} + c_1 e^{\lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x} + c_2 e^{\lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x} + c_3 e^{-\lambda x} + c_4 e^{\lambda \left(\frac{1}{2} - i \frac{\sqrt{3}}{2} \right) x} + c_5 e^{\lambda \left(-\frac{1}{2} - i \frac{\sqrt{3}}{2} \right) x} + \sinh(x-\xi) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi) + \int_0^x \frac{\left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-\xi)}{3\lambda^5} q(\xi) y_k(x, \lambda) d\xi \quad (5)$$

We apply Eq. 5 and in Eq. 2, then, we get. For $k = 0$ then:

$$y_0(0, \lambda) = 1, y'_0(0, \lambda) = 0, y''_0(0, \lambda) = 0,$$

$$y'''_0(0, \lambda) = 0, y^{(4)}_0(0, \lambda) = 0, y^{(5)}_0(0, \lambda) = 0$$

$$c_0 + c_1 + c_2 + c_3 + c_4 + c_5 = 1$$

$$\lambda c_0 + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \lambda c_1 + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \lambda c_2 -$$

$$\lambda c_3 + \left(-\frac{1}{2} - i \frac{\sqrt{3}}{2} \right) \lambda c_4 + \left(\frac{1}{2} - i \frac{\sqrt{3}}{2} \right) \lambda c_5 = 0$$

$$\lambda^2 c_0 + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \lambda^2 c_1 + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \lambda^2 c_2 +$$

$$\lambda^2 c_3 + \left(-\frac{1}{2} - i \frac{\sqrt{3}}{2} \right)^2 \lambda^2 c_4 + \left(\frac{1}{2} - i \frac{\sqrt{3}}{2} \right)^2 \lambda^2 c_5 = 0$$

$$\lambda^3 c_0 + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \lambda^3 c_1 + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \lambda^3 c_2 - \lambda^3 c_3 +$$

$$\left(-\frac{1}{2} - i \frac{\sqrt{3}}{2} \right)^3 \lambda^3 c_4 + \left(\frac{1}{2} - i \frac{\sqrt{3}}{2} \right)^3 \lambda^3 c_5 = 0$$

$$\lambda^4 c_0 + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \lambda^4 c_1 + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \lambda^4 c_2 +$$

$$\lambda^4 c_3 + \left(-\frac{1}{2} - i \frac{\sqrt{3}}{2} \right)^4 \lambda^4 c_4 + \left(\frac{1}{2} - i \frac{\sqrt{3}}{2} \right)^4 \lambda^4 c_5 = 0$$

$$\lambda^5 c_0 + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \lambda^5 c_1 + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \lambda^5 c_2 - \lambda^5 c_3 +$$

$$\left(-\frac{1}{2} - i \frac{\sqrt{3}}{2} \right)^5 \lambda^5 c_4 + \left(\frac{1}{2} - i \frac{\sqrt{3}}{2} \right)^5 \lambda^5 c_5 = 0$$

We can solve it for c_i and we get $c_i = 1/6$ for each $k = 0:5$ then y_0 has the form:

$$y_0 = \frac{1}{3} \left[\cosh \lambda x + \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x + \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x \right] + \frac{1}{3\lambda^5}$$

$$\int_0^x \left[\sinh \lambda(x-\xi) + \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-\xi) \right] q(\xi) y_0(\xi) d\xi$$

And we can use the same technique for $y_1(x, \lambda)$, $y_2(x, \lambda)$, $y_3(x, \lambda)$, $y_4(x, \lambda)$, $y_5(x, \lambda)$.

Corollary 1: For x in $[a, b]$ and $\lambda \neq 0$, $y_0(x, \lambda)$ and its derivatives in theorem 1 can be written as:

$$y_0 = \frac{1}{3}$$

$$\left[\cosh \lambda(x-a) + \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \right. \\ \left. \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^5} e^{|t|(x-a)} \right)$$

$$y'_0 = \frac{1}{3} \lambda \left[\begin{array}{l} \sinh \lambda(x-a) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ (x-a) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] + \\ 0 \left(\frac{1}{\lambda^4} e^{|t|(x-a)} \right)$$

$$y''_0 = \frac{1}{3} \lambda^3$$

$$\left[\begin{array}{l} \cosh \lambda(x-a) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \\ \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] + \\ 0 \left(\frac{1}{\lambda^3} e^{|t|(x-a)} \right)$$

$$y'''_0 = \frac{1}{3} \lambda^3$$

$$\left[\begin{array}{l} \sinh \lambda(x-a) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \\ \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] + \\ 0 \left(\frac{1}{\lambda^2} e^{|t|(x-a)} \right)$$

$$y^{(4)}_0 = \frac{1}{3} \lambda^4$$

$$\left[\begin{array}{l} \cosh \lambda(x-a) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ (x-a) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] + \\ 0 \left(\frac{1}{\lambda} e^{|t|(x-a)} \right)$$

$$y^{(5)}_0 = \frac{1}{3} \lambda^5$$

$$\left[\begin{array}{l} \sinh \lambda(x-a) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ (x-a) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] + \\ 0 \left(e^{|t|(x-a)} \right)$$

Proof: Since, $y_0(x, \lambda)$ in theorem 1 has the form:

$$y_0 = \frac{1}{3} \left[\cosh \lambda(x-a) + \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + \frac{1}{3\lambda^5} \\ \int_a^x \left[\sinh \lambda(x-\xi) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-\xi) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-\xi) \right] q(\xi) y_0(\xi) d\xi$$

Since, $|\cosh z| \leq e^{|Re(z)|}$ and $|\sinh z| \leq e^{|Re(z)|}$:

$$|\sinh \lambda(x-\xi)| \leq e^{|\sigma(x-\xi)|}, \left| \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-\xi) \right| \leq e^{\left| \left(\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)(x-\xi) \right|} \\ \left| \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-\xi) \right| \leq e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)(x-\xi) \right|}$$

where, $\lambda = \sigma + it$:

$$|\cosh \lambda x| \leq e^{|\sigma x|}, \left| \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)x \right| \leq e^{\left| \left(\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)x \right|}, \left| \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)x \right| \leq e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)x \right|} \\ y_0(x, \lambda) \leq \frac{1}{3} \left[|\cosh \lambda x| + \left| \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)x \right| + \left| \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)x \right| \right] + \frac{1}{3\lambda^5} \\ \int_a^x \left[|\sinh \lambda(x-\xi)| + \left| \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-\xi) \right| + \left| \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-\xi) \right| \right] q(\xi) y_0(\xi) d\xi$$

Then:

$$|y_0(x, \lambda)| \leq \frac{1}{3} \left[e^{|\sigma x|} + e^{\left| \left(\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)x \right|} + e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)x \right|} \right] + \frac{1}{3\lambda^5} \\ \int_a^x \left[e^{|\sigma(x-\xi)|} + \left| \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right| e^{\left| \left(\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)(x-\xi) \right|} + \left| \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right| e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)(x-\xi) \right|} \right] q(\xi) y_0(\xi, \lambda) d\xi$$

And since:

$$e^{|\sigma x|} \leq e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)x \right|}, e^{\left| \left(\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)x \right|} \leq e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)x \right|} \\ |y_0(x, \lambda)| \leq e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)x \right|} + \frac{1}{3\lambda^5} \\ \int_a^x \left[e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)(x-\xi) \right|} + e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)(x-\xi) \right|} \right] q(\xi) y_0(\xi, \lambda) d\xi$$

Put:

$$|y_0(x, \lambda)| = e^{\left| \left(-\frac{1}{2}\sigma - \frac{\sqrt{3}}{2}t \right)x \right|} f(x, \lambda)$$

$$f(x, \lambda) = y_0(x, \lambda) e^{\left| \left(\frac{1}{2}\sigma + \frac{\sqrt{3}}{2}t \right)(x-\xi) \right|}$$

$$f(x, \lambda) = y_0 = \frac{1}{3} \left[\cosh \lambda x + \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x + \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) x \right] e^{\left| \left(\frac{1}{2} \sigma + \frac{\sqrt{3}}{2} t \right) (x - \xi) \right|} +$$

$$\frac{1}{3\lambda^5} \int_a^x \left[\sinh \lambda (x - \xi) + \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x - \xi) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x - \xi) \right] e^{\left| \left(\frac{1}{2} \sigma + \frac{\sqrt{3}}{2} t \right) (x - \xi) \right|} q(\xi) f(\xi, \lambda) d\xi$$

Let, $M(\lambda)$ denote the maximum value of $|f(x, \lambda)|$ for x in $[a, b]$ then we obtain:

$$M(\lambda) \leq 1 + \frac{1}{|\lambda|^5} M(\lambda) \int_a^x |q(\xi)| d\xi$$

$$M(\lambda) \left(1 - \frac{1}{|\lambda|^5} M(\lambda) \int_a^x |q(\xi)| d\xi \right) \leq 1$$

$$M(\lambda) \leq \left\{ 1 - \frac{1}{|\lambda|^5} M(\lambda) \int_a^x |q(\xi)| d\xi \right\}^{-1}$$

$|f(x, \lambda)| \leq M(\lambda) = 0$ (1), $|\lambda| \rightarrow \infty$ and therefore:

$$y_0(x, \lambda) = 0 \left\{ e^{\left| \left(-\frac{1}{2} \sigma - \frac{\sqrt{3}}{2} t \right) (x - \xi) \right|} \right\}$$

So, the integral equation is:

$$0 \left\{ \frac{1}{\lambda^5} \int_a^x e^{\left| \left(-\frac{1}{2} \sigma - \frac{\sqrt{3}}{2} t \right) (x - \xi) \right|} e^{\left| \left(-\frac{1}{2} \sigma - \frac{\sqrt{3}}{2} t \right) (\xi - a) \right|} q(\xi) d\xi \right\} =$$

$$0 \left\{ \lambda^{-5} e^{\left| \left(-\frac{1}{2} \sigma - \frac{\sqrt{3}}{2} t \right) (x - a) \right|} \right\}$$

By the same way we get all derivatives.

Corollary 2: For x in $[a, b]$ and $\lambda \neq 0$, $y_1(x, \lambda)$ and its derivatives in theorem 1 can be written as:

$$y_1 = \frac{1}{3\lambda} \left[\sinh \lambda (x - a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right] +$$

$$\left[(x - a) + \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x - a) \right]$$

$$0 \left(\frac{1}{\lambda^6} e^{|t|(x-a)} \right)$$

$$y'_1 = \frac{1}{3}$$

$$\left[\begin{array}{l} \cosh \lambda (x - a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x - a) + \frac{1}{2} \\ \left(-1 + \frac{1}{32} i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x - a) \end{array} \right] +$$

$$0 \left(\frac{1}{\lambda^5} e^{|t|(x-a)} \right)$$

$$y''_1 = \frac{1}{3} \lambda$$

$$\left[\begin{array}{l} \sinh \lambda (x - a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ (x - a) + \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x - a) \\ 0 \left(\frac{1}{\lambda^4} e^{|t|(x-a)} \right) \end{array} \right]$$

$$y'''_1 = \frac{1}{3} \lambda^2$$

$$\left[\begin{array}{l} \cosh \lambda (x - a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \\ \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x - a) + \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \\ \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x - a) \\ 0 \left(\frac{1}{\lambda^3} e^{|t|(x-a)} \right) \end{array} \right]$$

$$y^{(4)}_1 = \frac{1}{3} \lambda^3$$

$$\left[\begin{array}{l} \sinh \lambda (x - a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ (x - a) + \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x - a) \\ 0 \left(\frac{1}{\lambda^2} e^{|t|(x-a)} \right) \end{array} \right]$$

$$y_1^{(5)} = \frac{1}{3} \lambda^4 \left[\cosh \lambda(x-a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda} e^{|t|(x-a)} \right)$$

Corollary 3: For x in $[a, b]$ and $\lambda \neq 0$, $y_2(x, \lambda)$ and its derivatives in theorem 1 can be written as:

$$\begin{aligned} y_2 &= \frac{1}{3\lambda^2} \left[\cosh \lambda(x-a) + \frac{1}{2} \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \frac{1}{2} \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^7} e^{|t|(x-a)} \right) \\ y'_2 &= \frac{1}{3\lambda} \left[\sinh \lambda(x-a) + \frac{1}{2} \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \frac{1}{2} \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^6} e^{|t|(x-a)} \right) \\ y''_2 &= \frac{1}{3} \left[\cosh \lambda(x-a) + \frac{1}{2} \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \frac{1}{2} \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^5} e^{|t|(x-a)} \right) \\ y'''_2 &= \frac{1}{3} \lambda \left[\sinh \lambda(x-a) + \frac{1}{2} \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \frac{1}{2} \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^4} e^{|t|(x-a)} \right) \\ y^{(4)}_2 &= \frac{1}{3} \lambda^2 \left[\cosh \lambda(x-a) + \frac{1}{2} \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \frac{1}{2} \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^3} e^{|t|(x-a)} \right) \\ y^{(5)}_2 &= \frac{1}{3} \lambda^3 \left[\sinh \lambda(x-a) + \frac{1}{2} \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \frac{1}{2} \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^2} e^{|t|(x-a)} \right) \end{aligned}$$

Corollary 4: For x in $[a, b]$ and $\lambda \neq 0$, $y_3(x, \lambda)$ and its derivatives in theorem 1 can be written as:

$$\begin{aligned} y_3 &= \frac{1}{3\lambda^3} \left[\sinh \lambda(x-a) - \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^8} e^{|t|(x-a)} \right) \\ y'_3 &= \frac{1}{3\lambda^2} \left[\cosh \lambda(x-a) - \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^7} e^{|t|(x-a)} \right) \\ y''_3 &= \frac{1}{3\lambda} \left[\sinh \lambda(x-a) - \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^6} e^{|t|(x-a)} \right) \\ y'''_3 &= \frac{1}{3} \left[\cosh \lambda(x-a) - \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + 0 \left(\frac{1}{\lambda^5} e^{|t|(x-a)} \right) \end{aligned}$$

$$y_3^{(4)} = \frac{1}{3\lambda} \left[\sinh \lambda(x-a) - \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^4} e^{|t|(x-a)} \right)$$

$$y_3^{(5)} = \frac{1}{3} \lambda^2 \left[\cosh \lambda(x-a) - \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^3} e^{|t|(x-a)} \right)$$

Corollary 5: For x in $[a, b]$ and $\lambda \neq 0$, $y_4(x, \lambda)$ and its derivatives in theorem 1 can be written as:

$$y_4 = \frac{1}{3\lambda^4} \left[\cosh \lambda(x-a) + \frac{1}{2} \left(1 - \frac{1}{32}i \right) \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^9} e^{|t|(x-a)} \right)$$

$$y'_4 = \frac{1}{3\lambda^3} \left[\sinh \lambda(x-a) + \frac{1}{2} \left(1 - \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^8} e^{|t|(x-a)} \right)$$

$$y''_4 = \frac{1}{3\lambda^2} \left[\cosh \lambda(x-a) + \frac{1}{2} \left(1 - \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^7} e^{|t|(x-a)} \right)$$

$$y'''_4 = \frac{1}{3\lambda} \left[\sinh \lambda(x-a) + \frac{1}{2} \left(1 - \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^6} e^{|t|(x-a)} \right)$$

$$y_4^{(4)} = \frac{1}{3} \left[\cosh \lambda(x-a) + \frac{1}{2} \left(1 - \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^5} e^{|t|(x-a)} \right)$$

$$y_4^{(5)} = \frac{1}{3} \lambda \left[\sinh \lambda(x-a) + \frac{1}{2} \left(1 - \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^4} e^{|t|(x-a)} \right)$$

Corollary 6: For x in $[a, b]$ and $\lambda \neq 0$, $y_5(x, \lambda)$ and its derivatives in theorem 1 can be written as:

$$y_5 = \frac{1}{3\lambda^5} \left[\sinh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) - \frac{1}{2} \left(1 + \frac{1}{32}i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^{10}} e^{|t|(x-a)} \right)$$

$$y'_5 = \frac{1}{3\lambda^4} \left[\cosh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) - \frac{1}{2} \left(1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \right] + 0 \left(\frac{1}{\lambda^9} e^{|t|(x-a)} \right)$$

$$y_5'' = \frac{1}{3\lambda^3}$$

$$\begin{aligned} & \left[\sinh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right] + \\ & \left[(x-a) - \frac{1}{2} \left(1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^2 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + \\ & 0 \left(\frac{1}{\lambda^8} e^{|t|(x-a)} \right) \end{aligned}$$

$$y_5''' = \frac{1}{3\lambda^2}$$

$$\begin{aligned} & \left[\cosh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right] + \\ & \left[(x-a) - \frac{1}{2} \left(1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^3 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + \\ & 0 \left(\frac{1}{\lambda^7} e^{|t|(x-a)} \right) \end{aligned}$$

$$y_5^{(4)} = \frac{1}{3\lambda}$$

$$\begin{aligned} & \left[\sinh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right] + \\ & \left[(x-a) - \frac{1}{2} \left(1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^4 \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + \\ & 0 \left(\frac{1}{\lambda^6} e^{|t|(x-a)} \right) \end{aligned}$$

$$y_5^{(5)} = \frac{1}{3}$$

$$\begin{aligned} & \left[\cosh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right] + \\ & \left[(x-a) - \frac{1}{2} \left(1 + \frac{1}{32}i \right) \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^5 \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + \\ & 0 \left(\frac{1}{\lambda^5} e^{|t|(x-a)} \right) \end{aligned}$$

Theorem 2: The formula of eigenfunctions $\psi_n(x)$ as $n \rightarrow \infty$ is:

$$\begin{aligned} \psi(x) = & \mp \frac{1}{3} \left(\frac{36864}{3071(b-a)} \right)^2 \\ & \left[\sinh \lambda(x-a) + \frac{1}{2} \left(1 + \frac{1}{32}i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right] + 0 \left(\frac{1}{\lambda^2} \right) \\ & \left[(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] \end{aligned}$$

where $a_{12} = 0$ and $a_{11} \neq 0$.

Proof: If $\psi(x) = c_0 y_0 + c_1 y_1 + c_2 y_2 + c_3 y_3 + c_4 y_4 + c_5 y_5$ then from first boundary condition we get $a_{11}c_0 + a_{12}c_1 + a_{13}c_2 + a_{14}c_3 + a_{15}c_4 + a_{16}c_5 = 0$, since, c_i cannot be zero for all i then. We can choose $c_0 = k_1 a_{12}$ and $c_1 = -k_1 a_{11}$ and $c_2 = k_2 a_{14}$ and $c_3 = -k_2 a_{13}$ and $c_4 = k_3 a_{16}$ and $c_5 = -k_3 a_{15}$ where $k_i \neq 0$ then from theorem 1 we obtain that:

$$\begin{aligned} \psi(x) = & \frac{1}{3} k_1 a_{12} \left[\begin{aligned} & \cosh \lambda(x-a) + \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ & (x-a) + \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \end{aligned} \right] + 0 \left(\frac{1}{\lambda^5} e^{|t|(x-a)} \right) - \\ & \frac{1}{3\lambda} k_1 a_{11} \left[\begin{aligned} & \sinh \lambda(x-a) + \frac{1}{2} \left(1 + \frac{1}{32}i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ & (x-a) + \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \end{aligned} \right] + \\ & 0 \left(\frac{1}{\lambda^6} e^{|t|(x-a)} \right) + \frac{1}{3\lambda^2} k_1 a_{14} \left[\begin{aligned} & \cosh \lambda(x-a) + \frac{1}{2} \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ & (x-a) + \frac{1}{2} \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \end{aligned} \right] + \\ & 0 \left(\frac{1}{\lambda^7} e^{|t|(x-a)} \right) - k_1 a_{13} \frac{1}{3\lambda^3} \left[\begin{aligned} & \sinh \lambda(x-a) - \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ & (x-a) + \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \end{aligned} \right] + \\ & 0 \left(\frac{1}{\lambda^8} e^{|t|(x-a)} \right) - k_1 a_{16} \frac{1}{3\lambda^4} \left[\begin{aligned} & \cosh \lambda(x-a) + \frac{1}{2} \left(1 - \frac{1}{32}i \right) \cosh \lambda \\ & \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) + \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \\ & \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \end{aligned} \right] + \\ & 0 \left(\frac{1}{\lambda^9} e^{|t|(x-a)} \right) - k_1 a_{15} \frac{1}{3\lambda^5} \left[\begin{aligned} & \sinh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \\ & \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) - \frac{1}{2} \left(1 + \frac{1}{32}i \right) \\ & \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \end{aligned} \right] + \\ & 0 \left(\frac{1}{\lambda^{10}} e^{|t|(x-a)} \right) \end{aligned}$$

$$\begin{aligned} \psi(x) = & \frac{1}{3} k_1 a_{12} \\ & \left[\cosh \lambda(x-a) + \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right] + 0 \left(\frac{1}{\lambda^5} \right) \\ & \left[(x-a) + \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] \end{aligned}$$

$$\begin{aligned} & -0\left(\frac{1}{\lambda}\right) + 0\left(\frac{1}{\lambda^6}\right) + 0\left(\frac{1}{\lambda^2}\right) + 0\left(\frac{1}{\lambda^7}\right) - 0\left(\frac{1}{\lambda^3}\right) \\ & + 0\left(\frac{1}{\lambda^8}\right) + 0\left(\frac{1}{\lambda^4}\right) + 0\left(\frac{1}{\lambda^9}\right) - 0\left(\frac{1}{\lambda^5}\right) + 0\left(\frac{1}{\lambda^{10}}\right) \\ \psi(x) = \frac{1}{3}k_1 a_{12} & \left[\begin{array}{l} \cosh \lambda(x-a) + \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \\ (x-a) + \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] + 0\left(\frac{1}{\lambda}\right) \end{aligned}$$

If $a_{12} = 0$ then $a_{11} \neq 0$:

$$\begin{aligned} \psi(x) = & -\frac{1}{3\lambda^2} k_2 a_{14} \\ & \left[\begin{array}{l} \cosh \lambda(x-a) + \frac{1}{2} \cosh \lambda \\ \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \end{array} \right] + 0\left(\frac{1}{\lambda^7} e^{|t|(x-a)}\right) \\ & \left[\begin{array}{l} \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] \end{aligned}$$

If $a_{12} = 0, a_{11} = 0, a_{14} = 0$ and $a_{13} \neq 0$:

$$\begin{aligned} \psi(x) = -\frac{1}{3\lambda} k_1 a_{11} & \left[\begin{array}{l} \sinh \lambda(x-a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \\ \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \\ \left(-1 + \frac{1}{32} i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] + 0\left(\frac{1}{\lambda^2}\right) \end{aligned}$$

$$\begin{aligned} \psi(x) = -\frac{1}{3\lambda^3} k_2 a_{13} & \left[\begin{array}{l} \sinh \lambda(x-a) - \sinh \lambda \\ \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \sinh \lambda \\ \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] + 0\left(\frac{1}{\lambda^8} e^{|t|(x-a)}\right) \end{aligned}$$

If $a_{12} = 0, a_{11} = 0$ and $a_{14} \neq 0$:

If $a_{12} = 0, a_{11} = 0, a_{14} = 0, a_{13} = 0$ and $a_{15} \neq 0$:

$$\begin{aligned} \psi(x) = -\frac{1}{3\lambda^4} k_3 a_{16} & \left[\begin{array}{l} \cosh \lambda(x-a) + \frac{1}{2} \left(1 - \frac{1}{32} i \right) \cosh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \cosh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \\ 0\left(\frac{1}{\lambda^9} e^{|t|(x-a)}\right) \end{array} \right] \end{aligned}$$

If $a_{12} = 0, a_{11} = 0, a_{14} = 0, a_{13} = 0, a_{16} = 0$ and $a_{15} \neq 0$:

$$\begin{aligned} \psi(x) = -\frac{1}{3\lambda^5} k_3 a_{15} & \left[\begin{array}{l} \sinh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \\ 0\left(\frac{1}{\lambda^{10}} e^{|t|(x-a)}\right) \end{array} \right] \end{aligned}$$

If $a_{12} = 0$ and $a_{11} \neq 0$:

$$\begin{aligned} \psi(x) = -\frac{1}{3\lambda} k_1 a_{11} & \left[\begin{array}{l} \sinh \lambda(x-a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] - 0\left(\frac{1}{\lambda^2}\right) \end{aligned}$$

Then :

$$\begin{aligned} k_1^{-1} \psi(x) = & -\frac{1}{3\lambda} a_{11} \\ & \left[\begin{array}{l} \sinh \lambda(x-a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)(x-a) \end{array} \right] + 0\left(\frac{1}{\lambda^2}\right) \end{aligned}$$

$$\begin{aligned}
 k_1^2 = k_1^2 \int_a^b \psi^2(x) dx &= \frac{1}{9} k_1^{-2} a_{11}^2 \int_a^b \frac{1}{\lambda^2} \left[\sinh \lambda(x-a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right]^2 dx + 0 \left(\frac{1}{\lambda^2} \right) = \\
 &\left(\begin{array}{l} \frac{\sin(\lambda(a-b)i) 1025i}{3071\lambda^3} + \frac{\sin(\lambda(a-b)2i) 1024i}{3071\lambda^3} + \frac{(a-b)}{\lambda^2} + \frac{1025 3^2 \sin \left(\frac{1}{3^2 \lambda(a-b)} \right)}{9213\lambda^3} + \\ \frac{1}{\lambda^3} \sin \left(\frac{1}{3^2 \lambda(a-b)} \right) \cos \left(\frac{\lambda(a-b)i}{2} \right) \left(\frac{2048 3^2}{3071} + \frac{64}{3071} \right) + \frac{1}{\lambda^3} \cos \left(\frac{1}{3^2 \lambda(a-b)} \right) \\ \sin \left(\frac{\lambda(a-b)i}{2} \right) \left(\frac{1}{3^2 64i} - \frac{2048i}{3071} \right) - \frac{1}{\lambda^3} \sin \left(\frac{1}{3^2 \lambda(a-b)} \right) \cos \left(\frac{\lambda(a-b)3i}{2} \right) \\ \left(\frac{2048 3^2}{9213} - \frac{64}{3071} \right) + \frac{1}{\lambda^3} \cos \left(\frac{1}{3^2 \lambda(a-b)} \right) \sin \left(\frac{\lambda(a-b)3i}{2} \right) \left(\frac{1}{9213} + \frac{2048i}{3071} \right) - \\ \frac{1}{\lambda^3} \sin \left(\frac{1}{3^2 \lambda(a-b)} \right) \cos \left(\lambda(a-b)i \right) \left(\frac{1023 3^2}{12284} - \frac{64}{3071} \right) + \frac{1}{\lambda^3} \cos \left(\frac{1}{3^2 \lambda(a-b)} \right) \\ \sin \left(\lambda(a-b)i \right) \left(\frac{1}{3^2 16i} + \frac{1023i}{12284} \right) \end{array} \right) + 0 \left(\frac{1}{\lambda^2} \right) = \\
 &= \frac{1}{9} k_1^2 a_{11}^2 \left(\begin{array}{l} \frac{3071}{4096} \left(0 \left(\frac{1}{\lambda^3} \right) + 0 \left(\frac{1}{\lambda^3} \right) + \frac{(a-b)}{\lambda^2} + 0 \left(\frac{1}{\lambda^3} \right) + 0 \left(\frac{1}{\lambda^3} \right) - \right) + 0 \left(\frac{1}{\lambda^2} \right) = \\ \frac{3071}{4096} \left(0 \left(\frac{1}{\lambda^3} \right) + 0 \left(\frac{1}{\lambda^3} \right) - 0 \left(\frac{1}{\lambda^3} \right) + 0 \left(\frac{1}{\lambda^3} \right) + 0 \left(\frac{1}{\lambda^3} \right) \right) + 0 \left(\frac{1}{\lambda^2} \right) = \end{array} \right) + 0 \left(\frac{1}{\lambda^2} \right) = \\
 &= \frac{1}{9} k_1^2 a_{11}^2 \frac{3071(a-b)}{4096 \lambda^2} + 0 \left(\frac{1}{\lambda^3} \right) + 0 \left(\frac{1}{\lambda^2} \right)
 \end{aligned}$$

$$k_1^2 = \frac{1}{9} k_1^2 a_{11}^2 \frac{3071(a-b)}{4096 \lambda^2} + 0 \left(\frac{1}{\lambda^2} \right)$$

$$\psi(x) = \mp \frac{1}{3} \left(\frac{36864}{3071(b-a)} \right)^2$$

Then:

$$\begin{aligned}
 a_{11} &= \pm \lambda \left(\frac{36864}{3071(b-a)} \right)^2 \\
 \psi(x) &= \mp \frac{1}{3\lambda} k_1 \lambda \left(\frac{36864}{3071(b-a)} \right)^2 \\
 &\left[\sinh \lambda(x-a) + \frac{1}{2} \left(1 + \frac{1}{32} i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right] + \\
 &\left[(x-a) + \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right] + \\
 &0 \left(\frac{1}{\lambda^2} \right) \\
 \psi(x) &= -\frac{1}{3\lambda^5} k_3 a_{15} \\
 &\left[\sinh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32} i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right] + 0 \left(\frac{1}{\lambda^{10}} \right) \\
 &\left[(x-a) - \frac{1}{2} \left(1 + \frac{1}{32} i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right]
 \end{aligned}$$

Then:

Then:

$$k_1^{-2} = \frac{1}{9} \frac{3071}{4096} a_{15}^2 \frac{1}{\lambda^{10}} (a-b) + O\left(\frac{1}{\lambda^{10}}\right)$$

$$\text{If } k_3 = 1 \text{ then } a_{15} = \pm \lambda^5 \left(\frac{36864}{3071(b-a)} \right)^2$$

$$\psi(x) = \mp \frac{1}{3} \left(\frac{36864}{3071(b-a)} \right)^2$$

$$\left[\sinh \lambda(x-a) - \frac{1}{2} \left(-1 + \frac{1}{32}i \right) \sinh \lambda \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \right]$$

$$+ \left[(x-a) - \frac{1}{2} \left(1 + \frac{1}{32}i \right) \sinh \lambda \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) (x-a) \right]$$

$$O\left(\frac{1}{\lambda^{10}}\right)$$

Theorem 3: Asymptotic formula for eigen values λ_n as $n \rightarrow \infty$ is:

$$\lambda = \frac{i\theta + 2n\pi i}{b-a}$$

Proof: Since:

$$\Delta(\lambda) = \begin{vmatrix} U_0(y_0) & U_0(y_1) & U_0(y_2) & U_0(y_3) & U_0(y_4) & U_0(y_5) \\ U_1(y_0) & U_1(y_1) & U_1(y_2) & U_1(y_3) & U_1(y_4) & U_1(y_5) \\ U_2(y_0) & U_2(y_1) & U_2(y_2) & U_2(y_3) & U_2(y_4) & U_2(y_5) \\ U_3(y_0) & U_3(y_1) & U_3(y_2) & U_3(y_3) & U_3(y_4) & U_3(y_5) \\ U_4(y_0) & U_4(y_1) & U_4(y_2) & U_4(y_3) & U_4(y_4) & U_4(y_5) \\ U_5(y_0) & U_5(y_1) & U_5(y_2) & U_5(y_3) & U_5(y_4) & U_5(y_5) \end{vmatrix}$$

Then, we can see that:

$$\Delta(\lambda) = \begin{vmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} & a_{26} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} & a_{36} \\ U_3(y_0) & U_3(y_1) & U_3(y_2)U_3(y_3) & U_3(y_4) & U_3(y_5) \\ U_4(y_0) & U_4(y_1) & U_4(y_2)U_4(y_3) & U_4(y_4) & U_4(y_5) \\ U_5(y_0) & U_5(y_1) & U_5(y_2)U_5(y_3) & U_5(y_4) & U_5(y_5) \end{vmatrix}$$

If we suppose that $X = \begin{vmatrix} a_{13} & a_{15} & a_{16} \\ a_{23} & a_{25} & a_{26} \\ a_{33} & a_{35} & a_{36} \end{vmatrix} \neq 0$ and since,

$$U_4(y_0) = -\frac{1}{3}\lambda^5 [3\cosh(\lambda(a-b)) + 3\sinh(\lambda(a-b))] + 0\left(e^{|t|(x-a)}\right)$$

the boundary conditions are linearly independent then:

$$\Delta(\lambda) = X \begin{vmatrix} U_3(y_0) & U_3(y_1) & U_3(y_3) \\ U_4(y_0) & U_4(y_1) & U_4(y_3) \\ U_5(y_0) & U_5(y_1) & U_5(y_3) \end{vmatrix}$$

Then we calculate $U_j(y_i)$ for $j = 3, 4, 5$ and $i = 0, \dots, 5$ by using corollary 2-7:

$$U_3(y_0) = \frac{1}{3}\lambda^5 [3\cosh(\lambda(a-b)) - 3\sinh(\lambda(a-b))] + 0\left(e^{|t|(x-a)}\right)$$

$$U_3(y_1) = \frac{1}{3}\lambda^4 [3\cosh(\lambda(a-x)) - 3\sinh(\lambda(a-x))] + 0\left(\frac{1}{\lambda}e^{|t|(x-a)}\right)$$

$$U_3(y_2) = \frac{1}{3}\lambda^3$$

$$\begin{bmatrix} 3\cosh(\lambda(a-x)) - 3\sinh(\lambda(a-x)) + \sin\left(\frac{\frac{1}{32}\lambda(a-x)}{2}\right) \\ \cosh\left(\frac{\lambda(a-x)}{2}\right)i + \frac{1}{4}\cosh\left(\frac{\lambda\left(\frac{1}{32}i-1\right)(a-x)}{2}\right)\left(\frac{1}{32}i-1\right) + \\ \frac{1}{4}\cosh\left(\frac{\lambda\left(\frac{1}{32}i+1\right)(a-x)}{2}\right)\left(\frac{1}{32}i-1\right) \\ 0\left(\frac{1}{\lambda^2}e^{|t|(x-a)}\right) \end{bmatrix}$$

$$U_3(y_3) = \frac{1}{3}\lambda^2 [3\cosh(\lambda(a-x)) - 3\sinh(\lambda(a-x))] + 0\left(\frac{1}{\lambda^3}e^{|t|(x-a)}\right)$$

$$U_3(y_4) = \frac{1}{3}\lambda [3\cosh(\lambda(a-x)) - 3\sinh(\lambda(a-x))] + 0\left(\frac{1}{\lambda^4}e^{|t|(x-a)}\right)$$

$$U_3(y_5) = \frac{1}{3} [3\cosh(\lambda(a-x)) - 3\sinh(\lambda(a-x))] + 0\left(\frac{1}{\lambda^5}e^{|t|(x-a)}\right)$$

$$U_4(y_0) = -\frac{1}{3}\lambda^5 [3\cosh(\lambda(a-b)) + 3\sinh(\lambda(a-b))] + 0\left(e^{|t|(x-a)}\right)$$

$$U_4(y_1) = -\frac{1}{3}\lambda^4 [-3\cosh(\lambda(a-x)) - 3\sinh(\lambda(a-x))] + 0\left(\frac{1}{\lambda}e^{|t|(x-a)}\right)$$

$$U_4(y_2) = -\frac{1}{3}\lambda^3$$

$$\begin{bmatrix} 3\cosh(\lambda(a-x)) + 3\sinh(\lambda(a-x)) - \sin\left(\frac{\frac{1}{32}\lambda(a-x)}{2}\right) \\ \cosh\left(\frac{\lambda(a-x)}{2}\right)i + \frac{1}{4}\cosh\left(\frac{\lambda\left(\frac{1}{32}i-1\right)(a-x)}{2}\right)\left(\frac{1}{32}i-1\right) + \\ \frac{1}{4}\cosh\left(\frac{\lambda\left(\frac{1}{32}i+1\right)(a-x)}{2}\right)\left(\frac{1}{32}i-1\right) \\ 0\left(\frac{1}{\lambda^2}e^{|t|(x-a)}\right) \end{bmatrix}$$

$$U_4(y_3) = -\frac{1}{3}\lambda^2 [-3\cosh(\lambda(a-x)) - 3\sinh(\lambda(a-x))] + 0\left(\frac{1}{\lambda^3}e^{|t|(x-a)}\right)$$

$$U_4(y_4) = -\frac{1}{3}\lambda [-3\cosh(\lambda(a-x)) - 3\sinh(\lambda(a-x))] + 0\left(\frac{1}{\lambda^4}e^{|t|(x-a)}\right)$$

$$U_4(y_5) = -\frac{1}{3}\lambda [-3\cosh(\lambda(a-x)) - 3\sinh(\lambda(a-x))] + 0\left(\frac{1}{\lambda^5}e^{|t|(x-a)}\right)$$

$$U_5(y_0) = \frac{1}{3}\lambda^5$$

$$\begin{bmatrix} 3\cosh\left(\frac{(\lambda(\sqrt{3}i-1))(a-b)}{2}\right) + 3\cosh\left(\frac{(\lambda(\sqrt{3}i+1))(a-b)}{2}\right) \\ 3\sinh\left(\frac{(\lambda(\sqrt{3}i-1))(a-b)}{2}\right) - 3\sinh\left(\frac{(\lambda(\sqrt{3}i+1))(a-b)}{2}\right) \\ 0\left(e^{|t|(x-a)}\right) \end{bmatrix}$$

$$U_5(y_1) = \frac{1}{3}\lambda^4 \left[\cosh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right)\left(\frac{3}{2}+\frac{3i}{64}\right) + \frac{1}{4} \cosh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right)\left(\frac{3}{2}+\frac{3i}{64}\right) + \sinh \right. \\ \left. \cosh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right)\left(\frac{3}{2}-\frac{3i}{64}\right) + \sinh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right)\left(\frac{3}{2}-\frac{3i}{64}\right) + \right. \\ \left. 0\left(\frac{1}{\lambda}e^{|t|(x-a)}\right) \right]$$

$$U_5(y_2) = \frac{1}{3}\lambda^3 \frac{1}{4} \\ \left[-\cosh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right)\left(\frac{1}{32i}-5\right) - \cosh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right)\left(\frac{1}{32i}-5\right) + \sinh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right) \right. \\ \left. + 0\left(\frac{1}{\lambda^2}e^{|t|(x-a)}\right) \right] \\ \left[\left(\frac{1}{32i}-5\right) + \sinh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right)\left(\frac{1}{32i}-5\right) \right]$$

$$U_5(y_3) = \frac{1}{3}\lambda^2 \\ \left[3\cosh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right) - 3\cosh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right) - 3\sinh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right) + \sinh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right) \right] + \\ 0\left(\frac{1}{\lambda^2}e^{|t|(x-a)}\right)$$

$$U_5(y_4) = \frac{1}{3}\lambda \\ \left[\cosh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right)\left(\frac{3}{2}+\frac{3i}{64}\right) + \cosh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right)\left(\frac{3}{2}+\frac{3i}{64}\right) + \sinh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right) \right. \\ \left. + 0\left(\frac{1}{\lambda^4}e^{|t|(x-a)}\right) \right] \\ \left[\left(\frac{3}{2}-\frac{3i}{64}\right) + \sinh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right)\left(-\frac{3}{2}+\frac{3i}{64}\right) \right]$$

$$U_5(y_5) = \frac{1}{3} \\ \left[\cosh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right)\left(-\frac{3}{2}-\frac{3i}{64}\right) + \cosh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right)\left(\frac{3}{2}-\frac{3i}{64}\right) + \sinh\left(\frac{\lambda\left(\frac{1}{32i}-1\right)(a-x)}{2}\right) \right. \\ \left. + 0\left(\frac{1}{\lambda^5}e^{|t|(x-a)}\right) \right] \\ \left[\left(\frac{3}{2}+\frac{3i}{64}\right) + \sinh\left(\frac{\lambda\left(\frac{1}{32i}+1\right)(a-x)}{2}\right)\left(-\frac{3}{2}+\frac{3i}{64}\right) \right]$$

$$\Delta(\lambda) = X$$

$$\left(\frac{\lambda^{11}}{27} \left[\begin{array}{l} \cosh \frac{(\lambda(\sqrt{3}i-1)(a-x))}{2} (96-i) + \cosh \frac{(\lambda(\sqrt{3}i+1)(a-x))}{2} (-96-i) + \sinh \frac{(\lambda(\sqrt{3}i-1)(a-x))}{2} (-96+i) \\ \sinh \frac{(\lambda(\sqrt{3}i+1)(a-x))}{2} (96+i) \left(\begin{array}{l} \cosh \lambda(a-x) \\ + \sinh \lambda(a-x) \end{array} \right) (\cosh \lambda(a-x) - \sinh \lambda(a-x)) \end{array} \right] \right)$$

$$\frac{1}{32} + 0\left(\lambda^6 e^{|t|(x-a)}\right) + 0\left(\lambda e^{|t|(x-a)}\right) + 0\left(\frac{1}{\lambda^2} e^{|t|(x-a)}\right)$$

$$\Delta(\lambda) = \frac{27}{32} X$$

$$\left(\lambda^{11} \left[\begin{array}{l} \cosh \frac{(\lambda(\sqrt{3}i-1)(a-x))}{2} (96-i) + \cosh \frac{(\lambda(\sqrt{3}i+1)(a-x))}{2} (-96-i) + \sinh \frac{(\lambda(\sqrt{3}i-1)(a-x))}{2} (-96+i) + \sinh \frac{(\lambda(\sqrt{3}i+1)(a-x))}{2} (96+i) \\ 0\left(\lambda^6 e^{|t|(x-a)}\right) + 0\left(\lambda e^{|t|(x-a)}\right) + 0\left(\frac{1}{\lambda^2} e^{|t|(x-a)}\right) \end{array} \right] \right) +$$

$$\Delta(\lambda) = \frac{27}{32} X e^{\left(\frac{-\sqrt{3}\lambda(a-b)i}{2} \right)} e^{\left(\frac{\lambda(a-b)}{2} \right)} \lambda^{11}$$

$$\left[\left(\left(e^{(\lambda(a-b))} (96-i) - 96+i \right) \right) + 0\left(\frac{1}{\lambda^5} e^{|t|(x-a)}\right) + 0\left(\frac{1}{\lambda^{10}} e^{|t|(x-a)}\right) + 0\left(\frac{1}{\lambda^{12}} e^{|t|(x-a)}\right) \right]$$

$$\Delta(\lambda) = \frac{27}{32} X e^{\left(\frac{-\sqrt{3}\lambda(a-b)i}{2} \right)} e^{\left(\frac{\lambda(a-b)}{2} \right)} \lambda^{11} \left[\left(\left(e^{(\lambda(a-b))} (96-i) - 96+i \right) \right) + 0\left(\frac{1}{\lambda^5} e^{3|t|(b-a)}\right) \right]$$

If $\Delta(\lambda) = 0$ then:

$$\left(e^{(\lambda(a-b))} (96-i) - 96+i + 0\left(\frac{1}{\lambda^5} e^{3|t|(b-a)}\right) \right) = 0$$

$$\text{Then } e^{(\lambda(a-b))} (96-i) - 96+i = 0\left(\frac{1}{\lambda^5} e^{3|t|(b-a)}\right) \text{ for large value of } |\lambda|$$

value of $|\lambda|$ we get:

$$e^{(\lambda(a-b))} = \frac{96+i}{96-i}$$

If $\theta = \tan^{-1}(96+i/96-i)$ then:

$$\lambda = \frac{1}{b-a} \left\{ \ln \left| \frac{96+i}{96-i} \right| + i\theta + 2n\pi i \right\}, \text{ for } n = 0, \pm 1, \pm 2, \dots$$

$$\lambda = \frac{1}{b-a} \{ \ln 1 + i\theta + 2n\pi i \}, \text{ for } n = 0, \pm 1, \pm 2, \dots$$

$$\lambda = \frac{i\theta + 2n\pi i}{b-a}, \text{ for } n = 0, \pm 1, \pm 2, \dots$$

In mathematics, spectral theory is an inclusive term for theories that extend a single square matrix's eigenvector and eigenvalue theory to a much broader theory of operator structure in a variety of mathematical spaces.

CONCLUSION

Aigunov, G.A., 1996. On the boundedness of orthonormal eigenfunctions of a class of Sturm-Liouville operators with a weight function of unbounded variation on a finite interval. Russ. Math. Surv., Vol. 51, No. 2.

Jwamer, K.H. and G.A. Aigounv, 2010. About uniform limitation of normalized eigenfunctions of t. regge problem in the case of weight functions, satisfying to lipschitz condition. J. Gen. Math. Notes, 1: 115-129.

Jwamer, K.H. and R.Q. Rasul, 2017. Estimations of the upper bound for the eigen-functions of the fourth order boundary value problem with smooth coefficients. Math. Sci. Lett., 6: 67-74.

- Menken, H., 2010. Accurate asymptotic formulas for eigenvalues and eigenfunctions of a boundary-value problem of fourth order. *Boundary Value Prob.*, Vol. 2010, No. 1, 10.1155/2010/720235
- Moller, M. and B. Zinsou, 2012. Spectral asymptotics of self-adjoint fourth order boundary value problems with eigenvalue parameter dependent boundary conditions. *Boundary Value Prob.*, Vol. 2012, No. 1. 10.1186/1687-2770-2012-106
- Naimark, M.A., 1967. *Linear Differential Operators*. Part 1. Frederick Ungar Publishing Company, New York, USA., Pages: 144.
- Tamarkin, J., 1928. Some general problems of the theory of ordinary linear differential equations and expansion of an arbitrary function in series of fundamental functions. *Math. Z.*, 27: 1-54.