

## New Types of Openness and Closed Graphs in Topological Space

Ali Khalaf Hussain Al-Hachami

Department of Mathematics, College of Education, Wassit University, Kut, Iraq

**Abstract:**  $\alpha$ -unequivocally  $\theta$ -continuity function and  $(\alpha, \theta)$ -closed graphs was examined by Chae *et al.* The goal of this study is to research a few of new portrayal and properties of  $\alpha$ -unequivocally  $\theta$ -continuity and  $(\alpha, \theta)$ -closed graphs. Besides we characterize new sort of a function called  $\alpha, \theta$ -open function which is more grounded than quasi  $\alpha$ -open and  $\alpha$ -open and we acquire a few portrayals and properties for it.

**Key words:** Characterize, properties, portrayals, function,  $\theta$ -continuity, ground

### INTRODUCTION

The concept of  $\alpha$ -open sets was introduced and investigated by Njastad (1965). Latterly, the concept of  $\alpha$ -unequivocally  $\theta$ -continuity function has studied by Chae *et al.* (1995). We know from Chae *et al.* (1995) that the type of  $\alpha$ -unequivocally  $\theta$ -continuity function is stronger than a unequivocally  $\theta$ -continuity function (Noiri, 1980) and a unequivocally  $\alpha$ -continuous function (Faro, 1987).

In this study we aim to investigate further properties and characterizations of  $\alpha$ -unequivocally  $\theta$ -continuity functions as well as  $\theta$ -closed graph (Chae *et al.*, 1995) and new types of function define called  $\alpha, \theta$ -open functions which is stronger than quasi  $\alpha$ -open and hence, unequivocally  $\alpha$ -open, some characterizations and properties are obtain for it.

**Preliminaries:** All through this study just  $X$  speaks to a topological space.

**Definition 2.1:** Let be an subset of a topological space  $(X, \tau)$  then is called:

- Regular open if  $A = (\bar{A})^\circ$  (Njastad, 1965)
- $\alpha$ -pen if  $A \subseteq (\bar{A}^\circ)^\circ$  (Levine, 1963)
- Semi-open if  $A \subseteq \overline{A^\circ}$  (Levine, 1963)
- $\theta$ -open if for each  $x \in A$ , there exist an open set  $U$  in  $X$  such that  $x \in U \subset \bar{U} \subset A$  (Velicko, 1968)
- $\theta$ -semi-open if for each  $x \in A$ , there exist an semi-open set  $U$  in  $X$  such that  $x \in U \subset \bar{U} \subset A$  (Noiri and Kang, 1984)

The supplements of the sets said above are their individual closed sets.

**Definition 2.2:** The set  $\alpha\bar{A} = \{p \in X: A \cap H \neq \emptyset \text{ for each } \alpha\text{-open set } H \text{ containing } p\}$ .

**Definition 2.3:** “A filter base  $\Psi$  is said to be  $\theta$ -convergent (Velicko, 1968) (resp.  $\alpha$ -convergent to a point  $x \in X$  if for each open (resp.  $\alpha$ -open) set  $G$  containing  $x$ , there exist an  $F \in \Psi$  such that  $F \subset \bar{G}$  (resp”.  $F \subset G$ ).

**Definition 2.4; (Maheshwari *et al.*, 1982):** “A subset  $A$  of a topological space  $(X, \tau)$  is called a feebly open set in  $X$  if there exist an open set  $U$  such that  $U \subset A \subset sCl(U)$  where is the semi-closure operator”.

**Remark 2.5; (Jankovic, 1985):** A subset  $A$  of a topological space  $(X, \tau)$  is called  $\alpha$ -open if and only if it is feebly open. It is notable that for a space  $(X, \tau)$ ,  $X$  can be retopologized by the family  $\tau^\alpha$  of all  $\alpha$ -open sets of  $X$  (Maheshwari *et al.*, 1982; Thakur, 1980) and furthermore the family  $\tau^\theta$  of all  $\theta$ -open set of  $X$  (Velicko, 1968) that is  $\tau^\theta$  (called  $\theta$ -topology) and  $\tau^\alpha$  (called an  $\alpha$ -topology) are topologies on  $X$  and it is clearly that  $\tau^\theta \subset \tau \subset \tau^\alpha$ . The family of all  $\alpha$ -open (resp.  $\theta$ -open and feebly-open) arrangements of  $X$  is indicated by  $\alpha O(X)$  (resp.  $\theta O(X)$  and  $FO(X)$ ).

**Definition 2.5; (Noiri and Kang, 1984):** A function  $f: X \rightarrow Y$  is said to be unequivocally  $\theta$ -continuous if for each  $x \in X$  and each open set  $H$  of  $Y$  containing  $f(x)$ , there exist an open set  $G$  of  $X$  containing  $x$  such that  $f(\bar{G}) \subset H$ .

**Definition 2.6; (Noiri and Kang, 1984):** A function  $f: X \rightarrow Y$  is said to be unequivocally  $\theta$ -continuous if for each open set  $H$  of  $Y$ ,  $f^{-1}(H)$  is  $\theta$ -open  $X$  in if and only if each closed set  $F$  of  $Y$   $f^{-1}(F)$ , is  $\theta$ -closed in  $X$ .

**Definition 2.7; (Maheshwari *et al.*, 1983):** A function  $f: X \rightarrow Y$  is said to be unequivocally  $\alpha$ -continuous (resp. faintly continuous (Long and Herrington, 1982), completely  $\alpha$ -irresolute and unequivocally  $\alpha$ -irresolute (Faro, 1987) if for each open (resp.  $\theta$ -open,  $\alpha$ -open and  $\alpha$ -open) set  $H$  of  $Y$ ,  $f^{-1}(H)$  is  $\alpha$ -open (resp. open, regular open and open) in  $X$ .

**Definition 2.8; (Noiri, 1973):** A function  $f: X \rightarrow Y$  is said to be semi-open (resp.  $\alpha$ -open (Maheshwari *et al.*, 1983), quasi  $\alpha$ -open (Thivagar, 1991; Abdul Jabbar, 2000),  $\theta$ s-open (Abdul-Jabbar, 2000) weakly  $\theta$ s-open and  $s^{**}$ -open (Ali, 2003) function if the image of each open, (resp. open  $\alpha$ -open, open,  $\theta$ -open and semi-open) set of  $G$  of  $X$ ,  $f(G)$  is semi-open (resp.  $\alpha$ -open, open,  $\theta$ -semi-open,  $\theta$ -semi-open and open) in  $Y$ .

**Definition 2.9; (Lee *et al.*, 1985):** A function  $f: X \rightarrow Y$  is said to be pre-feebly-open (resp. unequivocally  $\alpha$ -open (Thivagar, 1991),  $\alpha^{**}$ -open (Ali, 2003) function if the image of each  $\alpha$ -open set of  $G$  of  $X$ ,  $f(G)$  is  $\alpha$ -open in  $Y$ .

**Definition 2.10; (Baker, 1986):** Let  $A$  be a subset of a topological space  $(X, \tau)$  then  $A$  is called  $\theta$ -neighborhood of a point  $x$  in  $X$  if there exist an open set  $U$  such that  $x \in U \subset \bar{U} \subset A$ .

**Definition 2.11; (Lee *et al.*, 1985):** "A function  $f: X \rightarrow Y$  is said to be"  $\theta$ -open function if for each  $x \in X$  and each  $\theta$ -neighborhood  $A$  of  $X$ ,  $F(A)$  is  $\theta$ -neighborhood  $f(x)$ .

**Definition 2.12; (Singal and Arya, 1969):** A space  $X$  is said to be practically regular if for each regular closed set of  $X$  and each point  $x \in R$ , there exist disjoint open set  $U$  and  $V$  such that  $R \subset U$  and  $x \in V$ .

**Definition 2.13; (Faro, 1987):** A space  $X$  is said to be  $\alpha$ -hausdorff if for any  $x, y \in X$ ,  $x \neq y$ , there exist  $\alpha$ -open set  $G$  and  $H$  such that  $x \in G$ ,  $y \in H$  and  $G \cap H = \emptyset$ .

**Definition 2.14:** "A space  $X$  is said to be  $\theta$ -compact (resp.  $\alpha$ -compact (Jankovic *et al.*, 1988) if and only if every cover of  $X$  by  $\theta$ -open (resp.  $\alpha$ -open) sets has a finite subcover".

**Definition 2.15; (Porter and Thomas, 1969):** "A subset  $A$  of a topological space  $(X, \tau)$  is said to be quasi  $H$ -closed relative to  $X$  if  $\{E_i: i \in I\}$  each cover of  $A$  by open sets of  $X$ , there exist a finite subset  $I_0$  of  $I$  such that  $A \subset \bigcup \{E_i: i \in I_0\}$ ".

**Definition 2.16; (Porter and Thomas, 1969):** "A space  $X$  is said to be quasi  $H$ -closed if  $X$  is quasi  $H$ -closed relative to  $X$ ".

**Definition 2.17; (Noiri, 1975):** A function  $f: X \rightarrow Y$  is said to be  $\theta$ -closed (resp.  $s^{**}$ -closed (Long and Herring, 1977), semi-closed (Dube *et al.*, 1998),  $\theta$ s-closed (Abdul-Jabbar, 2000), almost unequivocally  $\theta$ s-closed and unequivocally  $\theta$ s-closed graph if and only if for  $x \in X$  each and each  $y \in Y$

such that  $y \neq f(x)$ , there exist an open (resp. semi-open, semi-open, semi-open, semi-open and semi-open)  $U$  containing  $x$  in  $X$  and an open (resp. open, semi-open, open, open and open) set  $V$  containing  $f(x)$  in  $Y$  such that  $(\bar{U} \times \bar{V}) \cap G(f) = \emptyset$  {resp.  $(U \times V) \cap G(f) = \emptyset$ ,  $(U \times V) \cap G(f) = \emptyset$ ,  $(\bar{U} \times \bar{V}) \cap G(f) = \emptyset$ ,  $(\bar{U} \times \bar{V}) \cap G(f) = \emptyset$  and  $(\bar{U} \times \bar{V}) \cap G(f) = \emptyset$  }.

## MATERIALS AND METHODS

### $\alpha$ -Unequivocally $\theta$ -coherence

**Definition 3.1:** By Chae *et al.* (1995) "A function  $f: X \rightarrow Y$  is said to be  $\alpha$ -unequivocally  $\theta$ -coherence if for each  $x \in X$  and each  $\alpha$ -open set  $H$  of  $Y$  containing  $f(x)$ , there exist an open set  $U$  of  $X$  containing  $x$  with the end goal that  $f(\bar{U}) \subset H$ ".

**Theorem 3.1:** For a function  $f: (X, \tau) \rightarrow (Y, \gamma)$  the accompanying proclamations are proportionality:

- $f$  is  $\alpha$ -unequivocally  $\theta$ -coherence
- $f: (X, \tau) \rightarrow (Y, \gamma)$  is unequivocally  $\alpha$ -irresolute

**Theorem 3.2:** In the event that a function  $f: X \rightarrow Y$   $\alpha$ -unequivocally  $\theta$ -coherence at that point for each  $x \in X$  and each  $\alpha$ -open set  $H$  of  $Y$  containing  $f(x)$ , there exist  $\theta$ -open set  $N$  of  $X$  containing  $x$  with the end goal that  $f(N) \subset H$ . The evidence of the above theorems are not hard and along these lines, they are precluded.

**Theorem 3.3:** For a function  $f: (X, \tau) \rightarrow (Y, \gamma)$  the accompanying articulations are proportionality:

- $f$  is  $\alpha$ -unequivocally  $\theta$ -coherence
- For each point  $x \in X$  and each filter base  $\Psi$  in  $X$   $\theta$ -converging to  $x$ , the filterbase  $f(\Psi)$  converges to  $f(x)$  in  $(Y, \alpha_0(Y))$
- For each point  $x \in X$  and each net  $\{x_\lambda\}_{\lambda \in \nabla}$  in  $X$   $\theta$ -converging to  $x$ , the net  $\{f(x_\lambda)\}_{\lambda \in \nabla}$  converges to  $f(x)$  in  $(Y, \alpha_0(Y))$
- For each point  $x \in X$  and each filterbase  $\Psi$  in  $X$   $\theta$ -converging to  $x$ , the filterbase  $f(\Psi)$   $\alpha$ -converges to  $f(x)$  in  $(Y, \gamma)$
- For each point  $x \in X$  and each net  $\{x_\lambda\}_{\lambda \in \nabla}$  in  $X$   $\theta$ -converging to  $x$ , the net  $\{f(x_\lambda)\}_{\lambda \in \nabla}$   $\alpha$ -converges to  $f(x)$  in  $(Y, \gamma)$

**Proof:** (i) $\Rightarrow$ (ii) $\Rightarrow$ (iii) and (i) $\Rightarrow$ (iv) $\Rightarrow$ (v) follows, immediately from Definition 3.1 and Theorem 2 of (Chae *et al.*, 1995).

**Lemma 3.1; (Andrijevic, 1984):** Let  $X$  be a topological space and  $A \subset X$ . At that point the accompanying are hold:

- $\alpha Cl(E) = E \cup Cl(Int(Cl(E)))$
- $\alpha Int(E) = E \cup Int(Cl(Int(E)))$

**Theorem 3.4:** For a function  $f: X \rightarrow Y$  the accompanying articulations are comparability:

- $f$  is  $\alpha$ -unequivocally  $\theta$ -coherence
- $f(Cl_\theta(A)) \subset Cl(Int(Cl(f(A))))$ , for every subset an of  $X$
- $Cl_\theta(f^{-1}(E)) \subset f^{-1}(Cl(Int(Cl(f(E))))$ , for every subset an of  $E$  of  $Y$
- $f^{-1}(Cl(Int(Cl(f(E)))) \subset Int_\theta(f^{-1}(E))$ , for every subset an of  $E$  of  $Y$

**Proof:** This follows from Lemma 3.1 and Theorem 2 of (Chae *et al.*, 1995).

**Theorem 3.5:** If a function  $f: X \rightarrow Y$  is  $\alpha$ -unequivocally  $\theta$ -coherence and if  $E$  is an open subset of  $X$ , then  $f|_E: E \rightarrow Y$  is  $\alpha$ -unequivocally  $\theta$ -coherence in the subspace  $E$ .

**Proof:** Let  $H$  be any  $\alpha$ -open subset of  $Y$ . Since,  $f$  is  $\alpha$ -unequivocally  $\theta$ -coherence. Therefore, by [7, theorem 2],  $f^{-1}(H) \in \theta_0(X)$ , so by Lemma 1.2.9 of (Abdul-Jabbar, 2000)  $(f|_E)^{-1}(H) = f^{-1}(H) \cap E \in \theta_0(E)$ . This implies that  $f|_E: E \rightarrow Y$  is  $\alpha$ -unequivocally  $\theta$ -coherence.

**Theorem 3.6:** For any two functions,  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$ , the accompanying are valid:

- $f$  is  $\alpha$ -unequivocally  $\theta$ -coherence and  $g$  is  $\alpha$ -continuous, then  $g \circ f$  is unequivocally  $\theta$ -coherence
- $f$  is faintly continuous and  $g$  is  $\alpha$ -unequivocally  $\theta$ -coherence, then  $g \circ f$  is unequivocally  $\alpha$ -irresolute

**Theorem 3.7:** “Let  $f: X \rightarrow Y$  be faintly continuous and  $\theta$ -open function and  $g: Y \rightarrow Z$  be a function. Then  $g \circ f: X \rightarrow Z$   $\alpha$ -unequivocally  $\theta$ -continuous if and only if  $g$  is  $\alpha$ -unequivocally  $\theta$ -coherence”.

**Proof:** Let  $g \circ f$   $\alpha$ -unequivocally  $\theta$ -coherence and  $H \in \alpha_0(Z)$ . Then  $(g \circ f)^{-1}(H) = f^{-1}(g^{-1}(H)) \in \theta_0(X)$ . Since,  $f$  is  $\theta$ -open function,  $f(f^{-1}(g^{-1}(H))) \in \theta_0(Y)$ . Hence,  $g^{-1}(H) \in \theta_0(Y)$ . Thus,  $g$  is  $\alpha$ -unequivocally  $\theta$ -coherence. It is easy to prove the opposite and is thus omitted.

**Theorem 3.8:** If  $g: Y \rightarrow Z$  be a one to one  $\alpha$ -open function on  $Y$  onto  $Z$  and  $g \circ f: X \rightarrow Z$  is  $\alpha$ -unequivocally  $\theta$ -continuous. Then  $f$  is unequivocally  $\theta$ -coherence.

**Proof:** Suppose  $g$  is  $\alpha$ -open function. Let  $H$  be an open subset of  $Y$ , since,  $g$  is one to one and onto, then the set  $g(H)$  is an  $\alpha$ -open subset of  $Z$ , since,  $g \circ f$  is  $\alpha$ -unequivocally  $\theta$ -coherence, it follows that  $(g \circ f)^{-1}(g(H)) = f^{-1}(g^{-1}(g(H))) = f^{-1}(H)$  is  $\theta$ -open in  $X$ . Thus,  $f$  is unequivocally  $\theta$ -continuous.

**Theorem 3.9:** If  $X$  is almost regular and  $f: X \rightarrow Y$  is completely  $\alpha$ -irresolute function  $f$  is  $\alpha$ -unequivocally  $\theta$ -coherence.

**Proof:** Let  $H$  be an  $\alpha$ -open subset of  $Y$ , since,  $f$  is completely  $\alpha$ -irresolute function, then  $f^{-1}(H)$  is regular open in  $X$  and from the fact that a space  $X$  is almost regular if and only if fore each  $x \in X$  and each regular open set  $f^{-1}(H)$  containing  $x$ , there exist a regular open set  $O$  such that  $x \in O \subset \bar{O} \subset f^{-1}(H)$  [31, theorem 2.2]. Therefore is  $\theta$ -open in  $X$  and by [7, theorem 2],  $f$  is  $\alpha$ -unequivocally  $\theta$ -continuous.

**Lemma 3.2; (Chae *et al.*, 1986):** Let  $\{X_\lambda: \lambda \in \Delta\}$  be a family of spaces and  $U_{\lambda i}$  be subset of  $X_{\lambda i}$  for each  $i = 1, 2, \dots, n$ . Then  $U = \prod_{i=1}^n U_{\lambda i} \times \prod_{\lambda \notin \Delta} X_\lambda$  is  $\alpha$ -open in  $\prod_{\lambda \in \Delta} X_\lambda$  if and only if  $U_{\lambda i} \in \alpha_0(X_{\lambda i})$  for each  $i = 1, 2, \dots, n$ .

**Theorem 3.10:** Let  $g_\lambda: X_\lambda \rightarrow Y_\lambda$  be a function for each  $\lambda \in \Delta$  and  $g: \prod X_\lambda \rightarrow \prod Y_\lambda$  a function defined by  $g(\{x_\lambda\}) = \{g_\lambda(x_\lambda)\}$  for eac  $\{x_\lambda\} \in \prod X_\lambda$ . If  $g$  is  $\alpha$ -unequivocally  $\theta$ -coherence, then  $g_\lambda$  is  $\alpha$ -unequivocally  $\theta$ -coherence for each  $\lambda \in \Delta$ .

**Proof:** “Let  $\beta \in \Delta$  and  $V_\beta \in \alpha_0(Y_\beta)$ . Then, by Lemma 3.2,  $V = V_\beta \times \prod_{\lambda \neq \beta} Y_\lambda$  is  $\alpha$ -open in  $\prod Y_\lambda$  and  $g^{-1}(V) = g_\beta^{-1}(V_\beta) \times \prod_{\lambda \neq \beta} Y_\lambda$  is  $\alpha$ -open in  $\prod X_\lambda$ . From Lemma 3.2,  $g_\beta^{-1}(V_\beta) \in \theta_0(X_\beta)$ .” Therefore,  $g_\beta$  is  $\alpha$ -unequivocally  $\theta$ -coherence.

**Remark 3.1:** It was known in [6, example 2.2] that  $V \in \alpha_0(X \times Y)$  may not, generally, be a union of sets of the form  $A \times B$  in the product space  $X \times Y$  where  $A \in \alpha_0(X)$  and  $B \in \alpha_0(Y)$ . Therefore, the converse of theorem 3.10 may not be true, generally.

**Theorem 3.11:** Let  $g: X \rightarrow Y_1 \times Y_2$  be  $\alpha$ -unequivocally  $\theta$ -coherence where  $X, Y_1$  and  $Y_2$  are any topological spaces. Let  $f_i: X \rightarrow Y_i$  defined as follows: For  $x \in X$ ,  $g(x) = (x_1, x_2)$ ,  $f_i(x) = x_i$  for  $i = 1, 2$ . Then  $f_i: X \rightarrow Y_i$  is  $\alpha$ -unequivocally  $\theta$ -continuous for  $i = 1, 2$ .

**Proof:** Let  $x$  be any point in  $X$  and  $H_i$  be any  $\alpha$ -open set in  $Y_i$  containing  $f_i(x) = x_i$ , then by Lemma 3.2,  $H_1 \times Y_2$  is  $\alpha$ -open  $Y_1 \times Y_2$  which contain  $(x_1, x_2)$ . Since,  $g$  is  $\alpha$ -unequivocally  $\theta$ -continuous, therefore, there sexist an open set  $U$  containing  $x$  such that  $g(Cl(U)) \subset H_1 \times Y_2$ . Then  $f_1(Cl(U)) \times f_2(Cl(U)) \subset H_1 \times Y_2$ . Therefore,  $f_1(Cl(U)) \subset H_1$ . Hence,  $f_1$   $\alpha$ -unequivocally  $\theta$ -coherence. Similar statement for  $f_2$  is  $\alpha$ -unequivocally  $\theta$ -coherence.

**Lemma 3.3:** Let  $X_1, X_2, \dots, X_n$  be  $n$  topological spaces and  $x = \prod_{i=1}^n x_i$ . Let  $E_i \in \theta_0(X_i)$  for  $i = 1, 2, \dots, n$  then  $\prod_{i=1}^n E_i \in \theta_0(\prod_{i=1}^n X_i)$ .

**Proof:** Let  $(x_1, x_2, \dots, x_n) \in \prod_{i=1}^n E_i$  then  $x_i \in E_i$  for  $i = 1, 2, \dots, n$ . Since,  $E_i \in \theta_0(X_i)$ , for  $i = 1, 2, \dots, n$ . Then there exist open sets  $U_i$  for  $i = 1, 2, \dots, n$  such that  $x_i \in U_i \subset \overline{U_i} \subset E_i$  for  $i = 1, 2, \dots, n$ . Therefore,  $(x_1, x_2, \dots, x_n) \in U_1 \times U_2 \times \dots \times U_n \subset \overline{U_1} \times \overline{U_2} \times \dots \times \overline{U_n} = Cl x_1 \times x_2 \times \dots \times x_n (U_1 \times U_2 \times \dots \times U_n) \subset \prod_{i=1}^n E_i$  and  $\prod_{i=1}^n U_i \in \tau(\prod_{i=1}^n E_i)$  is  $\theta$ -open set in  $\prod_{i=1}^n X_i$ .

**Theorem 3.12:** Let  $X_1, X_2, \dots, X_n$  and  $Z$  be topological spaces and  $\prod_{i=1}^n X_i \rightarrow Z$ . If given any point  $p$  of  $\prod_{i=1}^n X_i$ ,  $X_1, X_2, \dots, X_n$  be  $n$  topological spaces and  $\prod_{i=1}^n X_i$  and given any  $\alpha$ -open set  $U$  in  $Z$  containing  $f(p)$ , there exist  $\theta$ -open set  $E_i$  in  $X_i$  for  $i = 1, 2, \dots, n$  such that  $p \in \prod_{i=1}^n E_i$  and  $f(\prod_{i=1}^n E_i) \subset U$ . Then  $f$  is  $\alpha$ -unequivocally  $\theta$ -coherence.

**Proof:** Let  $p \in \prod_{i=1}^n X_i$  and  $U$  be any  $\alpha$ -open set in  $Z$  containing  $f(p)$ , there exist  $\theta$ -open set  $E_i$  in  $X_i$  for  $i = 1, 2, \dots, n$  such that  $p \in \prod_{i=1}^n E_i$  and  $f(\prod_{i=1}^n E_i) \subset U$ . Since,  $E_i \in \theta_0(X_i)$  for  $i = 1, 2, \dots, n$ . Therefore, by Lemma 3.3  $\prod_{i=1}^n E_i \in \theta_0(\prod_{i=1}^n X_i)$  for  $i = 1, 2, \dots, n$ . Thus,  $f$  is  $\alpha$ -unequivocally  $\theta$ -coherence.

## RESULTS AND DISCUSSION

**$\alpha\theta$ -Open function:** In this area new kind of function call  $\alpha\theta$ -open function study and we discover some portrayal and properties for it.

**Definition 4.1:** A function  $f: X \rightarrow Y$  is call  $\alpha\theta$ -open if and only if for each  $\alpha$ -open set  $G$  in  $X$ ,  $f(G) \in \theta_0(Y)$ . On the off chance that takes after quickly that each  $\alpha\theta$ -open function is quasi  $\alpha$ -open and thus, unequivocally  $\alpha$ -open, the opposite is not valid as observed from the accompanying illustration.

**Example 4.1:** Let  $X = \{a, b, c, d\}$  and  $\tau = \{x, \emptyset, \{a\}, \{c\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, c, d\}\}$ . The identity function  $i: (X, \tau) \rightarrow (X, \tau)$  is unequivocally  $\alpha$ -open but is not  $\alpha\theta$ -open function, since,  $\{a\} \in \alpha_0(X, \tau)$  but  $f(\{a\}) = \{a\} \notin \theta_0(X, \tau)$ . We discover a few portrayals and properties of  $\alpha\theta$ -open function.

**Theorem 4.1:** For any bijection function  $f: Y \rightarrow X$ , the accompanying are proportionate:

- The inverse function is  $\alpha$ -unequivocally  $\theta$ -coherence
- $f: Y \rightarrow X$  is  $\alpha\theta$ -open function

The following lemmas are used in sequel.

**Lemma 4.1; (Abdul-Jabbar, 2000):** The accompanying is valid, for each subset  $E$  of  $X$ :

$$X / Cl_0(E) = Int_\theta(X / E)$$

**Lemma 4.2:** The accompanying is valid for every subset  $E$  of  $X$ :

$$X / Cl_0(E) = \alpha Int_\theta(X / E)$$

**Theorem 4.2:** For a function  $f: Y \rightarrow X$  the accompanying are equal:

- $f$  is  $\alpha\theta$ -open function
- $f(\alpha Int(E)) \subset Int_\theta(f(E))$ , for each subset  $E$  of  $X$
- $\alpha Int(f^{-1}(W)) \subset f^{-1}(Int_\theta(W))$ , for each subset  $W$  of  $Y$
- $f^{-1}(Cl_\theta(W)) \subset \alpha Cl(f^{-1}(W))$ , for each subset  $W$  of  $Y$

**Proof:** (a) $\Rightarrow$ (b) Suppose  $f$  is  $\alpha\theta$ -open function and  $E \subset X$ . Since,  $\alpha Int \subset E$ ,  $f(\alpha Int(E)) \in \theta_0(Y)$  and  $f(\alpha Int(E)) \subset f(E)$  and hence,  $f(\alpha Int(E)) \subset Int_\theta(f(E))$ . Let  $W \subset Y$ . Then  $f^{-1}(W) \subset X$ , therefore, we apply (b), we obtain  $f(\alpha Int(f^{-1}(W))) \subset Int_\theta(f(f^{-1}(W)))$ . Then  $\alpha Int(f^{-1}(W)) \subset f^{-1}(Int_\theta(W))$ . (c) $\Rightarrow$ (d): let  $W \subset Y$ , then apply (c) to  $Y/W$ , we get  $\alpha Int(f^{-1}(W/Y)) \subset f^{-1}(Int_\theta(Y/W))$ . Then  $\alpha Int(X/f^{-1}(W)) \subset f^{-1}(Cl_\theta(W))$  which implies that  $X/\alpha Cl(f^{-1}(W)) \subset X/f^{-1}(Cl_\theta(W))$ . Hence  $f^{-1}(Cl_\theta(W)) \subset \alpha Cl(f^{-1}(W))$ . (d) $\Rightarrow$ (a): let  $G$  be any  $\alpha$ -open set in  $X$ . Then  $Y/f(G) \subset Y$ , apply (d), we obtain  $f^{-1}(Cl_\theta(Y/f(G))) \subset \alpha Cl(f^{-1}(Y/f(G)))$ . Then  $f^{-1}(Y/Int_\theta(f(G))) \subset \alpha Cl(X/G)$ . Which implies that  $X/f^{-1}(Int_\theta(f(G))) \subset X/Int_\theta = X/G$ . Therefore,  $G \subset f^{-1}(Int_\theta(f(G)))$ . Then  $f(G) \subset Int_\theta(f(G))$ . Therefore,  $f(G) \in \theta_0(Y)$ . Which completes the proof.

**Remark 4.1:** Let  $f: X \rightarrow Y$  be a bijective function. Then,  $f$  is  $\alpha\theta$ -open function if and only if  $f(F) \in \theta_0(Y)$ , for each  $\alpha$ -closed set  $F$  in  $X$ .

**Theorem 4.3:** If  $Y$  is regular space, then each  $s^{**}$ -open function is  $\alpha\theta$ -open.

**Proof:** Given  $G$  a chance to be any  $\alpha$ -open subset of  $X$ , then it is semi-open. Since,  $f$  is  $s^{**}$ -open function. Therefore,  $f(G)$  is open in  $Y$ . But  $Y$  is regular space, then by [1, Lemma 1.2.8]  $f(G)$  is  $\theta$ -open in  $Y$ . Which completes the proof.

**Theorem 4.4:** In the event that  $f: X \rightarrow Y$  is  $\theta$ -open function and  $E \subset X$  is an open set in  $X$ , at that point the  $f|_E: E \rightarrow Y$  is  $\alpha\theta$ -open function.

**Proof:** Let  $H$  be any  $\alpha$ -open set in the open subspace  $E$ . At that point, by [15, Theorem 3.7],  $H$  is  $\alpha$ -open in  $X$ . Since,  $f$  is  $\alpha\theta$ -open function. In this way,  $f(H)$  is  $\theta$ -open in  $Y$ . Hence,  $f|_E$  is  $\alpha\theta$ -open function.




Fig. 1: Growth of the graph

**Theorem 4.5:** Given  $f: X \rightarrow Y$  be a function and  $\{E_\alpha: \alpha \in \nabla\}$  be an open cover of  $X$ . If the restriction  $f|_{E_\alpha}: E_\alpha \rightarrow Y$  is  $\alpha\theta$ -open function for each  $\alpha \in \nabla$ , then  $f$  is  $\alpha\theta$ -open function.

**Proof:** Give  $H$  a chance to be any  $\alpha$ -open set in  $X$ . In this manner, by [15, Theorem 3.4],  $H \cap E_\alpha$  is  $\alpha$ -open in the subspace  $E_\alpha$  for each  $\alpha \in \nabla$ . Since,  $f|_{E_\alpha}$  is  $\alpha\theta$ -open function  $(f|_{E_\alpha})(H \cap E_\alpha)$  is  $\theta$ -open in  $Y$  and hence,  $f(H) = \cup \{(f|_{E_\alpha})(H \cap E_\alpha): \alpha \in \nabla\}$  is  $\theta$ -open in  $Y$ . This demonstrate  $f$  is  $\alpha\theta$ -open function.

**Remark 4.1:** Unmistakably  $\theta$ -compact and quasi  $H$ -closed equivalent from theorem 2.11 of (Ahmed and Yunis, 2002).

**Theorem 4.6:** In the event that  $f: X \rightarrow Y$  is  $\alpha\theta$ -open function and  $f(F)$  is  $\theta$ -compact relative to  $Y$ , then  $F$  is  $\alpha$ -compact subspace relative to  $X$ .

**Proof:** Let  $\{E_\alpha: \alpha \in \nabla\}$  be an open cover of  $F$ , then  $\{f(E_\alpha): \alpha \in \nabla\}$  is cover for  $f(F)$ . Since,  $f$  is  $\alpha\theta$ -open function. Therefore,  $f(E_\alpha) \in \theta\theta(Y)$  for each  $\alpha \in \nabla$ . Since,  $f(F)$  is  $\theta$ -compact relative to  $Y$ . Therefore, there exist a finite subfamily  $\{f(E_{\alpha_i}): i = 1, 2, \dots, n\}$  such that  $f(F) \subset \bigcup_{i=1}^n f(E_{\alpha_i})$ . Hence,  $F \subset \bigcup_{i=1}^n E_{\alpha_i}$ . Therefore,  $F$  is  $\alpha$ -compact subspace relative to  $X$ .

**Corollary 4.1:** If  $f: X \rightarrow Y$  is  $\alpha\theta$ -open surjective and  $Y$  is  $\theta$ -compact space, then  $X$  is  $\alpha$ -compact space.

**Theorem 4.7:** A function  $f: X \rightarrow Y$  is  $\alpha\theta$ -open if and only if for each subset  $S$  of  $Y$  and any  $\alpha$ -closed set  $F$  in  $X$  containing  $f^{-1}(S)$ , there exist a  $\theta$ -closed set  $M$  in  $Y$  containing  $S$  such that  $f^{-1}(M) \subset F$ .

**Proof:** Assume that  $f$  is  $\alpha\theta$ -open function. Let  $S \subset Y$  and  $F$  be an  $\alpha$ -closed set in  $X$  containing  $f^{-1}(S)$ . Put  $M = Y/f(X \setminus F)$ , then  $M$  is  $\theta$ -closed in  $Y$  and since,  $f^{-1}(S) \subset F$ , we have  $S \subset M$ . Since,  $f$  is  $\alpha\theta$ -open function and  $F$  is  $\alpha$ -closed in  $X$ ,  $M$  is  $\theta$ -closed in  $Y$ . It follows that  $f^{-1}(M) \subset F$ .

Conversely, let  $G$  be any  $\alpha$ -open subset of  $X$  and put  $S = Y/f$ . Then  $X/G$  is  $\alpha$ -closed set containing  $f^{-1}(G)$ . By hypothesis, there exist a  $\theta$ -closed set  $M$  in  $Y$  containing  $S$  such that  $f^{-1}(M) \subset X/G$ . Thus, we have  $f(G) \subset Y/M$ . On the other hand, we have  $f(G) = Y/S \supset Y/M$  and hence,  $f(G) = Y/M$ . Consequently,  $f(G)$  is  $\theta$ -open in  $Y$  and hence  $f$  is  $\alpha\theta$ -open function.

**Function with  $(\alpha, \theta)$ -closed graph:** In this area, we examine new properties of  $(\alpha, \theta)$ -closed graph (Chae *et al.*, 1995). Definition 5.1 (Chae *et al.*, 1995). Let,  $f(G) = \{(x, f(x)): x \in X\}$  be the graph of  $f: X \rightarrow Y$ , then is said to be  $(\alpha, \theta)$ -closed with respect to  $X \times Y$ , if for each point  $(x, y) \notin G(f)$ , there exist an open set  $U$  and an  $\alpha$ -open set  $H$  containing  $x$  and  $y$ , respectively, such that  $f(\bar{U}) \cap H = \emptyset$ . The accompanying diagram is a growth of the graph 4.1.1 of (Abdul-Jabbar, 2000). None of the suggestions is reversible (Fig. 1).

**Example 5.1:** Let  $X = \{a, b, c\}$  and,  $\tau = \{\varphi, X, \{a\}, \{a, b\}, \{a, c\}\}$ , then the function  $f: (X, \tau) \rightarrow (Y, \tau)$  defined as:  $f(x) = a$ , for each  $x \in X$  has  $\theta$ -closed graph which has not  $(\alpha, \theta)$ -closed graph.

**Theorem 5.1:** If  $f: X \rightarrow Z$  is a function with  $(\alpha\theta)$ -closed graph and  $f: X \rightarrow Z$   $\alpha$ -unequivocally  $\theta$ -coherence functions, then the set  $\{(x, y): f(x) = g(y)\}$  is  $\theta$ -closed in  $X \times Y$ .

**Proof:** Let  $E = \{(x, y): f(x) = g(y)\}$ . If  $(x, y) \in X \times Y/E$ , then  $f(x) \neq g(y)$ . Hence,  $(x, g(y)) \in (X, Z) \setminus G(f)$ . Since,  $f$  has  $(\alpha\theta)$ -closed graph. Therefore, there exist open set  $U$  and  $\alpha$ -unequivocally  $\theta$ -coherence of  $g$  implies that there is an open set  $V$  of  $X$  such that  $g(\bar{V}) \subset H$ . Therefore, we have  $f(\bar{U}) \times g(\bar{V}) = \emptyset$ . This established that  $f(\bar{U}) \times g(\bar{V}) \cap E = \emptyset$  which implies that  $(x, y) \notin Cl_\theta E$ .  $E$  is  $\theta$ -closed in  $X \times Y$ .

**Corollary 5.1:** If  $X$  is an Hausdorff space and  $f, g: X \rightarrow Y$  are  $\alpha$ -unequivocally  $\theta$ -coherence functions, then the set  $\{(x, y): f(x) = g(y)\}$  is  $\theta$ -closed in  $X \times Y$ .

**Theorem 5.2:** If  $f: X \rightarrow Y$  is any function with  $\theta$ -closed point inverses such that the image of each closure of open set is  $\alpha$ -closed, then has  $(\alpha\theta)$ -closed graph.

**Proof:** Let  $(x, y) \in X \times YG(f)$ . Then  $x \notin f^{-1}(y)$  and since,  $f^{-1}(y)$  is  $\theta$ -closed, there exist an open set  $U$  containing  $x$  such that  $\overline{U} \cap f^{-1}(y) = \emptyset$ . It follows that  $f(\overline{U})$  is  $\alpha$ -closed. Therefore, there is an  $\alpha$ -open set  $H$  in  $Y$  containing  $y$  such that  $f(\overline{U}) \cap H = \emptyset$ . Thus,  $f$  has  $(\alpha\theta)$ -closed graph.

**Theorem 5.3:** Let  $f: X \rightarrow Y$  be given function with  $(\alpha\theta)$ -closed graph, then for each  $x \in X$ ,  $\{f(x) = \cap \{\alpha\text{-Cl}(f(\overline{U})) : U \text{ is an open set of } x\}\}$  has.

**Proof:** Let the graph of the function be  $(\alpha\theta)$ -closed. Then it is claimed that for each  $x \in X$ ,  $\{f(x) = \cap \{\alpha\text{-Cl}(f(\overline{U})) : U \text{ is an open set of } x\}\}$ .

For if not, so, let  $y \neq f(x)$  such that  $y \in \cap \{\alpha\text{-Cl}(f(\overline{U})) : U \text{ is an open set of } x\}$ . Which implies that  $y \in \alpha\text{-Cl}(f(\overline{U}))$  for each open set of  $x$ ; it means that, for each  $\alpha$ -open set  $V$  of  $y$  in  $Y$ ,  $V \cap f(\overline{U}) \neq \emptyset$ . Thus, we obtain that  $(x, y) \in G(f)$  and there exist  $U$  and  $V$  such that  $V \cap f(\overline{U}) \neq \emptyset$  which implies that is contradiction. Thus,  $y = f(x)$ .

**Theorem 5.4:** Let  $f: X \rightarrow Y$  be a function with  $(\alpha\theta)$ -closed graph. If is quasi  $H$ -closed in  $X$ , then  $f(E)$  is has  $\alpha$ -closed in  $Y$ .

**Proof:** Let  $E$  be a quasi  $H$ -closed in  $X$ . Suppose that  $f(E)$  is not  $\alpha$ -closed in  $Y$ . Let  $y \notin f(E)$ . Therefore,  $y \neq f(x)$  for each  $x \in E$ . Since,  $f$  has  $(\alpha\theta)$ -closed graph. Therefore, there exist open set  $U_x$  and  $\alpha$ -open set  $H_x$  containing  $x$  and  $y$ , respectively such that  $f(\overline{U_x}) \cap H_x = \emptyset$ , for each  $x \in E$ . The family  $Q = \{U_x : x \in E\}$  is an open cover of  $E$ . Since,  $E$  is quasi  $H$ -closed, there exist a finite subfamily  $\{U_{x(i)}, \dots, U_{x(n)}\}$  of  $Q$  such that  $E \subset Y_{i=1}^n (\overline{U_{x(i)}})$ . Put  $H = \bigcap_{i=1}^n H_{x(i)}$ . Then:

$$f(E) \cap H \subset Y_{i=1}^n f(\overline{U_{x(i)}}) \cap H \subset Y_{i=1}^n f(\overline{U_{x(i)}}) \cap \left( \bigcap_{i=1}^n H_{x(i)} \right) = \emptyset$$

Since,  $H$  is an  $\alpha$ -open set containing  $y$ ,  $y \notin \alpha\text{-Cl}(f(E))$ . Therefore,  $\alpha\text{-Cl}(f(E)) \subset f(E)$ .

**Corollary 5.2:** The image of any quasi  $H$ -closed space in any space is  $\alpha$ -closed under functions with  $(\alpha\theta)$ -closed graphs.

**Theorem 5.4:** Let  $f: X \rightarrow Y$  be a given function. Then is  $G(f)$  is  $(\alpha\theta)$ -closed graph if and only if for each filter base  $\Psi$  in  $X$   $\theta$ -converges to some  $p$  in  $X$ ,  $f(\Psi)$   $\alpha$ -converges to some  $q$  in  $Y$ ,  $f(p) = q$ .

**Proof:** Suppose that Then  $G(f)$  is  $(\alpha\theta)$ -closed graph and let  $\Psi = \{E_\alpha : \alpha \in \nabla\}$  be a filter based in  $X$  such that  $\Psi \theta$ -converges to  $p$  and  $f(\Psi)$   $\alpha$ -converges to  $q$ . If  $f(p) \neq q$ , then  $(q, p) \notin G(f)$ . Thus, there exist open set  $U \subset X$  and  $\alpha$ -open set  $V \subset Y$  containing  $p$  and  $q$ , respectively, such that  $(\overline{U} \times V) \cap G(f) = \emptyset$ . Since,  $\Psi$   $\alpha$ -converges to  $p$  and  $f(\Psi)$   $\alpha$ -converges to  $q$ , there exist an  $E_\alpha \in \Psi$  such that  $E_\alpha \subset \overline{U}$  and  $f(E_\alpha) \subset V$ . Consequently,  $(\overline{U} \times V) \cap G(f) \neq \emptyset$  which is a contradiction.

Conversely, assume  $G(f)$  that is not  $(\alpha\theta)$ -closed graph. Then, there exist a point  $(p, q) \notin G(f)$  such that for each open set  $U \subset X$  and  $\alpha$ -open set  $V \subset Y$  containing  $p$  and  $q$ , respectively, such that  $(\overline{U} \times V) \cap G(f) \neq \emptyset$ . Define:

$$\Psi_1 = \{\overline{U_\alpha} : U_\alpha \text{ is an open set containing } p \text{ and } \alpha \in \nabla_1\}$$

$$\Psi_2 = \{V_\beta : V_\beta \text{ is an } \alpha\text{-open set containing } q \text{ and } \beta \in \nabla_2\}$$

$$\Psi_3 = \{E(\alpha, \beta) : E(\alpha, \beta) = (\overline{U_\alpha} \times V_\beta) \cap G(f), (\alpha, \beta) \in \nabla_1 \times \nabla_2\}$$

$\Psi = \{\Psi^*(\alpha, \beta) : (\alpha, \beta) \in \nabla_1 \times \nabla_2\}$  where  $\Psi^*(\alpha, \beta) = \{x \in U_x : (x, f(x)) \in E(\alpha, \beta)\}$ . Then  $\Psi$  is a filter base in  $X$  with property that  $\Psi$   $\alpha$ -converges to  $p$  and  $f(\Psi)$   $\alpha$ -converges to  $q$  and  $f(p) \neq q$ .

**Corollary 5.3:** A function  $f: X \rightarrow Y$  be has  $(\alpha\theta)$ -closed graph if and only if for each net  $X_\gamma$  in  $X$  such that  $X_\gamma \rightarrow \theta p \in X$  and  $f(x_\gamma) \rightarrow \alpha q \in Y$ ,  $f(p) = q$ .

## CONCLUSION

This study briefly described the  $\theta$ -open function, Quasi  $\theta$ -open and  $\theta$ -open their properties in this research.

## REFERENCES

- Abdul-Jabbar, A.M., 2000. Continuity, openness and closed graph in topological spaces. MSc Thesis, College of Science, Salahaddin-Erbil University, Erbil, Iraqi Kurdistan.
- Ahmed, N.K. and S.H. Yunis, 2002. Some equivalent concepts in topological spaces. Zanco., 14: 25-28.
- Ali, K.H., 2003. On semi-separation axioms. MSc Thesis, College of Education, Al-Mustansirah University, Baghdad, Iraq.
- Andrijevic, D., 1984. Some properties of the topology of  $\alpha$ -sets. Mat. Vesnik, 36: 1-10.
- Baker, C.W., 1986. Characterizations of some near-continuous functions and near-open functions. Intl. J. Math. Math. Sci., 9: 715-720.
- Chae, G.I., E. Hatir and S. Yuksel, 1995.  $\alpha$ -Strongly  $\theta$ -continuous functions. J. Nat. Sci., 5: 59-66.

- Chae, G.I., T. Noiri and D.W. Lee, 1986. On NA-continuous functions. *Kyungpook Math. J.*, 26: 73-79.
- Dube, K.K., L.J. Yoon and O.S. Panwar, 1998. A note of semi-closed graph. *ULT. Rep.*, 2: 379-383.
- Faro, G.L., 1987. On strongly  $\alpha$ -irresolute mappings. *Indian J. Pure Appl. Math.*, 18: 146-151.
- Jankovic, D., 1985. On semi separation properties. *Indian J. Pure Appl. Math.*, 16: 957-964.
- Jankovic, D.S., I.L. Reilly and M.K. Vamanamurthy, 1988. On strongly compact topological spaces. *Quest. Ans. Gen. Topol.*, 6: 29-40.
- Lee, H.W., G.I. Chae and D.W. Lee, 1985. Feebly irresolute functions. *Sungshin Women's Univ. Rep.*, 21: 273-280.
- Levine, N., 1963. Semi-open sets and semi-continuity in topological spaces. *Amer. Math. Monthly*, 70: 36-41.
- Long, P.E. and L. Herrington, 1982. The  $T_0$ -topology and faintly continuous functions. *Kyungpook Math. J.*, 22: 7-14.
- Long, P.E. and L.L. Herring, 1977. Functions with unequivocally graphs. *Bool. Univ. Mat. Ital.*, 12: 381-384.
- Maheshwari, S.N., G.I. Chae and P.C. Jain, 1982. Almost feebly continuous functions. *UIT. Rep.*, 13: 195-197.
- Maheshwari, S.N., I.A. Hasanein and S.N. El-Deeb, 1983.  $\alpha$ -Continuous and  $\theta$ -open mappings. *Acta Math. Hung.*, 41: 213-218.
- Njastad, O., 1965. On some classes of nearly open sets. *Pac. J. Math.*, 15: 961-970.
- Noiri, T. and S.M. Kang, 1984. On almost strongly  $\alpha$ -continuous functions. *Indian J. Pure Appl. Math.*, 15: 1-8.
- Noiri, T., 1973. Remarks on semi-open mappings. *Bull. Calcutta Math. Soc.*, 65: 197-201.
- Noiri, T., 1975. Properties of theta-continuous functions. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Nat.*, 58: 887-891.
- Noiri, T., 1980. On  $\alpha$ -continuous functions. *J. Korean Math. Soc.*, 16: 161-166.
- Porter, J. and J. Thomas, 1969. On H-closed and minimal hausdorff spaces. *Trans. Am. Math. Soc.*, 138: 159-170.
- Singal, M.K. and S.P. Arya, 1969. On almost-regular spaces. *Glasnik Mat.*, 4: 89-99.
- Thakur, S.M.S., 1980. On  $\alpha$ -irresolute functions. *Tamkang J. Math.*, 11: 209-214.
- Thivagar, M.L., 1991. Generalization of pairwise  $\alpha$ -continuous functions. *Pure Appl. Math. Sci.*, 33: 55-63.
- Velicko, N.V., 1968. H-closed topological spaces. *Amer. Math. Soc. Transl.*, 78: 103-118.