

## Occurrence of *Aspergillus* Species among Apparently Healthy Chickens in Poultry Farms in Kaduna State, Nigeria

<sup>1</sup>Clara N. Kwanashie, <sup>2</sup>Paul A. Abdu, <sup>1</sup>Haruna M. Kazeem and <sup>3</sup>Jarlath U. Umoh

<sup>1</sup>Department of Veterinary Microbiology, <sup>2</sup>Department of Veterinary Medicine,

<sup>3</sup>Department of Veterinary Public Health and Preventive Medicine,

Faculty of Veterinary Medicine, Ahmadu Bello University,

P.M.B. 1069 Zaria, Kaduna State, Nigeria

---

**Abstract:** The study was conducted to determine the occurrence of *Aspergillus* species among apparently healthy birds in poultry farms. Trachea swabs were collected from a total of 1500 birds in 52 commercial (10% of birds in each poultry farm visited) poultry farms were sampled. *Aspergillus* sp. was isolated from 718 (47.87%) of the birds. Six species of *Aspergillus* were isolated viz: *A. fumigatus* made up 52.37% (376) of the *Aspergillus* isolates followed by *A. flavus* 21.87% (157), *A. niger* 11.42% (82), *A. terreus* 8.64% (62), *A. restrictus* 2.79% (20) and *A. ochraceous* 2.92% (21). *Aspergillus* species were isolated throughout the year though with a higher incidence during the rainy season compared to the dry season.

**Key words:** *Aspergillus*, chickens, poultry farms, occurrence, Kaduna State, Nigeria

---

### INTRODUCTION

Fungi species belonging to the genus *Aspergillus* represent one of the most important among the fungi that affect poultry. *Aspergillus* species especially *Aspergillus fumigatus* is responsible for the disease called Aspergillosis in both poultry and humans (Hashempour *et al.*, 2011). Other species of the genus such as *A. flavus*, *A. niger* and *A. nidulans* have also been reported to cause the disease as well as been responsible for mycotoxicosis (aflatoxicosis) in poultry (Beernaert *et al.*, 2009). The warm and humid environment of the poultry house favors the growth of *Aspergillus* sp. *Aspergillus* are ubiquitous and opportunistic pathogens capable of growing on most substrates and producing spores that may contaminate the environment and air of the poultry house. The spores may get inhaled by birds and causing disease especially in immunocompromised birds or birds that inhale a high dose of the spores (Beernaert *et al.*, 2009). The disease represents an important threat to poultry production worldwide causing significant morbidity and mortality in both young and adult birds (Khosravinia *et al.*, 2009). The disease is characterized by non specific clinical signs affecting the respiratory system (dyspnea, gasping and cyanosis) and also causes weight loss, loss of appetite. Disseminated forms (via the hematogenous routes) of the disease affecting other organs can also occur (Beernaert *et al.*, 2009).

The study was conducted to determine the occurrence of *Aspergillus* species among apparently healthy birds in light of the threat they may pose to the birds and the poultry attendants.

### MATERIALS AND METHODS

**Study area:** The study area was farms located in Zaria and Kaduna of Kaduna State, Nigeria. Kaduna State is located between latitudes 11°32' and 9°20'N and longitude 8°50' and 6°51'E. The state is positioned in the Northern Guinea Savannah zone of Nigeria. The area is characterized by a cold dry season (November to February), hot-dry season (March to April) (both dry seasons) and the wet/rainy season (May to October) (Ayo *et al.*, 1999). The annual rainfall peaks in the month of August with the average of 146 mm. The average humidity is highest in August with 75.6 mm Hg<sup>-1</sup> and lowest at the months of December to January with 38.2 mm Hg<sup>-1</sup>. The mean temperatures for the zone are 10.7 and 38.75°C minimum and maximum respectively (Agbogu *et al.*, 2006).

**Study population and sampling:** The study population comprised of 10% (1,500) of the total population of birds (15,000) in 52 poultry farms visited. These farms were selected based on clinical records of the farms from the Poultry Ambulatory Unit of the Veterinary Teaching Hospital, Ahmadu Bello University, Zaria.

A total number of 1,500 chicken tracheal swabs were obtained from apparently healthy chickens. In each of the farm visited, trachea swabs were collected randomly from 10% of the chickens. Sampling was conducted all year round (January to December) to observe for seasonal variation in *Aspergillus* species isolation. All the samples were labeled and taken to the laboratory for culture and isolation of fungi.

**Fungal isolation and identification:** The fungi isolation was carried out using Sabouraud's Dextrose Agar (SDA) (Oxoid UK) medium, prepared according to the manufacturer's specification (Oxoid UK) with addition of 0.05 mg mL<sup>-1</sup> of chloramphenicol to control bacterial contamination (Ainstworth and Austwick, 1973). Each swab was inoculated on the SDA slant in 10 mL universal bottles, labeled, incubated at room temperature and at 37°C with daily observation for a period of 7 days. All the samples that had no fungal growth were discarded after 7 days (Ainstworth and Austwick, 1973).

The growths from the cultures were observed for the colonial morphologies i.e., size, color, topography and aerial growth and these characteristics were used to identify the genus and species. Growths were prepared for microscopic observation by staining them on glass slides using lactophenol cotton blue stain. The stained slides were observed under x10 magnification using a light microscope (Olympus USA). Characteristics of fungi such as hyphae, conidial heads and the arrangement of the conidia were also observed and used for the fungi identification.

## RESULTS AND DISCUSSION

*Aspergillus* species were isolated from 718 (47.87%) of the 1500 apparently healthy birds sampled. Six *Aspergillus* isolates were identified with *A. fumigatus* presenting as the major species isolated 376 (52.37%) followed by *A. flavus* 157 (21.87%), *A. niger* 82 (11.42%), *A. terreus* 62 (8.64%), *A. restrictus* 20 (2.79%) and *A. ochraceus* 21 (2.92%) (Table 1).

*Aspergillus* species was also isolated both in the rainy and dry season. *A. fumigatus* had the highest rate of occurrence/isolation in both the rainy 262 (59.95%) and dry seasons 114 (40.57%). This is followed by *A. flavus* 72 (16.48%) and 85 (30.25%) and *A. niger* 38 (8.70%) and 44 (15.66%) in the rainy and dry seasons respectively (Table 2).

The results of this study have shown that *Aspergillus* species are present among apparently healthy birds. This species of fungi has been reported to be one of the most common fungal infections of poultry

Table 1: Distribution of *Aspergillus* species isolated from apparently healthy chickens in 52 poultry farms across Kaduna State, Nigeria

<i>Aspergillus</i> species	No. of isolates (%)
<i>A. fumigatus</i>	376 (52.37)
<i>A. flavus</i>	157 (21.87)
<i>A. niger</i>	82 (11.42)
<i>A. terreus</i>	62 (8.64)
<i>A. restrictus</i>	20 (2.79)
<i>A. ochraceus</i>	21 (2.92)
Total	718.00

Table 2: Distribution of *Aspergillus* species isolated from apparently healthy chickens during the rainy and dry seasons in 52 poultry farms across Kaduna State, Nigeria

<i>Aspergillus</i> species	No. of isolates (%)	
	Rainy (May to Oct.)	Dry season (Nov. to Apr.)
<i>A. fumigatus</i>	262 (59.95)	114 (40.57)
<i>A. flavus</i>	72 (16.48)	85 (30.25)
<i>A. niger</i>	38 (8.70)	44 (15.66)
<i>A. terreus</i>	29 (6.64)	33 (11.74)
<i>A. restrictus</i>	15 (3.43)	5 (1.78)
<i>A. ochraceus</i>	21 (4.84)	0.00
Total	437.00	281.00

(Greenacre *et al.*, 1992). The spores of *Aspergillus* especially *A. fumigatus* has been reported to be highly thermotolerant growing at temperatures between 15-53°C (Greenacre *et al.*, 1992) which is consistent with the average minimum (10.7°C) and maximum (38.75°C) temperature for this zone (Agbogun *et al.*, 2006). This may likely be the reason for the presence of the fungi all year round. *A. fumigatus* accounts for about 95% of aspergillus cases in birds followed by *A. flavus* (Deem, 2003). *Aspergillus* species have been reported to produce toxic metabolites which are responsible for the clinicopathologic manifestations seen in cases of aspergillosis. *A. fumigatus* produces gliotoxin, *A. flavus* produces aflatoxin B1 while *A. niger* produces oxalic acid (Joseph, 2000). The high prevalence of these species is of economic importance as is an indication of possibility of aspergillosis which is associated with a high morbidity and mortality (Abdu *et al.*, 1995). This is the first time of isolating *A. ochraceus* from apparently healthy chickens in this environment this species is highly important in aflatoxicosis and of public health concern. *A. terreus* and *A. niger* though not commonly isolated are pathogens of birds and man (Redig *et al.*, 1980). Some of these fungi especially those that produce mycotoxins can contaminate the feed of the birds which they may subsequently consume resulting in mycotoxicosis.

The study has shown that *Aspergillus* sp. are present among apparently healthy birds with a high prevalence. The presence of these fungi may pose a potential threat to the health of the birds and source of economic loss to the farmer in the event of an outbreak.

## CONCLUSION

Fungal infection in poultry is very difficult and expensive to treat with a guarded prognosis (Joseph, 2000). Therefore, prevention is the most effective way of combating this disease. Proper ventilation and regular cleaning with fungicides especially after removal of fecal or decaying matter is very important. Stressors which have been shown to play an important role in the development of fungal disease should be minimized.

## REFERENCES

- Abdu, P.A., S.U. Hassan, C.N. Kwanashie and N.D.G. Ibrahim, 1995. Aspergillosis in fowls. Bull. Anim. Health Prod., 43: 93-94.
- Agbogbo, V.N., V.J. Umoh, C.A. Okufo, S.I. Smith and J.B. Ameh, 2006. Study of the bacteriological and physicochemical indicators of pollution of surface waters in Zaria, Nigeria. Afr. J. Biotechnol., 5: 732-737.
- Ainstworth, G.C. and P.K.C. Austwick, 1973. Fungal Diseases of Animals. 2nd Edn., Commonwealth Agricultural Bureaux, UK., pp: 37-88.
- Ayo, J.O., S.B. Oladele, S. Ngam, A. Fayomi and S.B. Afolayan, 1999. Diurnal fluctuations in rectal temperature of the Red Sokoto goat during the harmattan season. Res. Vet. Sci., 66: 7-9.
- Beermaert, L.A., F. Pasmans, K. Baert, L. van Waeyenberghe, K. Chiers, F. Haesebrouck and A. Martel, 2009. Designing a treatment protocol with voriconazole to eliminate *Aspergillus fumigatus* from experimentally inoculated pigeons. Vet. Microbiol., 139: 393-397.
- Deem, S.L., 2003. Fungal diseases of birds of prey. Vet. Clin. North Am. Exotic Anim. Pract., 6: 363-376.
- Greenacre, C.B., K.S. Latimer and B.W. Ritchie, 1992. Leg paresis in a black palm cockatoo (*Probosciger aterrimus*) caused by aspergillosis. J. Zoo Wildlife Med., 23: 122-126.
- Hashempour, A., M.H.S. Zali, R. Delshad, V.R. Karamad, V. Farzayi and M. Kalbkhani, 2011. A study on the existence of *Aspergillus* in birds in the farms around Urmia-Iran. J. Stored Prod. Postharvest Res., 2: 235-236.
- Joseph, V., 2000. Aspergillosis in raptors. Semin. Avian Exotic Pet Med., 9: 66-74.
- Khosravinia, H., M.H. Gharoni and M. Darvishnia, 2009. Litter mycology and the impacts of litter type and preslaughter feed withdrawal on crop bacterial community in broiler chicken. Afr. J. Microbiol. Res., 3: 844-850.
- Redig, P.T., M.R. Fuller and D.L. Evans, 1980. Prevalence of *Aspergillus fumigatus* in free-living goshawks (*Accipiter gentilis atricapillus*). J. Wildlife Dis., 16: 169-174.