

## Toluene Exposure and its Association to Hippuric Acid and Non-Systematic Symptoms among Outdoor Workers in Urban Area, Bangkok, Thailand

<sup>1</sup>Tanasorn Tunsaringkarn, <sup>2</sup>Tassanee Prueksasit,  
<sup>1</sup>Saowanee Semathong, <sup>1</sup>Anusorn Rungsiyothin and <sup>1</sup>Kalaya Zapaung  
<sup>1</sup>College of Public Health Sciences, <sup>2</sup>Department of General Science,  
Faculty of Science, Chulalongkorn University, 10330 Bangkok, Thailand

**Abstract:** Volatile Organic Compounds (VOCs) are mainly from motor vehicle emissions which cause of serious problems of air pollution in Bangkok Metropolitan, Thailand. Toluene is the most abundance of aromatic hydrocarbon found in ambient air. This study aimed to evaluate toluene exposures and its association to non-systematic symptoms among outdoor workers in Pathumwan District, central Bangkok. A cross-sectional study was conducted in urban area of Bangkok during September 2012 to March 2013. Two hundred and eighty seven outdoor workers of security guardsmen, motorcycle drivers and street vendors were included in this study for interviewing and urine collection. Thirty six ambient air samples were collected for toluene analyses by Gas Chromatography with Flam Ionized Detector (GC/FID). The results showed that average ambient air toluene was 91.1  $\mu\text{g}/\text{m}^3$  which street vendor was the highest exposed as well as average hippuric acid level was 0.4 g/gCr. Non-systematic symptoms prevalence of workers were headache, dizziness, weakness and throat irritation at 42.5, 16.6, 36.6 and 15.3%, respectively which they were highest in street vendors. Toluene exposure level was significantly associated to hippuric acid ( $p = 0.001$ ) and non-systematic symptoms of dizziness and throat irritation ( $p < 0.01$ ). However, hippuric acid was not associated to any non-systematic symptoms. There were trended higher toluene exposures at roadside area of street vendors and motor cycle drivers than security guardsmen in university area as well as non-specific symptoms. There were the associations between ambient air toluene level and urinary biomarker of hippuric acid and non-systematic symptom of dizziness.

**Key words:** Toluene, hippuric acid, non-systematic symptom, outdoor worker, dizziness

---

### INTRODUCTION

Toluene is formed during the combustion of organic materials and widely used as a solvent in paints, varnishes, pesticide formulations, printing inks, dyes, adhesives, sealants, cleaning agents, nail polish and for chemical extractions (Alberta Environment, 2005). Volatile Organic Compounds (VOCs) constitute one of the most important major air pollutant in the atmospheric. Toluene is the most abundance of aromatic hydrocarbon found in ambient air and can also adversely affect to human health (US EPA, 1994) especially, urban area which vehicle exhaust was considered to be the dominant source (Ameno *et al.*, 1992; Al-Khulaifi *et al.*, 2014). Toluene is rapidly absorbed into the bloodstream by inhalation exposure. Once absorbed, it is rapidly distributed to white adipose tissue, adrenals, skin, kidneys, liver, lung and brain (WHO, 1985; EPA, 1993). Hippuric acid is the

primary urinary metabolite of toluene can be used as an indicator of toluene exposure (Duydu *et al.*, 1999; Bahrami *et al.*, 2005). It is well established that neurotoxicity and neurobehavioural deficits are the principal effects of long-term inhalation exposure to toluene in both humans and experimental animals (Brugnone *et al.*, 1995; ATSDR, 2000) but insufficient information on possible exposure to other chromosome-damaging agents (IARC, 1999; EPA, 2005; Tunsaringkarn *et al.*, 2011). Most human carcinogens are genotoxic but all of genotoxic agents have not been carcinogenic in human. Rapid increasing population as well as motor vehicle number per year which most them use biofuel as gasohol and biodiesel, Light Petroleum Gas (LPG) as government policies leading to serious problem of outdoor workers in Bangkok such as mortality, hospital admissions cardiovascular and respiratory disease, asthma attacks, acute bronchitis, respiratory

symptoms and restrictions in activity (Kampa and Castanas, 2008; WHO, 2014). The global burden of disease estimated that outdoor air pollution was account for 1.4% of total mortality, 0.5% of all Disability-Adjusted Life Years (DALYs) and 2% of all cardiopulmonary disease (Ezzati *et al.*, 2002; WHO, 2002; Cohen *et al.*, 2004). The WHO estimates indicated that 81% of the attributable deaths from air pollution (WHO, 2002). Outdoor workers in urban area are also directly exposed to toluene in atmosphere. It should have risk of long-term effects on non-systematic symptoms. This study aimed to evaluate toluene in ambient air and its association to hippuric acid and non-systematic symptoms among outdoor workers in Urban Area, Central Bangkok, Thailand.

**MATERIALS AND METHODS**

**Study sites:** The study was carried out in 32 samples from 16 monitoring sites consist of 10 sites of motorcycle drivers and street vendors on the main roadsides of Rama road 1, Phythai road, Henrydunant road and 6 sites of security guardsmen in Chulalongkorn University which all sites are in Pathumwan area, central Bangkok, Thailand. Main study sites of roadsides were located at Community Pharmacy Clerkship (CPC), Bangkok Bank Public Company Limited (BBL), Krung Thai Bank Public

Company Limited (KTB), Siam Commercial Bank Public Company Limited (SCB), Faculty of Science (FS) and study sites in university were located at Satit Chula (SC), university entrance-exit gate 1-4 and checkpoint behind Faculty of Science as shown in Fig. 1.

**Population study:** Total of 287 outdoor workers were included by random in this study including 76 security guardsmen of university, 76 motorcycle drivers and 135 street vendors. All inclusion criteria of workers were healthy, age over 18 years old, work period >3 months and signed consent from and interviewed the characteristics of age, weight, height, period of employment, working hour a day, working day a week and non-systematic symptoms at last 3 months. This research was approved by the Ethical Review Committee for Research Involving Human Research Subjects, Health Science Group, Chulalongkorn University (COA No. 089/2012).

**Ambient air sample collection and analyses:** Total of 32 ambient air monitoring study sites were collected for 8 h working time of workers during September 2012 to March 2013. Ambient air samples were collected by active charcoal tube with flow rate 100 mL min<sup>-1</sup> for toluene analyses by GC/FID (Tunsaringkarn *et al.*, 2012). Sample tubes were kept at 4°C during transportation to the laboratory and were stored in a refrigerator until further analysis.



Fig. 1: Study site map; ● motorcycle drivers and street vendors at roadsides; ★ security guards in university

The charcoal tube was extracted with Carbondisulphide (CS) and the sample solution was then analyzed by Gas Chromatography (GC) with an FID detector (Varian, CP3800) and column RP Amide Discovery C16 (25 m×4.0 mm with 0.5 µm packing). The initial oven temperature was set to 40°C, increased at a rate of 10°C min<sup>-1</sup> 100°C and held at that temperature for 2 min; FID temperature was set at 225°C. The carrier gas was he pushed at a flow rate of 1.0 mL min<sup>-1</sup>. Toluene levels was analyzed under the relative intensity of chromatographic signal for 20 min. Quantification was performed using the External Calibration Method. The coefficients of toluene determination (R<sup>2</sup>) ranges were 0.991-0.999. Each sample was analyzed by triplicate.

**Urinary sample collections and analyses:** The cross-sectional study of 287 urine samples of all outdoor workers were collected by glass bottle after 8 h working time (8.00 am to 15.00 pm). Urine samples were kept at 4°C during transportation to laboratory, stored at -20°C and later analyzed for Creatinine (Cr) (Slot, 1965), hippuric acid (Ogata and Taguchi, 1987). All urinary samples were analyzed at a standard Spacial Laboratory, Bangkok, Thailand. Urinary creatinine concentrations were accepted between 0.3 and 3.0 g L<sup>-1</sup> (ACGIH, 2010) as the World Health Organization (WHO) has adopted guideline. All measured values were corrected by urinary Cr concentrations for clinical chemistry analysis (Barr *et al.*, 2005).

**Statistical analysis:** All data were statistically analyzed using the SPSS 17.0 for Windows Software. Descriptive statistics was used to understand the characteristics, exposures and urinary biomarkers concentrations. The comparison of ambient air toluene and urinary biomarkers

of hippuric acid were done using independent t-test. The association between ambient air VOCs levels and urinary biomarkers of outdoor workers were assessed by Linear Regression Model. A statistically significant difference was accepted at p<0.05 as other medical studies.

**RESULTS**

**Characteristic, ambient air toluene, hippuric acid of outdoor workers:** Average age, BMI, period of employment, working hour a day and working day a week of all workers were 40.3 years, 24.5 kg/m<sup>2</sup>, 8.5 years, 10.8 h and 6.5 days, respectively (Table 1). Age and period of employment of security guardsmen were significantly higher than motorcycle drivers (p<0.05) and period of employment of security guardsmen was significantly higher than street vendors (p<0.01). Working hour a day of security guardsmen was higher than motorcycle drivers (p<0.01) and was higher than street vendors (p<0.05). While working day a week of security guardsmen was higher than street vendors (p<0.05). Average ambient air toluene was 91.1 µg/m<sup>3</sup> while hippuric acid of all workers was 0.4 g/gCr.

**Non-systematic symptoms of outdoor workers:** Non-systematic symptoms of all outdoor workers were headache, dizziness, weakness and throat irritation at 42.5, 16.6, 36.6 and 15.3%, respectively (Table 2). All symptoms were highest in street vendor group.

**Association between ambient air toluene and hippuric acid and non-systematic symptoms and association between hippuric acid and non-systematic symptoms of outdoor workers:** The ambient air toluene was significantly associated to hippuric acid (p = 0.001) and

Table 1: General characteristic, ambient air toluene exposure and of hippuric acid among outdoor workers

Parameters	Mean±SD			Average
	Security guardsman (n = 76)	Motorcycle driver (n = 76)	Street vender (n = 135)	
Age (years)	43.5±0.4*	37.9±0.80*	39.4±0.80	40.3±2.90
BMI (kg/m <sup>2</sup> )	24.5±0.4	24.4±0.20	24.8±0.10	24.5±0.20
Period of employment (years)	12.8±0.1* **	7.0±0.20*	5.7±0.20**	8.5±3.80
Working hour per day (h)	10.5±0.3**	12.0±0.10**, <sup>†</sup>	10.1±0.10 <sup>†</sup>	10.8±1.00
Working day per week (days)	6.4±0.1*	6.6±0.10	6.1±0.10*	6.5±0.20
Ambient air toluene (µg/m <sup>3</sup> )	31.7±4.3	88.5±76.0	157.1±77.4	91.1±57.6
Hippuric acid (g/gCr)	0.3±0.3	0.3±0.30	0.5±0.50	0.4±0.10

\*<sup>†</sup>Significant difference between worker group at p<0.05; \*\*significant difference between worker group at p<0.01

Table 2: Non-systematic symptoms among outdoor workers

Non-systematic symptoms	n (%)			
	Security guardsman (n = 76)	Motorcycle driver (n = 76)	Street vender (n = 135)	Total (n = 287)
Headache	17 (22.4)	29 (38.2)	76 (56.3)	122 (42.5)
Dizziness	5 (6.6)	9 (11.8)	34 (25.2)	48 (16.6)
Weakness	15 (6.6)	30 (19.7)	61 (25.9)	106 (36.6)
Throat irritation	8 (10.5)	10 (13.2)	29 (21.5)	44 (15.3)

Table 3: Association between toluene exposures, hippuric acid and non-systematic symptom parameters of outdoor workers

Independence parameters	Dependence parameters	Linear regression model results <sup>#</sup>		
		Standardize coefficient	95% CI	p-values
Toluene	Hippuric acid	0.311	0.001-0.004	0.001
	Headache	0.000	0.000-0.001	0.059
	Dizziness	0.000	0.000-0.001	0.006
	Weakness	0.000	-0.003-0.002	0.636
	Throat irritation	0.000	0.000-0.001	0.002
Hippuric acid	Headache	0.020	-0.173-0.218	0.818
	Dizziness	0.034	-0.111-0.165	0.696
	Weakness	0.024	-0.169-0.222	0.790
	Throat irritation	-0.116	-0.221-0.046	0.197

Adjust by age, BMI, period of employment, working hours a day and working days a week

non-systematic symptoms of dizziness ( $p < 0.01$ ) and throat irritation ( $p < 0.01$ ) (Table 3). In addition, ambient air toluene was nearly significantly associated to headache symptom ( $p = 0.059$ ). However, hippuric acid was not associated to any non-systematic symptom of outdoor workers.

### DISCUSSION

Most of all workers had normal BMI while their work employment was 8.5 years. The motorcycle drivers worked for longest time a day (12.0 h) but street vendors was highest exposed to toluene ( $157.1 \mu\text{g}/\text{m}^3$ ). The average toluene ( $91.1 \mu\text{g}/\text{m}^3$ ) in this study was in urban air concentration range  $5\text{-}150 \mu\text{g}/\text{m}^3$  which present toluene concentration was higher than previous study in Bangkok 2007 ( $36.1 \mu\text{g}/\text{m}^3$ ) (Laowagul *et al.*, 2008). Moreover, this toluene level was higher in the comparison of previous reported data in urban cities of other countries;  $4\text{-}36 \mu\text{g}/\text{m}^3$  in Hong Kong (Lee *et al.*, 2002),  $27 \mu\text{g}/\text{m}^3$  in Taiwan (Hsieh and Tsai, 2003),  $39.2 \mu\text{g}/\text{m}^3$  in Algeria (Kerbachi *et al.*, 2006),  $13.0\text{-}33.1 \mu\text{g}/\text{m}^3$  in ITO, Delhi (Kumar and Tyagi, 2006) and  $11.52 \mu\text{g}/\text{m}^3$  in Suzhou, China (Wang *et al.*, 2010). It may be higher if it close to industrial emission sources (WHO, 2000). The acute and chronic effects of toluene on Central Nervous System (CNS) are the effects of the most concern (Dann *et al.*, 1989; WHO, 2000).

This study found that most of workers had non-systematic symptoms of 42.5% headache, 36.6% weakness, 16.6% dizziness and 15.3% throat irritation which were also found in gasoline station workers (Tunsaringkarn *et al.*, 2012). The street vendor had the highest of these symptoms while the security guardsman was the lowest symptoms. Ambient air toluene was significantly associated with dizziness ( $p < 0.01$ ) and throat irritation symptoms ( $p < 0.01$ ). In addition, ambient air toluene was strongly associated with hippuric acid ( $p = 0.001$ ). It can supported that hippuric acid is used as

biomarker of toluene exposure (Pagnotto and Lieberman, 1967; Kawai *et al.*, 2008) which hippuric acid excretion has been correlated with the Time Weighted Average concentration (TWA) of toluene during the workshift (Duydu *et al.*, 1999). But insufficiency of urinary hippuric acid to indicate the low exposures to toluene as previous study (Angerer and Kramer, 1997; Duydu *et al.*, 1999). Average hippuric acid concentration of all workers in this study was  $0.4 \text{ g/gCr}$  which was the same value of press workers from factories in Bangkok ( $0.37 \text{ g/gCr}$ ; range  $0.06\text{-}1.98 \text{ g/gCr}$ ; Wiwanitkit *et al.*, 2002) but higher than gas service station workers in Chonburi Province ( $0.28 \text{ g/gCr}$ ; Yimrungruang *et al.*, 2008). Moreover, toluene showed the association with non-specific symptoms of dizziness and throat irritation which supported other study including headache, dizziness, fatigue, muscular weakness, drowsiness and incoordination with staggering gait, skin paresthesia, collapse and coma (US EPA, 1980). The previous study of the subjective symptoms were measured with the psychological/neurological questionnaire (Psychologisch-Neurologischer Fragebogen, PNF) have been reported in a significant dose-response relation ( $50\text{-}100 \text{ ppm}$ ) and long term exposure ( $>12$  years) which scored significantly higher on a symptom index than shorter exposure ( $1\text{-}12$  years) (Ukai *et al.*, 1993). About 20% of absorbed toluene dose is excreted unchanged by the lungs which the rest is mainly metabolized to hippuric acid and very small percentage of the toluene is metabolized to cresol (Angerer, 1979). So, hippuric acid can found in urine which occupational worker was limited at  $1.6 \text{ g/gCr}$  (ACGIH, 2009a, b). This study found hippuric acid was lower than limited level and it was not associated with any symptoms in this study as Kim *et al.* (1989) and Hashim *et al.* (2006) studies found that there were no correlation between hippuric acid concentrations and subjective symptoms and pulmonary function parameters.

However, the air quality guideline could also be based on the odour threshold. The peak concentrations of toluene in air should be kept below the odour detection threshold level of  $1 \text{ mg}/\text{m}^3$  ( $0.266 \text{ ppm}$ ) as a 30 min average as WHO recommended (WHO, 2000).

### CONCLUSION

Average ambient air toluene concentration (8 h average) which widespread in the environmental of urban area, Bangkok was  $91.1 \mu\text{g}/\text{m}^3$  ( $24.2 \text{ ppb}$ ) with a range of  $28.7\text{-}142.2 \mu\text{g}/\text{m}^3$  ( $7.6\text{-}37.8 \text{ ppb}$ ). Outdoor workers of guardsmen, motorcycle drivers and street vendors exposed to considerably higher levels than the general population. Most non-systematic symptoms prevalence of outdoor workers found headache, weakness, dizziness

and throat irritation respectively which dizziness and throat irritation were associated with ambient air toluene exposure. As well as urine hippuric acid was strongly associated with toluene exposure.

However, urine hippuric acid was good an indicator of toluene exposure but did not seem to be a good predictor of non-systematic symptoms of outdoor workers. It should be future studied in large population group.

#### ACKNOWLEDGEMENTS

This research was funded by the National Research Council of Thailand (NRCT) and facilities supported by College of Public Health Sciences, Chulalongkorn University. Researchers would like to thank the Ethical Review Committee for Research Involving Human Research Subjects, Health Science Group, Chulalongkorn University for their suggestion and approval of this study. Researchers also thank to all workers for their kind collaborations and lastly, thank researcher for advice and editing the manuscript.

#### REFERENCES

- ACGIH., 2009a. Guide to occupational exposure values. Proceedings of the American Conference of Governmental Industrial Hygienists, June 11, 2009, Cincinnati, Ohio.
- ACGIH., 2009b. Introduction to biological exposure indices. In TLVs and BEIs Based on the Documentation of the Threshold Limit Values for Chemical Substances and Physical Agents and Biological Exposure Indices. Cincinnati, Ohio.
- ACGIH., 2010. Introduction to biological exposure indices. In TLVs and BEIs Based on the Documentation of the Threshold Limit Values for Chemical Substances and Physical Agents and Biological Exposure Indices. Cincinnati, Ohio.
- ATSDR., 2000. Toxic profile for toluene: Health effect. Agency for Toxic Substances and Disease Registry. <http://www.atsdr.cdc.gov/toxprofiles/tp56-c2.pdf>.
- Al-Khulaifi, N.M., H.F. Al-Mudhaf, R. Alenezi, A.S.I. Abu-Shady and M.I. Selim, 2014. Seasonal and temporal variations in volatile organic compounds in indoor and outdoor air in Al-Jahra city, Kuwait. *J. Environ. Protect.*, 5: 310-326.
- Alberta Environment, 2005. Alberta ambient air quality objectives: Toluene. May 1, 2005. <http://environment.gov.ab.ca/info/library/6657.pdf>.
- Ameno, K., T. Kiriu, C. Fuke, S. Ameno, T. Shinohara and I. Ijiri, 1992. Regional brain distribution of toluene in rats and in a human autopsy. *Arch. Toxicol.*, 66: 153-156.
- Angerer, J. and A. Kramer, 1997. Occupational chronic exposure to organic solvents. XVI. Ambient and biological monitoring of workers exposed to toluene. *Int. Arch. Occup. Environ. Health*, 69: 91-96.
- Angerer, J., 1979. Occupational chronic exposure to organic solvents: VII. Metabolism of toluene in man. *Int. Arch. Occupat. Environ. Health*, 43: 63-67.
- Bahrami, A.R., A. Jonidi-Jafari, B. Folladi, H. Mahjub, Q. Sadri and M.M. Zadeh, 2005. Comparison of urinary O-cresol and hippuric acid in drivers, gasoline station workers and painters exposed to toluene in West of Iran. *Pak. J. Biol. Sci.*, 8: 1001-1005.
- Barr, D.B., L.C. Wilder, S.P. Caudill, A.J. Gonzalez, L.L. Needham and J.L. Pirkle, 2005. Urinary creatinine concentrations in the U.S. population: Implications for urinary biologic monitoring measurements. *Environ. Health Perspect.*, 113: 192-200.
- Brugnone, F., M. Gobbi, K. Ayyad, C. Giuliani, M. Cerpelloni and L. Perbellini, 1995. Blood toluene as a biological index of environmental toluene exposure in the normal population and in occupationally exposure workers immediately after exposure and 16 hours later. *Int. Arch. Occup. Environ. Health*, 66: 421-425.
- Cohen, A.J., H.R. Anderson, B. Ostro, K.D. Pandey and M. Krzyanowski *et al.*, 2004. Mortality Impacts of Urban Air Pollution. In: Comparative Quantification of Health Risks: Global and Regional Burden of Disease Attributable to Selected Major Risk Factors, Ezzati, M., A.D. Lopez, A. Rodgers and C.J.L. Murray (Eds.). Vol. 2, Chapter 17, World Health Organization, Geneva, pp: 1353-1434.
- Dann, T., W. Wang and A. Etlinger, 1989. Volatile organic compounds in Canadian ambient air: A new emphasis. Pollution Measurement Division Report No. PMD 89-26, Environment Canada, Ottawa. <http://www.ijc.org/files/publications/B12.pdf>.
- Duydu, Y., S. Suzen, N. Erdem, H. Uysal and N. Vural, 1999. Validation of hippuric acid as a biomarker of toluene exposure. *Bull. Environ. Contamin. Toxicol.*, 63: 1-8.
- EPA., 1993. Ambient concentration summaries for clean air act title III hazardous air pollutants. Final Report. US Environmental Protection Agency, EPA Report No. EPA/600/R-94/090, Washington, DC.
- EPA., 2005. Toxicological review of toluene (CAS No. 108-88-3). In Support of Summary Information on the Integrated Risk Information System (IRIS). EPA/635/R-05/004. September 2005. <http://www.epa.gov/iris/toxreviews/0118tr.pdf>.

- Ezzati, M., A.D. Lopez, A. Rodgers, S. van der Hoorn, C.J. Murray and The Comparative Risk Assessment Collaborating Group, 2002. Selected major risk factors and global and regional burden of disease. *Lancet*, 360: 1347-1360.
- Hashim, J.H., Z. Hashim, M.F. Jabarudin, N. Ismail, L. Naing, M.H. Jaafar and R. Nordin, 2006. Toluene exposure and respiratory problems among spray painters in an automotive industry. *Epidemiology*, 17: S172-S173.
- Hsieh, C.C. and J.H. Tsai, 2003. VOC concentration characteristics in Southern Taiwan. *Chemosphere*, 50: 545-556.
- IARC., 1999. Toluene diisocyanates (Group 2B). International Agency for Research on Cancer (IARC)-Summaries and Evaluations, Volume 71, pp: 865.
- Kampa, M. and E. Castanas, 2008. Human health effects of air pollution. *Environ. Pollut.*, 151: 362-367.
- Kawai, T., H. Ukai, O. Inoue, Y. Maejima and Y. Fukui *et al.*, 2008. Evaluation of biomarkers of occupational exposure to toluene at low levels. *Int. Arch. Occupat. Environ. Health*, 81: 253-262.
- Kerbachi, R., M. Boughedaoui, L. Bounoua and M. Keddad, 2006. Ambient air pollution by aromatic hydrocarbons in Algiers. *Atmospheric Environ.*, 40: 3995-4003.
- Kim, J.J., J.O. Ham, K.D. Ahn, B.K. Lee, T.S. Nam and N.W. Paik, 1989. A study on urinary hippuric acid and subjective symptoms in toluene exposed workers. *Korean J. Occup. Environ. Med.*, 1: 206-217.
- Kumar, A. and S.K. Tyagi, 2006. Benzene and toluene profiles in ambient air of Delhi as determined by active sampling and GC analysis. *J. Sci. Indust. Res.*, 65: 252-257.
- Laowagul, W., H. Garivait, W. Limpaseni and K. Yoshizumi, 2008. Ambient air concentrations of benzene, toluene, ethylbenzene and xylene in Bangkok, Thailand during April-August in 2007. *Asian J. Atmospheric Environ.*, 2: 14-25.
- Lee, S.C., M.Y. Chiu, K.F. Ho, S.C. Zou and X. Wang, 2002. Volatile Organic Compounds (VOCs) in urban atmosphere of Hong Kong. *Chemosphere*, 48: 375-382.
- Ogata, M. and T. Taguchi, 1987. Quantitation of urinary metabolites of toluene, xylene, styrene, ethylbenzene and phenol by automated high performance liquid chromatography. *Int. Arch. Occupat. Environ. Health*, 59: 263-272.
- Pagnotto, L.D. and L.M. Lieberman, 1967. Urinary hippuric acid excretion as an index of toluene exposure. *Am. Indust. Hyg. Assoc. J.*, 28: 129-134.
- Slot, C., 1965. Plasma creatinine determination a new and specific Jaffe reaction method. *Scand. J. Lab. Invest.*, 17: 381-387.
- Tunsaringkam, T., J. Suwansaksri, P. Ketkaew and W. Siriwong, 2011. Blood toluene and genotoxicity in gasoline station workers in Bangkok: A preliminary study. *J. Health Res.*, 25: 161-164.
- Tunsaringkam, T., W. Siriwong, A. Rungsiyothin and S. Nopparatbundit, 2012. Occupational exposure of gasoline station workers to BTEX compounds in Bangkok, Thailand. *Int. J. Occupat. Environ. Med.*, 3: 117-125.
- US EPA., 1980. Toluene, health and environmental effects profile No. 160. US Environmental Protection Agency, Office of Solid Waste, Washington, D.C., April 30, 1980.
- US EPA., 1994. Chemicals in the environment: Toluene (Cas No. 108-88-3). OPPT Chemical Fact Sheet, EPA 749-F-94-021, Office of Pollution Prevention and Toxics, US Environmental Protection Agency, August 1994. [http://www.epa.gov/chemfact/f\\_toluene.txt](http://www.epa.gov/chemfact/f_toluene.txt).
- Ukai, H., T. Watanabe, H. Nakatsuka, T. Satoh and S.J. Liu *et al.*, 1993. Dose-dependent increase in subjective symptoms among toluene-exposed workers. *Environ. Res.*, 60: 274-289.
- WHO, 2002. The world health report 2002: Reducing risks, promoting healthy life. World Health Report, World Health Organization, Geneva, Switzerland. <http://www.who.int/whr/2002/en/>.
- WHO., 1985. Toluene. Environmental Health Criteria, No. 52, World Health Organization, Geneva. <http://www.inchem.org/documents/ehc/ehc/ehc52.htm>.
- WHO., 2000. Chapter 5.14: Toluene. WHO Regional Office for Europe, Copenhagen, Denmark. [http://www.euro.who.int/\\_data/assets/pdf\\_file/0020/123068/AQG2ndEd\\_5\\_14Toluene.PDF](http://www.euro.who.int/_data/assets/pdf_file/0020/123068/AQG2ndEd_5_14Toluene.PDF).
- WHO., 2014. Ambient (outdoor) air quality and health. Fact sheet No. 313. World Health Organization. <http://www.who.int/mediacentre/factsheets/fs313/en/>.
- Wang, F., F. Costabile, F. Liu, W. Hong, D. Fang and I. Allegrini, 2010. Ambient BTX measurements in Suzhou, China. *Environ. Monit. Assess.*, 168: 21-31.
- Wiwanitkit, V., J. Suwansaksri, S. Srita and A. Fongsoongnern, 2002. High levels of hippuric acid in urine of Thai press workers. *Southeast Asian J. Trop. Med. Public Health*, 33: 624-627.
- Yimrungruang, D., V. Cheevaporn, T. Boonphakdee, P. Watchalayann and H.F. Helander, 2008. Characterization and health risk assessment of volatile organic compounds in gas service station workers. *EnvironmentAsia*, 2: 21-29.