

Association of Vitamin D Level with Children Fractures: A Prospective Cohort Study

¹Rami Jahmani, ²Liqaa Raffee, ⁴Alaa Mutasim Babiker Ali, ³Mohammed Alorjani,

¹K.M. Bashaireh, ¹Audat Ziad and ¹Ahmad Radaideh

¹Department of Special Surgery,

²Department of Accident and Emergency Medicine,

³Department of Pathology, Faculty of Medicine,

Jordan University of Science and Technology, 22110 Irbid, Jordan

⁴Omdurman Islamic University, Khartoum-Sudan

Abstract: Fractures have increased prevalence among children which requires searching for intervening options. To evaluate the association of vitamin D with fractures in children. The study design was a prospective design. The study included 70 patients with diagnosed fracture by X-ray and 49 control subjects. Data were collected using a clinical data sheet and laboratory finding of vitamin D. Data were analyzed using SPSS Version 21. The mean age of study group was 6.57±4.66 years and that of control group was 6.88±5.47 years. Males were more frequent in both study and control groups 58.6 and 59.18%, respectively. The mean level of vitamin D in study group was 23.22±14.68 ng/mL and this was slightly higher than that of control group 22.52±12.23 ng/mL. According to anatomic site, fractures in upper limb 75.7% were more prevalent than that in lower limb 24.3%. The type of fracture was either complete 88.6% or green stick 11.4%. No significant association was obtained of vitamin D with fractures in children ($p>0.05$). Among patients who have fractures, a significant association between vitamin D level and each of age, anatomic location of structure and type of fracture. The present study showed, the lack of association between vitamin D level and fractures in children.

Key words: Children, vitamin D, fracture, type of fracture, anatomic, site

INTRODUCTION

An emerging trend has been witnessed in the middle of the 20th century, so that, the mortality of infants and children resulting from infections and nutritional disturbances has dramatically decreased in Europe. On the other hand the role of accidents in death rate and morbidity has become more important (Mayranpaa, 2012).

Several studies have indicated that fractures are prevalent among children accounting for 10-25% of all pediatric injuries (Sibert *et al.*, 1981; Landin, 1983; Jones and Cooley, 2002; Mattila *et al.*, 2004; Spady *et al.*, 2007).

The term "vitamin D" was first used in 1922 when McCollum and colleagues reported the identification of a new lipid-soluble substance which they named vitamin D. This finding appeared to be of great importance as the new vitamin showed to be capable of

preventing rickets, a bone disease which at that time had reached epidemic proportion (Hansen *et al.*, 2001; Holick, 2006).

The identification of vitamin D led to intensive search for the active substrate of this vitamin in 1931. Askew and colleagues succeeded in isolating vitamin D₂ from irradiated plant sterols, later in 1936, vitamin D₃ was identified by Windaus and colleagues but it was not until the 1971 where (1,25 (OH)₂ D₃) the metabolically active form of vitamin D₃ was identified by Holick and colleagues (Holick, 2003; Kochupillai, 2008). Vitamin D regulates colonic proliferation, differentiation, apoptosis and angiogenesis and the colonic vitamin D system is regulated by several known natural factors, the most important one is nutritional calcium (Cross *et al.*, 2009).

Cranney *et al.* (2007) conducted a study to review and analyze the literature about the association between vitamin D and bone fracture. The results showed a

deficiency in studies about infants and recommended conducting additional high quality studies in infants, children, premenopausal women and diverse racial or ethnic groups. However, the evidence for an association was inconsistent for fractures in infants.

Contreras *et al.* (2014) conducted a study to evaluate the potential of healthy children who had fractures due to accidental trauma to be deficient of vitamin D compared to their counterparts without fractures. Findings of the study showed that the level of vitamin D in fracture group was 26.7 ng/mL and that of control group was 25.45 ng/mL. No statistical variations were obtained ($p = 0.84$).

Minkowitz *et al.* (2017) conducted a study to explore the possibility of existence of association between the level of vitamin D and life style of children with fractures and those without fractures. Results showed that both fracture and non-fracture patients and similar levels of vitamin D 27.5 ± 8.9 - 27.4 ± 9.1 ng/mL, respectively. No significant association was observed ($p = 0.941$).

Study objectives: The aim of this study was to evaluate the association of vitamin D level with fractures in children.

MATERIALS AND METHODS

- Study design: a single-center, prospective study design was conducted
- Study population: children from 0-16 years who admitted or visit OPD clinic at King Abdullah University Hospital with fracture confirmed by X-ray
- Study sample: study sample included 70 patients with diagnosed fracture and 49 control persons
- Study area: King Abdullah University Hospital, Jordan University of Science and Technology
- Study period: January 2017-March 2017
- Study tool: clinical data sheet with vitamin D level assessment
- Inclusion criteria: any child with fracture under the age of 18 years and his caregiver signed a consent form to carry out the study
- Exclusion criteria: participant not fulfilling the inclusion criteria

Data analysis: Data were collected in excel sheet to create raw data and then the data were exported into SPSS Version 21 for statistical analysis. Data were analyzed based on frequency and percentages for categorical variables such as gender, Means (M) and Standard Deviation (SD) for continuous variables such as age. The association between vitamin D level among study and control groups was computed based on independent t-test. Significance was considered at $p < 0.05$.

RESULTS AND DISCUSSION

Demographic characteristics of participants: Table 1 and Fig. 1 and 2 shows that the mean age of study group was 6.57 ± 4.66 years and that of control group was 6.88 ± 5.47 years. Males were more frequent in both study and control groups 58.6 and 59.18%, respectively.

Clinical characteristics of study participants: Table 2 shows the mean level of vitamin D in study group was 23.22 ± 14.68 ng/mL and this was slightly higher than that of control group 22.52 ± 12.23 ng/mL (Fig. 3). According to anatomic site, fractures in upper limb 75.7% were more prevalent than that in lower limb 24.3% (Fig. 4). The type of fracture was either complete 88.6% or green stick 11.4% (Fig. 5).

Table 1: Demographic characteristics of participants

| Variables | Description |
|------------------------------------|-------------|
| Age-study group (M±SD) years | 6.57±4.66 |
| Age-control group (M±SD) years | 6.88±5.47 |
| Gender-study group (N, %) | |
| Males | 41 (58.6) |
| Females | 29 (41.4) |
| Gender-control group (N, %) | |
| Males | 29 (59.18) |
| Females | 20 (40.82) |




Fig. 1: The frequency of study participants by gender




Fig. 2: The frequency of study participants by age

Table 2: Clinical characteristics of study participants

| Variables | Description |
|---|-------------|
| Vitamin D-study group (MSD), ng/mL | 23.22±14.68 |
| Vitamin D-control group (M±SD), ng/mL | 22.52±12.23 |
| Fracture by anatomic region (N, %) | |
| Upper limb | 53 (75.7) |
| Lower limb | 17 (24.3) |
| Type of fracture (N, %) | |
| Complete | 62 (88.6) |
| Green stick | 8 (11.4) |




Fig. 3: Prevalence of vitamin D in study groups




Fig. 4: Fracture by anatomic region

The association between vitamin D levels in study and control groups: Table 3 shows, no significant differences between the level of vitamin D in study group 23.22±14.26 ng/mL and control group 22.52±12.23 ng/mL $p = 0.775$.

The association between vitamin D and study variables (one way ANOVA): The level of vitamin D is subdivided into two main categories: ≤ 30 and > 30 ng/mL. Table 4 shows, the level of vitamin D is associated significantly with age $p = 0.006$ and anatomic location of fracture $p = 0.000$ and type of fracture $p = 0.000$.

Table 3: The association between vitamin D levels in study and control groups (independent t-test)

| Pairs | M | SD | p-value |
|-------------------------|-------|-------|---------|
| Vitamin D-study group | 23.22 | 14.26 | 0.775 |
| Vitamin D-control group | 22.52 | 12.23 | - |




Fig. 5: Frequency of fractures according to their type

This study was conducted to investigate the association of vitamin D level with fractures in children. The data of this study showed that there was no association between vitamin D and fracture in children ($p = 0.775$). This finding is in line with other studies that reported no association between vitamin D and fracture (Contreras *et al.*, 2014; Minkowitz *et al.*, 2017).

The efforts in this study failed to prove this association, probably due to have similar levels of vitamin D in both groups. Although, the general trend in Jordan is to find population deficient in vitamin D as it is the general status in this study it seems that vitamin D deficiency may need to be less than in this study to impact fractures. According to the study by Cranney *et al.* (2007) there is a deficiency in studies about infants and recommended conducting additional high quality studies in infants, children, premenopausal women and diverse racial or ethnic groups. However, the evidence for an association was inconsistent for fractures in infants.

The findings of this study showed that among patients with fractures there was a strong association between vitamin D and each of age $p = 0.006$ anatomic location $p = 0.000$ and type of fracture $p = 0.000$. These findings may add new piece of information for literature because these findings are not well addressed in literature. It is plausible to claim a role of vitamin D in severity of fractures due to the effect of vitamin D in bone structures as reported in several studies (Cranney *et al.*, 2007; Contreras *et al.*, 2014; Minkowitz *et al.*, 2017).

Table 4: The association between vitamin D and study variables (one way ANOVA)

| Variables | Sum of squares | df | Mean square | F-values | Sig. |
|------------------------------------|----------------|----|-------------|----------|-------|
| Age | | | | | |
| Between groups | 1.851 | 1 | 1.851 | 8.112 | 0.006 |
| Within groups | 15.520 | 68 | 0.228 | | |
| Total | 17.371 | 69 | | | |
| Vitamin D-anatomic location | | | | | |
| Between groups | 8785.506 | 1 | 8785.506 | 113.579 | 0.000 |
| Within groups | 5259.925 | 68 | 77.352 | | |
| Total | 14045.431 | 69 | | | |
| Vitamin D-type of fracture | | | | | |
| Between groups | 8785.506 | 1 | 8785.506 | 113.579 | 0.000 |
| Within groups | 5259.925 | 68 | 77.352 | | |
| Total | 14045.431 | 69 | | | |

CONCLUSION

The present study showed the lack of association between vitamin D level and fractures in children. Within patients who have fractures, a significant association between vitamin D level and each of age, anatomic location of structure and type of fracture.

REFERENCES

- Contreras, J.J., B. Hiestand, J.C. O'Neill, R. Schwartz and M. Nadkarni, 2014. Vitamin D deficiency in children with fractures. *Pediatr. Emergency Care*, 30: 777-781.
- Cranney, A., T. Horsley, S. O'Donnell, H. Weiler and L. Puil *et al.*, 2007. Effectiveness and Safety of Vitamin D in Relation to Bone Health. U.S. Department of Health and Human Services, USA., ISBN: 9781587632655, Pages: 235.
- Cross, H.S., T. Nittke and M. Peterlik, 2009. Modulation of vitamin D synthesis and catabolism in colorectal mucosa: A new target for cancer prevention. *Anticancer Res.*, 29: 3705-3712.
- Hansen, C.M., L. Binderup, K.J. Hamberg and C. Carlberg, 2001. Vitamin D and cancer: Effects of 1, 25 (OH) 2D3 and its analogs on growth control and tumorigenesis. *Front. Biosci.*, 6: 820-848.
- Holick, M.F., 2003. Vitamin D: A millenium perspective. *J. Cell. Biochem.*, 88: 296-307.
- Holick, M.F., 2006. Resurrection of vitamin D deficiency and rickets. *J. Clin. Invest.*, 116: 2062-2072.
- Jones, G. and H.M. Cooley, 2002. Symptomatic fracture incidence in those under 50 years of age in Southern Tasmania. *J. Paediatrics Child Health*, 38: 278-283.
- Kochupillai, N., 2008. The physiology of vitamin D: Current concepts. *Indian J. Med. Res.*, 127: 256-262.
- Landin, L.A., 1983. Fracture patterns in children: Analysis of 8,682 fractures with special reference to incidence, etiology and secular changes in a Swedish urban population 1950-1979. *Acta Orthopaedica Scand*, 54: 3-109.
- Mattila, V., J. Parkkari, P. Kannus and A. Rimpela, 2004. Occurrence and risk factors of unintentional injuries among 12-to 18-year-old Finns: A survey of 8219 adolescents. *Eur. J. Epidemiol.*, 19: 437-444.
- Mayranpaa, M., 2012. Fractures in children: Epidemiology and associated bone health characteristics. Master Thesis, University of Helsinki, Helsinki, Finland.
- Minkowitz, B., B. Cerase, E. Poletick, J.T. Nguyen and N.D. Formoso *et al.*, 2017. Low vitamin D levels are associated with need for surgical correction of pediatric fractures. *J. Pediatr. Orthopaedics*, 37: 23-29.
- Sibert, J.R., G.B. Maddocks and B.M. Brown, 1981. Childhood accidents: An endemic of epidemic proportion. *Arch. Dis. Childhood*, 56: 225-227.
- Spady, D.W., D.L. Saunders, D.P. Schopflocher and L.W. Svenson, 2007. Patterns of injury in children: A population-based approach. *Pediatr.*, 113: 522-529.