

Incidence of Fetal Wastage in Cattle Slaughtered at the Oko-Oba Abattoir and Lairage, Agege, Lagos, Nigeria

A. Ibironke Ademola

¹Department of Paraclinical Studies, Faculty of Veterinary Science,
University of Pretoria, South Africa

Abstract: The aim of this study is to determine the proportion of fetuses destroyed due to the slaughter of pregnant cows at the Oko-Oba abattoir and lairage, Agege, Lagos, Nigeria. A retrospective study was conducted on abattoir records kept by the Lagos State Department of Veterinary Services (LSDVS) over a 4 year period. The study showed an incidence rate ranging from 1.50-2.10% over the 4 year study period (Average of 1.8%). This practice constitutes a huge drain on the animal protein availability for human consumption in a developing country like Nigeria. It is therefore recommended that appropriate legislation be put in place and enforced to control the slaughter of pregnant animals. Comprehensive ante-mortem inspection should also be carried out by abattoir staff on all animals intended for slaughter for human consumption.

Key words: Fetal wastage, cattle, abattoir, Nigeria, incidence, consitutes

INTRODUCTION

The United Nations in 2003 estimated Nigeria's population to be 124,009,000 making Nigeria the most populous country in Africa and one of the top ten most populous countries in the world. As a result of this huge population, there is a high demand for animal protein to meet the dietary protein requirements of the populace. Meeting this high protein requirement is therefore a great challenge for a developing country like Nigeria.

Meat from cattle slaughtered at the various abattoirs in the country, constitute the largest source of animal protein for the Nigerian populace (Idahor *et al.*, 2009b). There is however, a disturbing trend in the bid to provide meat for the consumption of the human populace which involves the slaughter of pregnant female animals. The slaughter of pregnant animals for meat is unethical (Khan and Khan, 1989).

This practice frustrates the efforts of geneticist, nutritionist and livestock breeders and is a drain on breeding animals thus widening the gap of animal protein between the ever increasing human populations (Khan and Khan, 1989).

These fetuses are discovered during routine post-mortem meat inspection and are totally condemned by the meat inspection officers at the abattoir. Previous researches have shown the occurrence of slaughter of pregnant animals in different countries of the world and

these reports have shown some very disturbing figures. This trend poses a threat to efforts to meet the dietary protein requirements of many developing countries like Nigeria. A study conducted at the Faisalabad abattoir revealed 11.65, 8.61, 21.28 and 19.22% in buffaloes, cattle, sheep and goats, respectively for frequency of pregnant animals slaughtered (Khan and Khan, 1989). A study by Tchoumboue (1984) reported that 16% of slaughtered cattle in Yaoundé abattoir in Cameroon were pregnant while Ndi *et al.* (1993) reported an average of 22.1% female cattle pregnant in Bamenda and as much as 45% in Yaounde.

A similar situation was also reported by Al-Dahash and David, 1977. Occurrence of fetal wastage has also been reported in various abattoirs in different parts of Nigeria. A 2.6% rate of slaughter of pregnant cattle was reported in Enugu (Wosu, 1988) while 0.32% of pregnant cattle were reportedly slaughtered at the Doma abattoir (Idahor *et al.*, 2009a).

Idahor *et al.*, (2009b) also reported 4.6, 10 and 29.5% for cows, does and ewes, respectively at the Lafia abattoir. A similar trend has also been reported in Gombe state (Mohammad *et al.*, 2008), Sokoto state (Maigandi *et al.*, 2008; Mohammad *et al.*, 2007), Bauchi and Jos (Sanusi *et al.*, 2006) while Abdulkadir *et al.* (2008) reported an incidence rate of 3.9% for slaughter of pregnant animals in Makurdi abattoir, Benue state from 1997-2002.

There was however, no published report on fetal wastage from the Oko-Oba abattoir in Lagos which is one of the largest and best organized urban abattoir in one of the most populous cities in West Africa. The Oko-Oba abattoir receives and processes the highest number of cattle to beef in Nigeria hence its importance.

The aim of this study is therefore to determine the rate of fetal losses encountered during routine post-mortem meat inspection at the Oko-Oba abattoir and Lairage, Agege, Lagos, Nigeria and some possible implications of these losses.

MATERIALS AND METHODS

Study area and animals: The Oko-Oba abattoir and lairage is located at Agege (longitude 3°17'01" and latitude 6°39'32), a suburb of Lagos state in the western part of Nigeria. It is one of the largest and best organized abattoirs in Nigeria and receives cattle from various parts of Nigeria, mainly northern Nigeria and even from the countries in the West Africa sub-region including Niger, Chad, Burkina Faso, Mali and Cameroon (Cadmus *et al.*, 2006; Ibiroko *et al.*, 2010).

Although the abattoir has the daily maximum handling capacity of more than one thousand and three hundred heads of cattle, it presently operate with the slaughter of an average of around 1000 heads of cattle daily (Ibiroko *et al.*, 2010).

Since most abattoir slaughters are done early in the morning in Nigeria due to lack of good storage facilities and good preservation systems, the abattoir appears congested in the mornings, a situation that makes it seem incapable of handling the number currently slaughtered on a daily basis.

The abattoir receives animals from wide geographical areas of West Africa and it provides meat to a cosmopolitan population of the city of Lagos, the choice of the abattoir as a point for data collection is therefore considered representative for monitoring animal disease incidences and patterns that present in an average Nigerian abattoir.

Data collection: The retrospective data covering periods from 2004-2008 were collected and analyzed based on abattoir records available through the effort of the meat inspectors and veterinary officers of the Lagos State Department of Veterinary Services (LSDVS).

Data of period prior to August 2004 were excluded due to inaccuracy and inconsistency. The LSDVS kept effective and accurate records from 2004 onward following

a government directive to commence the monitoring of key zoonotic and infectious diseases at the abattoir. All of the records of abattoir slaughtering of cattle, carcass and offal condemnation were obtained for a period of 3 years (August, 2004-August, 2007). Daily visits were made to the abattoir by the researchers between 15th November, 2007 and 15th January, 2008 to ascertain the current status and scope of the problems on the slaughter slabs. Oral interviews were held with butchers, cattle traders and meat inspection officers at the abattoir.

RESULTS AND DISCUSSION

A total of 1,170,492 cattle were slaughtered at this abattoir over the survey period with the monthly slaughter figure ranging between 21,875 and 30,070. Of the 1,170,492 cattle slaughtered at the abattoir, 392,062 were cows (females) and the rest were bulls (males). Over the survey period, the rate of slaughter of pregnant cattle ranged between 1.50 and 2.10% (average rate of 1.8%) with a total of 7,406 fetuses condemned from pregnant females in the abattoir.

Table 1 shows the slaughter figures, number of fetuses condemned and the percentages. The result of this study confirms that pregnant cows are frequently slaughtered at the Oko-Oba abattoir. This study revealed an incidence rate ranging from 1.5-2.1% (average of 1.8%) over the 4 years studied. The result of this study is lower than results obtained from previous similar studies (Tchoumboue, 1984; Wosu, 1988; Ndi *et al.*, 1993; Abdulkadir *et al.*, 2008; Idahor *et al.*, 2009a, b).

Visits to the abattoir showed that meat inspection staff available on the slaughter slabs seemed to be overwhelmed by the number and rate of slaughter carried out at the abattoir especially during the mid-morning rush period when meat inspection was observed to be carried out haphazardly due to staff inadequacy.

As a result of this, some fetuses may be missed and smuggled away from the abattoir without being detected. This may partly explain the low figures recorded for pregnancy in slaughtered cows at the abattoir.

Under conditions on anonymity, some butchers also confessed to the researchers that they smuggled undetected fetuses outside the abattoir and sell this to dog breeders and owners who use it to supplement their dog's feed and pay varying sums of money for the fetuses; a practice that should be discouraged as it can spread diseases like brucellosis to both the dog and humans. This may also explain why recorded figures are lower than expected.

Table 1: Incidence of fetal wastage at the Oke-Oba abattoir

Periods	Total cattle slaughtered	Total cows (females) slaughtered	Percentage of cows (Females) slaughtered	No. of fetuses condemned	Percentage of fetuses condemned to number of cows
Aug. 2004-Dec. 2004	156,953	51,813	33.01	778	1.50
Jan. 2005-Dec. 2005	381,855	128,025	33.53	2,692	2.10
Jan. 2006- Dec. 2006	402,139	133,719	33.25	2,592	1.94
Jan. 2007- Aug. 2007	229,545	78,505	34.20	1,344	1.71
Total	1,170,492	392,062	-	7,406	Average = 1.8%

CONCLUSION

Slaughtering of pregnant animals for meat purpose is unethical and is contrary to the rules of slaughter under which only unproductive, infertile, sterile, old or accidentally injured animals are allowed to slaughter (Khan and Khan, 1989). It also frustrates the scientific endeavors of geneticist, nutritionist and livestock breeders working for the propagation of animal species (Khan and Khan, 1989).

This practice also ultimately reduces the quantity and quality of animal protein available for human consumption which worsens an already bad situation in most developing countries. The practice of slaughter of pregnant animals should therefore be discouraged as it is even cruel to the animals and contrary to the principles of animal welfare.

RECOMMENDATIONS

The following recommendations are however made to stem the tide:

- Legislation on the prohibition of slaughter of pregnant animals should be passed into law and enforced by the law enforcement agents. Stiff penalties should be stipulated for non-compliance
- There is need for more meat inspection staff at the abattoirs to meet the rate of slaughter and carry out more comprehensive meat inspection
- Ante-mortem inspection should be comprehensively carried out on animal before they are passed as fit for slaughter. This process will help detect pregnant cow and reduce the current level of wastage.
- Butchers and the general populace should be educated on the implications of slaughtering pregnant animals or handling animal fetuses
- In more advanced countries of the world, serum progesterone levels of animals meant for slaughter can be measured to detect pregnant animals prior to slaughter
- Endoscopy can also be employed in sheep and goat to detect pregnancy in the more developed countries of the world (Khan and Khan, 1989)

ACKNOWLEDGEMENTS

The researchers wishes to extend his gratitude to Dr (Mrs). Taiwo Adejuwon and Dr. M.O. Adeyemi of the Lagos State Department of Veterinary Services and the staff of Harmony Butcheries for their cooperation during the period of this study.

REFERENCES

- Abdulkadir, U., E.Z. Jiya and S.A. Kosu, 2008. Survey of fetal wastages: A case study of makurdi abattoir in benue state from 1997 to 2002. Pak. J. Nutr., 7: 450-452.
- Al-Dahash, S.Y.A. and J.S.E. David, 1977. The incidence of ovarian activity, pregnancy and bovine genital abnormalities shown by an abattoir survey. Vet. Record., 101: 296-299.
- Cadmus, S.I.B., I.F. Ijagbone, H.E. Oputa, H.K. Adesokan and J.K. Stack, 2006. Serological survey of brucellosis in livestock animals and workers in Ibadan, Nigeria. Afr. J. Biomed. Res., 9: 163-168.
- Ibironke, A.A., C.M.E. McCrindle, T.A. Adejuwon and S.I.B. Cadmus, 2010. Losses associated with mortality of cattle and camels during transport to Oke-Oba abattoir, Lagos state, Nigeria. Eur. J. Trans. Myol. Basic Applied Myol., 1: 13-16.
- Idahor, K.O., J.N. Omeje and A. Akwe, 2009a. Embryonic losses from slaughtering of pregnant ruminant animals at Doma abattoir. Proceedings of the 14th Annual Conference of Animal Science Association of Nigeria, Sept. 14-17, LAUTECH Ogbomoso, Nigeria, pp: 240-242.
- Idahor, K.O., J.N. Omeje, V.E. Agu, P. Audi, S.R. David and B.D. Luka, 2009b. Awareness of fetal losses from ruminants slaughtered at Lafia abattoir. J. Life Phys. Sci., 3: 44-48.
- Khan, M.Z. and A. Khan, 1989. Frequency of pregnant animals slaughtered at Faisalabad abattoir. J. Islamic Acad. Sci., 2: 82-82.
- Maigandi, S.A., F. Bibi-Farouk and I.R. Mohammed, 2008. Seasonal fetal recovery in Camels (*Camelus dromedarius*) slaughtered in Sokoto state abattoir in semi-arid zone of Nigeria. Proceedings of the 13th Annual Conference of Animal Science, (ACAS'08), ABU, Zaria, pp: 208-211.

- Mohammad, B.F., I.Y. Haruna, A.M. Abdulsamad and J.M. Bichi, 2008. Fetal wastage in Northern Nigeria: The case of Gombe abattoir, Gombe State. Proceedings of the 13th Annual Conference of Animal Science, (ACAS'08), ABU, Zaria, pp: 124-127.
- Mohammad, I.R., R. Ashiru and A.Y. Abdulahi, 2007. Implications of slaughter of pregnant ewes and does to future stock in the semi-arid urban abattoir. *J. Anim. Vet. Adv.*, 6: 819-822.
- Ndi, C., N.E. Tambi and N.W. Aghari, 1993. Reducing calf wastage from slaughter of pregnant cows in Cameroon. *World Anim. Rev.*, http://www.fao.org/documents/art_dett.asp?lang=en&art_id=54949.
- Sanusi, M., M. Abubakar and B. Luka, 2006. Incidence of fetal wastage in ruminant animals slaughtered at Bauchi and Jos abattoirs. Proceedings of the 31st Annual of the Nigerian Society of Animal Production, March 12-15, Bayero University, Kano, Nigeria, pp: 102-106.
- Tchoumboue, J., 1984. Calves lost through pregnant cow slaughtering: A particular case in Yaoundé abattoir, Cameroon. *Revue Elevage Medecine Veterinaire Pays Tropicaux*, 37: 70-72.
- Wosu, L.D., 1988. Calf wastage through slaughtering of pregnant cows in Enugu abattoir, Nigeria. *Revue Elevage Medecine Veterinaire Pays Tropicaux*, 41: 97-98.